



Medical Policy

Chimeric Antigen Receptor Therapy for Hematologic Malignancies

Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

Policy Number: 066

BCBSA Reference Number: 8.01.63

Related Policies

- Adoptive Immunotherapy, #[455](#)
- CAR T-Cell Therapy Services for Diffuse Large B-cell Lymphoma (axicabtagene cilleucel or tisagenlecleucel) Prior Authorization Request Form, #[924](#)
- CAR T-Cell Therapy Services for B-cell Acute Lymphoblastic Leukemia (tisagenlecleucel) Prior Authorization Request Form, #[925](#)
- CAR T-Cell Therapy Services for Mantle Cell Lymphoma (Brexucabtagene Autoleucel) Prior Authorization Request Form, #[940](#)
- Cellular Immunotherapy for Prostate Cancer, #[268](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Prior Authorization Request Form: Chimeric Antigen Receptor Therapy for Hematologic Malignancies

This form must be completed and faxed to: Medical and Surgical: 888-973-0726; Medicare Advantage: 1-800-447-2994.

- Click here for CAR T-Cell Therapy Services for **Diffuse Large B-cell Lymphoma** (axicabtagene cilleucel or tisagenlecleucel) Prior Authorization Request Form #[924](#)
- Click here for CAR T-Cell Therapy Services for **B-cell Acute Lymphoblastic Leukemia** (tisagenlecleucel) Prior Authorization Request Form #[925](#)
- Click here for CAR T-Cell Therapy Services for **Mantle Cell Lymphoma** (Brexucabtagene Autoleucel) Prior Authorization Request Form #[940](#)

Tisagenlecleucel (Kymriah): B-cell acute lymphoblastic leukemia

Tisagenlecleucel may be considered **MEDICALLY NECESSARY** for relapsed^a (second or later) or refractory^b patients if they meet **ALL** of following criteria 1 through 5:

1. Confirmed diagnosis of CD19-positive B-cell acute lymphoblastic leukemia with morphologic marrow tumor involvement ($\geq 5\%$ lymphoblasts)
2. Are up to 25 years old at the time of infusion
3. Have not received prior treatment with tisagenlecleucel or any other gene therapy or are being considered for treatment with any other gene therapy
4. Have adequate organ function with no significant deterioration in organ function expected within 4 weeks after apheresis
5. Do not have any of the following:
 - Burkitt lymphoma
 - Active hepatitis B, C, or any uncontrolled infection
 - Grade 2 to 4 graft-versus-host disease
 - Concomitant genetic syndrome with the exception of Down syndrome
 - Received allogeneic cellular therapy, such as donor lymphocyte infusion, within 6 weeks prior to tisagenlecleucel infusion
 - Patient has active central nervous system 3 acute lymphoblastic leukemia (ie, white blood cell count ≥ 5 cells/ μ L in cerebrospinal fluid with presence of lymphoblasts).^a
 - The healthcare facility that dispenses and administers Kymriah is enrolled and complies with the Risk Evaluation and Mitigation Strategy known as Kymriah REMS, including:
 - Onsite, immediate access to tocilizumab
 - Availability of a minimum of two doses of tocilizumab for each patient for administration within 2 hours after Kymriah infusion, if needed for treatment of cytokine release syndrome
 - Assurance that healthcare providers who prescribe, dispense or administer Kymriah are trained in the management of cytokine release syndrome and neurologic toxicities.

^a Relapsed disease describes the reappearance of leukemia cells in the bone marrow or peripheral blood after the attainment of a complete remission with chemotherapy and/or allogeneic cell transplant.

^b Refractory (resistant) disease is defined as those patients who fail to obtain complete response with induction therapy, ie, failure to eradicate all detectable leukemia cells ($<5\%$ blasts) from the bone marrow and blood with subsequent restoration of normal hematopoiesis ($>25\%$ marrow cellularity and normal peripheral blood counts).

*Central nervous system (CNS) disease for B-cell acute lymphoblastic leukemia is defined by the following groups:

- CNS 1: Absence of blasts on cerebrospinal fluid cytospin preparation, regardless of the white blood cell (WBC) count
- CNS 2: WBC count of less than 5/mL and blasts on cytospin findings
- CNS 3: WBC count of 5/mL or more and blasts on cytospin findings and/or clinical signs of CNS leukemia (eg, facial nerve palsy, brain/eye involvement, hypothalamic syndrome).

Tisagenlecleucel (Kymriah): Non-Hodgkin lymphoma

Tisagenlecleucel may be considered **MEDICALLY NECESSARY** for relapsed^c or refractory^c patients with aggressive types of non-Hodgkin lymphoma if they meet 1 through 6:

1. Are adults (age ≥ 18) at the time of infusion
2. Histologically confirmed diagnosis of diffuse large B-cell lymphoma, not otherwise specified; high-grade B-cell lymphoma **or** diffuse large B-cell lymphoma arising from follicular lymphoma.
3. Received adequate prior therapy including **ALL** of the following:
 - Anti-CD20 monoclonal antibody for CD20-positive tumor
 - Anthracycline-containing chemotherapy regimen
 - For subjects with transformed follicular lymphoma, prior chemotherapy for follicular lymphoma and subsequently have chemorefractory disease after transformation to diffuse large B-cell lymphoma:

- If patient has a history of allogeneic stem cell transplant, has no signs of active graft versus host disease
 - No active autoimmune disease requiring systemic immunosuppression
4. Have adequate organ and bone marrow function as determined by the treating oncologist/hematologist
 5. Have not received prior CD19-directed CAR T-cell therapy treatment or any other gene therapy or are being considered for treatment with any other gene therapy
 6. Do not have primary central nervous system lymphoma

AND

The healthcare facility that dispenses and administers Yescarta is enrolled and complies with the Risk Evaluation and Mitigation Strategy including:

- Onsite, immediate access to tocilizumab, **AND**
- Availability of a minimum of two doses of tocilizumab for each patient for administration within 2 hours after Yescarta infusion, if needed for treatment of cytokine release syndrome, **AND**
- Assurance that healthcare providers who prescribe, dispense or administer Yescarta are trained in the management of cytokine release syndrome and neurologic toxicities.

Tisagenlecleucel is considered **INVESTIGATIONAL** for the treatment of relapsed or refractory primary mediastinal large B-cell lymphoma.

^cRelapsed or refractory disease, defined as progression after 2 or more lines of systemic therapy (which may or may not include therapy supported by autologous cell transplant).

Axicabtagene ciloleucel (Yescarta): Non-Hodgkin lymphoma

Axicabtagene ciloleucel infusion is considered **MEDICALLY NECESSARY** for relapsed or refractory^c patients with aggressive types of non-Hodgkin lymphoma if they meet 1 through 6:

1. Are adults (age ≥18) at the time of infusion
2. Histologically confirmed diagnosis of diffuse large B-cell lymphoma, not otherwise specified; **or** primary mediastinal large B-cell lymphoma **or** high-grade B-cell lymphoma **or** diffuse large B-cell lymphoma arising from follicular lymphoma.
3. Received adequate prior therapy including all of the following:
 - Anti-CD20 monoclonal antibody for CD20-positive tumor
 - Anthracycline-containing chemotherapy regimen
 - For subjects with transformed follicular lymphoma, prior chemotherapy for follicular lymphoma and subsequently have chemorefractory disease after transformation to diffuse large B-cell lymphoma
4. Have adequate organ and bone marrow function as determined by the treating oncologist/hematologist
5. Have not received prior CD19-directed CAR T-cell therapy treatment or any other gene therapy or are being considered for treatment with any other gene therapy **AND**
6. Do not have primary central nervous system lymphoma.

^c Relapsed or refractory disease is defined as progression after 2 or more lines of systemic therapy (which may or may not include therapy supported by autologous cell transplant).

Brexucabtagene Autoleucel: Mantle Cell Lymphoma

Brexucabtagene autoleucel is considered **MEDICALLY NECESSARY** for relapsed or refractory mantle cell lymphoma^d if they meet criteria 1 through 5:

1. Are adults (age ≥18) at the time of infusion, **AND**
2. Histologically confirmed diagnosis of mantle cell lymphoma, **AND**
3. Received adequate prior therapy including chemotherapy and an anti-CD20 antibody, **OR** a Bruton tyrosine kinase inhibitor (example ibrutinib or acalabrutinib), **AND**

4. Have adequate organ and bone marrow function as determined by the treating oncologist/hematologist, **AND**
5. Have not received prior FDA approved CD19-directed CAR T-cell therapy treatment or any other gene therapy or are being considered for treatment with any other gene therapy.

^d Relapsed or refractory disease is defined as disease progression after last regimen or failure to achieve a partial remission or complete remission to the last regimen

All CAR-T therapies are considered **INVESTIGATIONAL** for all other indications.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is required .*
Commercial PPO and Indemnity	Prior authorization is required .*

*Prior Authorization Request Form: Chimeric Antigen Receptor Therapy for Hematologic Malignancies

This form **must** be completed and faxed to: Medical and Surgical: 888-973-0726; Medicare Advantage: 1-800-447-2994.

- [Click here for CAR T-Cell Therapy Services for the Treatment of Diffuse Large B-cell Lymphoma](#) (axicabtagene cilteucel or tisagenlecleucel) Prior Authorization Request Form #924
- [Click here for CAR T-Cell Therapy Services for B-cell Acute Lymphoblastic Leukemia](#) (tisagenlecleucel) Prior Authorization Request Form #925
- [Click here for CAR T-Cell Therapy Services for Mantle Cell Lymphoma](#) (Brexucabtagene Autoleucel) Prior Authorization Request Form #940

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above **medical necessity criteria MUST** be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

HCPCS Codes

HCPCS codes:	Code Description
C9399	Unclassified drugs or biologicals
J3490	Unclassified drugs
J3590	Unclassified biologics
J9999	Not otherwise classified, antineoplastic drugs

Q2041	Axicabtagene ciloleucel, up to 200 million autologous anti-cd19 car positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose
Q2042	Tisagenlecleucel, up to 600 million car-positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose
Q2053	Brexucabtagene autoleucel, up to 200 million autologous anti-cd19 car positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose

ICD-10 Procedure Codes

ICD-10-PCS procedure codes:	Code Description
XW033C3	Introduction of Engineered Autologous Chimeric Antigen Receptor T-cell Immunotherapy into Peripheral Vein, Percutaneous Approach, New Technology Group 3
XW043C3	Introduction of Engineered Autologous Chimeric Antigen Receptor T-cell Immunotherapy into Central Vein, Percutaneous Approach, New Technology Group 3
XW23346	Transfusion of Brexucabtagene Autoleucel Immunotherapy into Peripheral Vein, Percutaneous Approach, New Technology Group 6

The following CPT codes are considered investigational for **Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:**

CPT Codes

CPT codes:	Code Description
0537T	Chimeric antigen receptor T-cell (CAR-T) therapy; harvesting of blood-derived T lymphocytes for development of genetically modified autologous CAR-T cells, per day
0538T	Chimeric antigen receptor T-cell (CAR-T) therapy; preparation of blood-derived T lymphocytes for transportation (eg, cryopreservation, storage)
0539T	Chimeric antigen receptor T-cell (CAR-T) therapy; receipt and preparation of CAR-T cells for administration
0540T	Chimeric antigen receptor T-cell (CAR-T) therapy; CAR-T cell administration, autologous

Description

B-cell ALL is a malignancy (clonal) of the bone marrow in which the early lymphoid precursors of the white blood cells (called lymphoblasts) proliferate and replace the normal hematopoietic cells of the marrow. This results in overcrowding of the bone marrow, as well as the peripheral organs (particularly the liver, spleen, and lymph nodes) by the lymphoblasts. As a consequence, the leukemic blasts displace the normal hematopoietic bone marrow and cause cytopenias in all 3 cell lineages (anemia, thrombocytopenia, granulocytopenia). Leukostasis affecting brain and lung may also occur. Death occurs commonly due to severe pancytopenia and resulting infections. Refractory (resistant) disease is defined as those patients who fail to obtain a complete response with induction therapy, ie, failure to eradicate all detectable leukemia cells (<5% blasts) from the bone marrow and blood with subsequent restoration of normal hematopoiesis (>25% marrow cellularity and normal peripheral blood counts). Relapsed disease describes the reappearance of leukemia cells in the bone marrow or peripheral blood after the attainment of complete remission. Minimal residual disease (MRD) refers to the presence of disease in cases deemed to be in complete remission by conventional pathologic analysis. MRD positivity is defined as the presence of 0.01% or more ALL cells and has been shown to be a strongest prognostic factor to predict the risk of relapse and death when measured during and after induction therapy in both newly diagnosed

and relapsed ALL. In a meta-analysis of 20 studies of 11249 pediatric ALL, Berry et al (2017) reported a hazard ratio for event-free survival in MRD-negative patients compared with MRD-positive patients of 0.23 (95% confidence interval, 0.18 to 0.28).¹

Approximately 5000 cases of B-cell ALL are diagnosed every year in the United States,² and approximately 620 pediatric and young adult patients with B-cell ALL will relapse each year in the United States.³ B-cell ALL is largely a disease of the young, with approximately 60% of cases occurring in patients younger than 20 years old with a median age at diagnosis of 15 years.²

Treatment

While treatable in 85% cases, approximately 15% of children and young adults with ALL will relapse and 2% to 3% of ALL patients are primary refractory.⁴ Retreatment of refractory or relapsed ALL is generally unsuccessful and associated with a high mortality rate.⁵ The 2-year survival rate among patients with ALL who relapse after hematopoietic cell transplantation is 15%.⁶

The U.S. Food and Drug Administration (FDA) approved clofarabine (as a single agent or in combination) in 2004 and blinatumomab in 2014 for relapsed and refractory ALL. Reported median objective response rates in the pivotal trials of the 2 agents were 19.7% and 33%, the median durations of response were 2.5 months and 6 months, and median overall survival durations were 3 months and 7.5 months, respectively.^{7,8} Note that the percentages of patients treated with 3 or more prior treatments of clofarabine and blinatumomab trial were 62% and 7%, respectively. Nevertheless, treatment options for patients with relapsed or refractory ALL are limited, associated with poor outcomes and high toxicity and the disease remains incurable.

Diffuse Large B Cell Lymphoma (DLBCL)

DLBCL is the most common histologic subtype of non-Hodgkin lymphoma and accounts for approximately 25% of non-Hodgkin lymphoma cases.⁹ DLBCL exhibits large heterogeneity in morphologic, genetic, and clinical aspects and multiple clinicopathologic entities are defined by the 2016 World Health Organization classification, which are sufficiently distinct to be considered separate diagnostic categories. The incidence of DLBCL is approximately 7 cases per 100,000 persons per year.¹⁰

Treatment

Treatment in the first-line setting include multiple chemotherapy and/or immunotherapy options that typically include rituximab. While majority of patients responds well to first-line immunochemotherapy combinations containing rituximab, 10 to 15% have primary refractory disease within 3 months after treatment initiation and another 20 to 35% have a relapse.¹¹ Of those who relapse or are refractory, 40 to 60% of patients may respond to second-line chemotherapy. Treatment of relapsed/refractory cases is generally stratified according to hematopoietic cell transplant eligibility. There is general consensus that salvage therapy followed by autologous transplantation is the preferred treatment for medically eligible patients with a first relapse of DLBCL or primary refractory DLBCL. Approximately 50% of patients who relapse or are refractory to first line agents proceed to undergo autologous hematopoietic stem-cell transplantation, and of these, approximately 30 to 40% remain progression-free 3 years after transplantation.^{12,13,14,15,16} Patients who are ineligible for as second-line therapy that includes high-dose chemotherapy and hematopoietic stem-cell transplantation, prognosis is often poor with a median overall survival of 4.4 months. Overall survival at 1 year is 23% and 16% at year 2.¹⁶ Patients who relapse after autologous transplantation, option are limited that includes allogeneic hematopoietic stem-cell transplantation. However, the procedure can only be performed if patient is chemo-responsive and a donor is available. Further the procedure is associated with a high risk of complications. Mortality risk unrelated to disease relapse is 23% at 1 year.^{17,18,19} FDA has also approved agents for refractory/relapsed DLBCL include pembrolizumab (Keytruda), polutuzumab vedotin-piiq (Polivy), selinexor (Xpovio) and tafasitamab-cxix (Monjuvi).

Mantle Cell Lymphoma (MCL)

MCL is a rare B-cell malignancy classified as an aggressive form of non-Hodgkin lymphoma that arises from cells originating in the “mantle zone” of the lymph node and typically affects men over the age of 60.

It accounts for approximately 3-6% of all non-Hodgkin lymphoma in the United States and differs from diffuse large B cell lymphoma (another subtype of non-Hodgkin lymphoma).^{20,21,22} In 2018, the overall incidence of MCL in the U.S was 3,500 with 5-year and 10-year prevalence of 12,000 and 18,000 cases. Median age at the time of diagnosis is 68, a majority of patients are non-Hispanic white males and more than 70% of patients present with stage IV disease.^{23,24} The majority (75%) of cases initially present with lymphadenopathy while presentation is extranodal in the remaining 25 percent. In most cases of MCL, chromosomal translocation results in aberrant expression of cyclin D1, leading to cell cycle dysregulation.²⁵ Many signaling pathways are constitutively activated and/or deregulated in MCL, including the B-cell receptor signaling pathway.²⁶

Treatment

In first line treatment of mantle cell lymphoma, preferred chemotherapy regimens often differ in affected patients due to the rarity and heterogeneity of the disease. There is no standard of care that exists for second-line and higher chemotherapy when a patient has relapsed or refractory MCL.²⁷ Second line therapies typically depend on the front line therapy utilized, comorbidities, the tumor's sensitivity to chemotherapy, and overall risk-benefit. Potential salvage regimens include ibrutinib, acalabrutinib, lenalidomide, combination chemotherapy, bortezomib, temsirolimus.

Despite availability of multiple treatments, MCL is not curable (with the possible exception of hematopoietic cell transplantation). Median overall survival in modern trials incorporating intensive therapy is 8 to 10 years with no plateau in the survival curve. Shorter survival times are seen with less intensive therapy. Multiple prognostic indices are used in MCL patients to guide course of treatment. First-line treatment of MCL can consist of aggressive therapy or less-aggressive therapy, depending on patient status at baseline.²⁶ It generally consists of chemotherapy in combination with rituximab. Only 30% to 40% of patients have a durable long term remission after chemo immunotherapy first line therapy.²⁸ Progression is common, with a median time to treatment failure of less than 18 months. Virtually all patients will have refractory or recurrent disease. Treatment of recurrent MCL is difficult, due to the rapid development of chemotherapy resistance. There are multiple preferred chemotherapy regimens that may be offered, and choice is primarily made based on prior treatment history, patient comorbidities and performance status. The expected toxicities of a given regimen as well as clinician's experience with the regimens are additional considerations. A preferred order for their use has not been established. Most of these regimens have not been compared directly in randomized trials. Given the limited efficacy of these agents and paucity of data comparing these various treatment options, participation in a clinical trial is encouraged whenever possible. Complete response rates in previously treated or relapsed MCL are generally low (<30%) and have limited response durations.¹³ Among patients who have disease progression after the receipt of Bruton's kinase inhibitor (BTK) therapy, the reported objective response rate ranges from 25 to 42% and a median overall survival of 6 to 10 months with salvage therapies.^{29,30,31,32} Allogeneic stem-cell transplantation may be an option for selected patients. However, non-relapse-related mortality remains high at 10 to 24%.³³

While the clinical course of MCL is generally aggressive, a small proportion of patients with low stage and low-risk disease may have an indolent course, managed by observation, splenectomy, or treatment with alkylating agents analogous to the treatment of patients with small lymphocytic lymphoma or follicular lymphoma.

Commercial Chimeric Antigen Receptor T Cell Therapies Available in the United States

As of August 2020, there are 3 CAR T cell therapies approved by the FDA for the treatment of cancer. All 3 are CD19-targeting CAR T cell immunotherapies in which patients own T-cells are genetically engineered using a viral vector to express a synthetic receptor called chimeric antigen receptor. Once injected, the genetically modified T-cells selectively target and bind to CD19 antigen expressed on the surface of B cells and tumors derived from B cells. Axicabtagene ciloleucel and brexucabtagene autoleucel are approved to treat subsets of patients with lymphoma while tisagenlecleucel is approved for treatment of subsets of patients with leukemia and lymphoma.

Summary

Chimeric antigen receptor (CAR) T cells are genetically engineered cells that represent a novel class of cancer immunotherapy. In general, the process of autologous CAR T-cell therapy begins with harvesting white blood cells from the patient via leukapheresis followed by T-cell receptor activation and genetic engineering via retroviral or lentiviral transduction. After the CAR T cells are generated, they are expanded to clinically relevant numbers, undergo quality control testing, and are cryopreserved. Commercial CAR T-cell products are manufactured at a centralized facility, necessitating transfer of the apheresis product to the manufacturing site, and the final cryopreserved CAR T-cell product back to the treatment facility. Typically, the patient undergoes lymphodepleting chemotherapy to create a favorable immune environment for CAR T-cell activity prior to receiving a single intravenous infusion of the product. Three commercial CAR T cell products have been approved by the U.S. Food and Drug Administration for the treatment of cancer. Tisagenlecleucel is approved for treatment of subsets of patients with leukemia and lymphoma and axicabtagene ciloleucel and brexucabtagene autoleucel are approved to treat subsets of patients with lymphoma.

Tisagenlecleucel

For individuals who are up to 25 years of age with relapsed or refractory B-cell acute lymphoblastic leukemia who receive tisagenlecleucel, the evidence includes a single-arm prospective trial. Relevant outcomes are OS, DSS, QOL, and treatment-related mortality and morbidity. The pivotal single-arm trials reported a 81% response rate (measured by complete response or complete remission with incomplete blood count) in heavily pretreated patients. All patients who achieved complete remission or complete remission with incomplete blood count were also minimal residual disease-negative, which is predictive of survival in acute lymphoblastic leukemia patients. After a median follow-up of 13.1 months, the median duration of response was not reached. The observed benefits seen with tisagenlecleucel were offset by a high frequency and severity of adverse events. Cytokine release syndrome was observed in more than half (77%) of the patients, and approximately 88% had an adverse event at grade 3 or higher. Long-term follow-up, real-world evidence, and post-marketing studies are required to assess the generalizability of tisagenlecleucel efficacy and safety outside of the clinical trial setting. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who are adults with a histologically confirmed diagnosis of aggressive non-Hodgkin lymphoma (eg, diffuse large B-cell lymphoma not otherwise specified, high-grade B-cell lymphoma, transformed follicular lymphoma) who receive tisagenlecleucel, the evidence includes a single-arm prospective trial. Relevant outcomes are OS, DSS, QOL, and treatment-related mortality and morbidity. The pivotal single-arm trial reported a 52% overall response rate (measured by complete or partial responses) in heavily pretreated patients. After a median follow-up of 14 months, the median duration of response was not reached. The observed benefits were offset by a high frequency and severity of adverse events. Any grade cytokine release syndrome was observed in 58% of the patients, and 63% had an adverse event at grade 3 or higher. Long-term follow-up, real-world evidence, and post-marketing studies are required to assess the generalizability of tisagenlecleucel efficacy and safety outside of the clinical trial setting. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Axicabtagene Ciloleucel

For individuals who are adults with a histologically confirmed diagnosis of aggressive non-Hodgkin lymphoma (eg, diffuse large B-cell lymphoma not otherwise specified, high-grade B-cell lymphoma, primary mediastinal large B-cell lymphoma, transformed follicular lymphoma) who receive axicabtagene ciloleucel, the evidence includes a single-arm prospective trial. Relevant outcomes are OS, DSS, QOL, and treatment-related mortality and morbidity. The pivotal single-arm trial reported a 83% overall response rate (measured by complete or partial remission) in heavily pretreated patients. After a median follow-up of 27.1 months, the median duration of response was 11.1 months. The observed benefits were offset by a high frequency and severity of adverse events. Cytokine release syndrome was observed in more than half of the patients, and 98% had an adverse event at grade 3 or higher. Long-term follow-up and real-world evidence are required to assess the generalizability of axicabtagene ciloleucel efficacy and safety outside of the clinical trial setting. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Brexucabtagene Autoleucl

For individuals who are adults with relapsed or refractory mantle cell lymphoma, the evidence includes one phase II single-arm study. Relevant outcomes are overall survival, disease-specific survival, quality of life, treatment-related mortality, and morbidity. ZUMA-2 study enrolled adult patients with relapsed refractory MCL who were heavily pre-treated. Of 74 patients enrolled, therapy was successfully manufactured for 71 (96%) and administered to 68 (92%). Results were reported for pre-specified 60 evaluable patients with a median follow-up (as of the July 24, 2019 data cutoff date) of 12.3 months (range, 7.0 to 32.3). The primary efficacy analysis demonstrated an objective response rate of 87% with a 62% rate of complete response. The median duration of response, progression-free survival, and median overall survival (OS) were not reached. Fifty-seven percent of patients remained in remission at data cutoff, and the estimated 12-month PFS and OS rates were 61% and 83%, respectively. Among patients who have disease progression after Bruton's kinase inhibitor therapy, the reported objective response rate ranges from 25 to 42% with a median overall survival of 6 to 10 months with salvage therapies. In the absence of randomized controlled trial, it is difficult to draw comparisons with currently available salvage treatment. In some clinical scenarios, patients who received CAR-T therapy through a clinical trial have demonstrated clinically significant improvements in overall response when treated with FDA approved CAR-T's. No notable study limitations were identified. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Policy History

Date	Action
4/2021	Clarified coding information.
3/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
1/2021	Clarified coding information.
12/2020	BCBSA National medical policy review. New medically necessary indications described for brexucabtagene autoleucl for mantle cell lymphoma. Clarified coding information. Effective 12/1/2020.
12/2019	New policy created from policy #455 Adoptive Immunotherapy. FDA-approved tisagenlecleucel (Kymriah) and axicabtagene ciloleucel (Yescarta) therapies were moved from policy #455 to create a new standalone policy #066 Chimeric Antigen Receptor Therapy for Hematologic Malignancies. Policy statements unchanged.
3/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
1/2019	Clarified coding information.
8/2018	BCBSA National medical policy review. Tisagenlecleucel added to the second medically necessary policy statement with modified criteria. Effective 8/1/2018.
6/2018	BCBSA National medical policy review. Policy statement clarified, changing "2 or 3" to "3", to read: "Patient has active central nervous system 3 acute lymphoblastic leukemia (ie, white blood cell count ≥ 5 cells/ μ L in cerebrospinal fluid with presence of lymphoblasts)." Prior Authorization Information reformatted.
4/2018	Clarified coding information.
2/2018	BCBSA National medical policy review. Policy criteria for Kymriah and Yescarta clarified.
1/2018	Clarified coding information. Preauthorization request form for Yescarta and Kymriah added.
11/2017	Medical policy criteria for Yescarta clarified. Effective 11/17/2017.
11/2017	BCBSA National medical policy review. New medically necessary indications added for Kymriah (tisagenlecleucel). Effective 11/7/2017. New medically necessary indications added for Yescarta (axicabtagene ciloleucel). Effective 11/7/2017. Clarified coding information.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

References

1. Berry DA, Zhou S, Higley H, et al. Association of Minimal Residual Disease With Clinical Outcome in Pediatric and Adult Acute Lymphoblastic Leukemia: A Meta-analysis. *JAMA Oncol.* Jul 13 2017; 3(7): e170580. PMID 28494052
2. Hunger SP, Mullighan CG. Acute Lymphoblastic Leukemia in Children. *N Engl J Med.* Oct 15 2015; 373(16): 1541-52. PMID 26465987
3. Maude SL, Teachey DT, Porter DL, et al. CD19-targeted chimeric antigen receptor T-cell therapy for acute lymphoblastic leukemia. *Blood.* Jun 25 2015; 125(26): 4017-23. PMID 25999455
4. Pui CH, Carroll WL, Meshinchi S, et al. Biology, risk stratification, and therapy of pediatric acute leukemias: an update. *J Clin Oncol.* Feb 10 2011; 29(5): 551-65. PMID 21220611
5. Tallen G, Ratei R, Mann G, et al. Long-term outcome in children with relapsed acute lymphoblastic leukemia after time-point and site-of-relapse stratification and intensified short-course multidrug chemotherapy: results of trial ALL-REZ BFM 90. *J Clin Oncol.* May 10 2010; 28(14): 2339-47. PMID 20385996
6. Bajwa R, Schechter T, Soni S, et al. Outcome of children who experience disease relapse following allogeneic hematopoietic SCT for hematologic malignancies. *Bone Marrow Transplant.* May 2013; 48(5): 661-5. PMID 23128573
7. Jeha S, Gaynon PS, Razzouk BI, et al. Phase II study of clofarabine in pediatric patients with refractory or relapsed acute lymphoblastic leukemia. *J Clin Oncol.* Apr 20 2006; 24(12): 1917-23. PMID 16622268
8. von Stackelberg A, Locatelli F, Zugmaier G, et al. Phase I/Phase II Study of Blinatumomab in Pediatric Patients With Relapsed/Refractory Acute Lymphoblastic Leukemia. *J Clin Oncol.* Dec 20 2016; 34(36): 4381-4389. PMID 27998223
9. Swerdlow SH, Campo E, Pileri SA, et al. The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood.* May 19 2016; 127(20): 2375-90. PMID 26980727
10. Morton LM, Wang SS, Devesa SS, et al. Lymphoma incidence patterns by WHO subtype in the United States, 1992-2001. *Blood.* Jan 01 2006; 107(1): 265-76. PMID 16150940
11. Sehn LH, Gascoyne RD. Diffuse large B-cell lymphoma: optimizing outcome in the context of clinical and biologic heterogeneity. *Blood.* Jan 01 2015; 125(1): 22-32. PMID 25499448
12. Crump M, Neelapu SS, Farooq U, et al. Outcomes in refractory diffuse large B-cell lymphoma: results from the international SCHOLAR-1 study. *Blood.* Oct 19 2017; 130(16): 1800-1808. PMID 28774879
13. Gisselbrecht C, Glass B, Mounier N, et al. Salvage regimens with autologous transplantation for relapsed large B-cell lymphoma in the rituximab era. *J Clin Oncol.* Sep 20 2010; 28(27): 4184-90. PMID 20660832
14. Sehn LH, Assouline SE, Stewart DA, et al. A phase 1 study of obinutuzumab induction followed by 2 years of maintenance in patients with relapsed CD20-positive B-cell malignancies. *Blood.* May 31 2012; 119(22): 5118-25. PMID 22438256
15. Crump M, Kuruvilla J, Couban S, et al. Randomized comparison of gemcitabine, dexamethasone, and cisplatin versus dexamethasone, cytarabine, and cisplatin chemotherapy before autologous stem-cell transplantation for relapsed and refractory aggressive lymphomas: NCIC-CTG LY.12. *J Clin Oncol.* Nov 01 2014; 32(31): 3490-6. PMID 25267740
16. Van Den Neste E, Schmitz N, Mounier N, et al. Outcome of patients with relapsed diffuse large B-cell lymphoma who fail second-line salvage regimens in the International CORAL study. *Bone Marrow Transplant.* Jan 2016; 51(1): 51-7. PMID 26367239
17. Rigacci L, Puccini B, Doderio A, et al. Allogeneic hematopoietic stem cell transplantation in patients with diffuse large B cell lymphoma relapsed after autologous stem cell transplantation: a GITMO study. *Ann Hematol.* Jun 2012; 91(6): 931-9. PMID 22245922

18. Lazarus HM, Zhang MJ, Carreras J, et al. A comparison of HLA-identical sibling allogeneic versus autologous transplantation for diffuse large B cell lymphoma: a report from the CIBMTR. *Biol Blood Marrow Transplant.* Jan 2010; 16(1): 35-45. PMID 20053330
19. Van Den Neste E, Schmitz N, Mounier N, et al. Outcomes of diffuse large B-cell lymphoma patients relapsing after autologous stem cell transplantation: an analysis of patients included in the CORAL study. *Bone Marrow Transplant.* Feb 2017; 52(2): 216-221. PMID 27643872
20. A clinical evaluation of the International Lymphoma Study Group classification of non-Hodgkin's lymphoma. The Non-Hodgkin's Lymphoma Classification Project. *Blood.* Jun 01 1997; 89(11): 3909-18. PMID 9166827
21. Zhou Y, Wang H, Fang W, et al. Incidence trends of mantle cell lymphoma in the United States between 1992 and 2004. *Cancer.* Aug 15 2008; 113(4): 791-8. PMID 18615506
22. Teras LR, DeSantis CE, Cerhan JR, et al. 2016 US lymphoid malignancy statistics by World Health Organization subtypes. *CA Cancer J Clin.* Nov 12 2016; 66(6): 443-459. PMID 27618563
23. Fu S, Wang M, Lairson DR, et al. Trends and variations in mantle cell lymphoma incidence from 1995 to 2013: A comparative study between Texas and National SEER areas. *Oncotarget.* Dec 22 2017; 8(68): 112516-112529. PMID 29348844
24. Federal Register / Vol. 85, No. 104 / Friday, May 29, 2020 / Proposed Rules; KTEX19; Page 175. Federal Register / Vol. 85, No. 104 / Friday, May 29, 2020 / Proposed Rules. Accessed July 24, 2020. <https://www.govinfo.gov/content/pkg/FR-2020-05-29/pdf/2020-10122.pdf>
25. Argatoff LH, Connors JM, Klasa RJ, et al. Mantle cell lymphoma: a clinicopathologic study of 80 cases. *Blood.* Mar 15 1997; 89(6): 2067-78. PMID 9058729
26. Jares P, Colomer D, Campo E. Molecular pathogenesis of mantle cell lymphoma. *J Clin Invest.* Oct 2012; 122(10): 3416-23. PMID 23023712
27. Campo E, Rule S. Mantle cell lymphoma: evolving management strategies. *Blood.* Jan 01 2015; 125(1): 48-55. PMID 25499451
28. Flinn IW, van der Jagt R, Kahl B, et al. First-Line Treatment of Patients With Indolent Non-Hodgkin Lymphoma or Mantle-Cell Lymphoma With Bendamustine Plus Rituximab Versus R-CHOP or R-CVP: Results of the BRIGHT 5-Year Follow-Up Study. *J Clin Oncol.* Apr 20 2019; 37(12): 984-991. PMID 30811293
29. Cheah CY, Seymour JF, Wang ML. Mantle Cell Lymphoma. *J Clin Oncol.* Apr 10 2016; 34(11): 1256-69. PMID 26755518
30. Martin P, Maddocks K, Leonard JP, et al. Postibrutinib outcomes in patients with mantle cell lymphoma. *Blood.* Mar 24 2016; 127(12): 1559-63. PMID 26764355
31. Jain P, Kanagal-Shamanna R, Zhang S, et al. Long-term outcomes and mutation profiling of patients with mantle cell lymphoma (MCL) who discontinued ibrutinib. *Br J Haematol.* Nov 2018; 183(4): 578-587. PMID 30175400
32. Epperla N, Hamadani M, Cashen AF, et al. Predictive factors and outcomes for ibrutinib therapy in relapsed/refractory mantle cell lymphoma-a real world study. *Hematol Oncol.* Dec 2017; 35(4): 528-535. PMID 28066928
33. Robinson SP, Boumendil A, Finel H, et al. Long-term outcome analysis of reduced-intensity allogeneic stem cell transplantation in patients with mantle cell lymphoma: a retrospective study from the EBMT Lymphoma Working Party. *Bone Marrow Transplant.* May 2018; 53(5): 617-624. PMID 29335632
34. National Comprehensive Cancer Network (NCCN). Acute Lymphoblastic Leukemia. Version 1.2020. 2020 Jan 15; https://www.nccn.org/professionals/physician_gls/pdf/ped_all.pdf. Accessed August 28, 2020.
35. Lee DW, Santomaso BD, Locke FL, et al. ASTCT Consensus Grading for Cytokine Release Syndrome and Neurologic Toxicity Associated with Immune Effector Cells. *Biol Blood Marrow Transplant.* Apr 2019; 25(4): 625-638. PMID 30592986
36. National Cancer Institute. U.S. Department of Health and Human Services. Common Terminology Criteria for Adverse Events (CTCAE) Version 5.0. 2017 Nov 27; https://ctep.cancer.gov/protocolDevelopment/electronic_applications/docs/CTCAE_v5_Quick_Reference_8.5x11.pdf. Accessed September 2, 2019.
37. Maude SL, Laetsch TW, Buechner J, et al. Tisagenlecleucel in Children and Young Adults with B-Cell Lymphoblastic Leukemia. *N Engl J Med.* Feb 01 2018; 378(5): 439-448. PMID 29385370

38. National Comprehensive Cancer Network (NCCN). B-Cell Lymphomas. Version 4.2020. 2020 Aug 13; <https://www.nccn.org/professionals/physician/gls/pdf/b-cell.pdf>. Accessed August 29, 2020.
39. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. Feb 10 2007; 25(5): 579-86. PMID 17242396
40. Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol*. Sep 20 2014; 32(27): 3059-68. PMID 25113753
41. Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in Adult Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med*. Jan 03 2019; 380(1): 45-56. PMID 30501490
42. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol*. Jan 2019; 20(1): 31-42. PMID 30518502
43. Kite Pharma Inc. Prescribing Label: Yescarta (axicabtagene ciloleucel) suspension for intravenous infusion. 2017; <https://www.fda.gov/downloads/BiologicsBloodVaccines/CellularGeneTherapyProducts/ApprovedProducts/UCM581226.pdf>. Accessed September 2, 2019.
44. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. *N Engl J Med*. Dec 28 2017; 377(26): 2531-2544. PMID 29226797
45. Novartis Pharmaceuticals. Prescribing Label: Kymriah (tisagenlecleucel) suspension for intravenous infusion. 2018; <https://www.pharma.us.novartis.com/sites/www.pharma.us.novartis.com/files/kymriah.pdf>. Accessed September 2, 2019.
46. Tecartus (brexucabtagene autoleucel) suspension for intravenous infusion. Prescribing label. Accessed on September 1, 2020. <https://www.gilead.com/-/media/files/pdfs/medicines/oncology/tecartus/tecartus-pi.pdf>
47. Wang M, Munoz J, Goy A, et al. KTE-X19 CAR T-Cell Therapy in Relapsed or Refractory Mantle-Cell Lymphoma. *N Engl J Med*. Apr 02 2020; 382(14): 1331-1342. PMID 32242358
48. Centers for Medicare & Medicaid Services (CMS). Proposed Decision Memo for Chimeric Antigen Receptor (CAR) T-cell Therapy for Cancers (CAG-00451N). 2017 Feb 15; <https://www.cms.gov/medicare-coverage-database/details/nca-proposed-decision-memo.aspx?NCAId=291>. Accessed September 2, 2019.