



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy Facet Joint Denervation

Table of Contents

- [Policy: Commercial](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [References](#)
- [Information Pertaining to All Policies](#)

Policy Number: 140

BCBSA Reference Number: 7.01.116 (For Plans internal use only)

Related Policies

- Diagnosis and Treatment of Sacroiliac Joint Pain, [#320](#)
- Facet Arthroplasty, [#174](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Non-pulsed radiofrequency denervation of cervical facet joints (C3-4 and below) and lumbar facet joints may be **MEDICALLY NECESSARY** when all of the following criteria are met:

- No prior spinal fusion surgery in the vertebral level being treated, AND
- Disabling low back (lumbosacral) or neck (cervical) pain, suggestive of facet joint origin as evidenced by absence of nerve root compression as documented in the medical record on history, physical and radiographic evaluations; and the pain is not radicular, AND
- Pain has failed to respond to three (3) months of conservative management which may consist of therapies such as nonsteroidal anti-inflammatory medications, acetaminophen, manipulation, physical therapy, and a home exercise program, AND
- A trial of controlled diagnostic medial branch blocks* AND
- If there has been a prior successful radiofrequency (RF) denervation, a minimum time of six (6) months has elapsed since prior RF treatment (per side, per anatomical level of the spine.)

*A successful trial of controlled diagnostic medial branch blocks consists of 2 separate positive blocks on different days with local anesthetic only (no steroids or other drugs), or a placebo-controlled series of blocks, under fluoroscopic guidance, that has resulted in at least a 50% reduction in pain.

Radiofrequency denervation is **INVESTIGATIONAL** for the treatment of chronic spinal/back pain for all uses that do not meet the criteria listed above, including but not limited to treatment of thoracic facet joint pain.

All other methods of denervation are **INVESTIGATIONAL** for the treatment of chronic spinal/back pain, including, but not limited to pulsed radiofrequency denervation, laser denervation, chemodenervation (e.g., alcohol, phenol, or high-concentration local anesthetics), and cryodenervation.

Therapeutic medial branch blocks are **INVESTIGATIONAL**.

If there has been a prior successful radiofrequency (RF) denervation, additional diagnostic medial branch blocks for the same level of the spine are **NOT MEDICALLY NECESSARY**.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is not required .
Commercial PPO and Indemnity	Prior authorization is not required .

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:

CPT Codes

CPT codes:	Code Description
64633	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance (fluoroscopy or CT); cervical or thoracic, single facet joint
64634	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance (fluoroscopy or CT); cervical or thoracic, each additional facet joint
64635	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance (fluoroscopy or CT); lumbar or sacral, single facet joint
64636	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance (fluoroscopy or CT); lumbar or sacral, each additional facet joint

Description

Percutaneous radiofrequency (RF) facet denervation is used to treat neck or back pain originating in facet joints with degenerative changes. Diagnosis of facet joint pain is confirmed by response to nerve blocks. Patients generally are sedated for the RF procedure. The goal of facet denervation is long-term pain relief. However, the nerves regenerate and, therefore, repeat procedures may be required.

Facet joint denervation is performed under local anesthetic and with fluoroscopic guidance. A needle or probe is directed to the median branch of the dorsal ganglion innervating the facet joint, where multiple thermal lesions are produced, typically by an RF generator. A variety of terms may be used to describe

RF denervation (eg, rhizotomy, rhizolysis). In addition, the structures to which the RF energy is directed may be referred to as facet joint, facet nerves, medial nerve or branch, median nerve or branch, or dorsal root ganglion.

Alternative methods of denervation include pulsed RF, laser, chemodenervation, and cryoablation. Pulsed RF consists of short bursts of electric current of high voltage in the RF range but without heating the tissue enough to cause coagulation. RF is suggested as a possibly safer alternative to thermal RF facet denervation. Temperatures do not exceed 42°C at the probe tip versus temperatures in the 60°C range reached in thermal RF denervation, and tissues may cool between pulses. It is postulated that transmission across small unmyelinated nerve fibers is disrupted but not permanently damaged, while large, myelinated fibers are not affected. With chemical denervation, injections with a diluted phenol solution, a chemical ablating agent, are injected into the facet joint nerve.

Summary

Percutaneous radiofrequency (RF) facet denervation is used to treat neck and back pain originating in facet joints with degenerative changes. Diagnosis of facet joint pain is confirmed by response to nerve blocks. The goal of facet denervation is long-term pain relief. However, the nerves regenerate and, therefore, repeat procedures may be required.

For individuals with suspected facet joint pain who receive diagnostic medial branch blocks, the evidence includes systematic reviews, a small, randomized trial, and observational studies. Relevant outcomes are other test performance measures, symptoms, and functional outcomes. There is considerable controversy about the role of these blocks, the number of positive blocks required, and the extent of pain relief obtained. Studies have reported the use of single or double blocks and at least 50% or 80% improvement in pain and function. This evidence has suggested that there are relatively few patients who exhibit pain relief following 2 nerve blocks, but that these select patients may have pain relief for several months following RF denervation. Other large series have reported the prevalence and false-positive rates following controlled diagnostic blocks, although there are issues with the reference standards used in these studies because there is no criterion standard for the diagnosis of facet joint pain. There is level I evidence for the use of medial branch blocks for diagnosing chronic lumbar facet joint pain and level II evidence for diagnosing cervical and thoracic facet joint pain. The evidence available supports a threshold of at least 75% to 80% pain relief to reduce the false-positive rate. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with facet joint pain who receive RF ablation, the evidence includes systematic reviews and randomized controlled trials (RCTs). Relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. While the evidence is limited to RCTs with small sample sizes, RF facet denervation appears to provide at least 50% pain relief in carefully selected patients. Diagnosis of facet joint pain is difficult. However, response to controlled medial branch blocks and the presence of tenderness over the facet joint appear to be reliable predictors of success. When RF facet denervation is successful, repeat treatments appear to have similar success rates and duration of pain relief. Thus, the data indicate that, in carefully selected individuals with lumbar or cervical facet joint pain, RF treatments can improve outcomes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with facet joint pain who receive therapeutic medial nerve branch blocks or alternative methods of facet joint denervation, the evidence includes a systematic review, randomized trials without a sham control, and uncontrolled case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and medication use. Pulsed RF does not appear to be as effective as conventional RF denervation, and there is insufficient evidence to evaluate the efficacy of other methods of denervation (eg, alcohol, laser, cryodenervation) for facet joint pain or the effect of therapeutic medial branch blocks on facet joint pain. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
1/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
1/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2018	Annual policy review. New references added
1/2016	Annual policy review. New references added
1/2015	Clarified coding information.
12/2014	Annual policy review. New references added.
6/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
3/2014	Annual policy review. New investigational indications described. Effective 3/1/2014. Coding information clarified.
6/2013	Annual policy review. New investigational indications described. Effective 6/1/2013.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
6/2011	Reviewed - Medical Policy Group – Orthopedics, Rehabilitation and Rheumatology. No changes to policy statements.
5/2011	Updated description based on annual policy review.
1/2011	Updated to clarify non-coverage of pulsed radiofrequency denervation.
10/2010	Annual policy review. No changes to policy statements.
7/2010	Reviewed - Medical Policy Group – Orthopedics, Rehabilitation Medicine and Rheumatology. No changes to policy statements.
7/2010	Updated to clarify the information in the coverage sections.
11/1/2009.	Medical Policy #140 issued. Effective 11/1/2009.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Boswell MV, Manchikanti L, Kaye AD, et al. A Best-Evidence Systematic Appraisal of the Diagnostic Accuracy and Utility of Facet (Zygapophysial) Joint Injections in Chronic Spinal Pain. Pain Physician. Jul-Aug 2015; 18(4): E497-533. PMID 26218947
2. Falco FJ, Datta S, Manchikanti L, et al. An updated review of the diagnostic utility of cervical facet joint injections. Pain Physician. Nov-Dec 2012; 15(6): E807-38. PMID 23159977
3. Falco FJ, Manchikanti L, Datta S, et al. Systematic review of the therapeutic effectiveness of cervical facet joint interventions: an update. Pain Physician. Nov-Dec 2012; 15(6): E839-68. PMID 23159978
4. Falco FJ, Manchikanti L, Datta S, et al. An update of the systematic assessment of the diagnostic accuracy of lumbar facet joint nerve blocks. Pain Physician. Nov-Dec 2012; 15(6): E869-907. PMID 23159979
5. Falco FJ, Manchikanti L, Datta S, et al. An update of the effectiveness of therapeutic lumbar facet joint interventions. Pain Physician. Nov-Dec 2012; 15(6): E909-53. PMID 23159980

6. Cohen SP, Strassels SA, Kurihara C, et al. Randomized study assessing the accuracy of cervical facet joint nerve (medial branch) blocks using different injectate volumes. *Anesthesiology*. Jan 2010; 112(1): 144-52. PMID 19996954
7. Cohen SP, Stojanovic MP, Crooks M, et al. Lumbar zygapophysial (facet) joint radiofrequency denervation success as a function of pain relief during diagnostic medial branch blocks: a multicenter analysis. *Spine J*. May-Jun 2008; 8(3): 498-504. PMID 17662665
8. Pampati S, Cash KA, Manchikanti L. Accuracy of diagnostic lumbar facet joint nerve blocks: a 2-year follow-up of 152 patients diagnosed with controlled diagnostic blocks. *Pain Physician*. Sep-Oct 2009; 12(5): 855-66. PMID 19787011
9. Manchikanti L, Pampati S, Cash KA. Making sense of the accuracy of diagnostic lumbar facet joint nerve blocks: an assessment of the implications of 50% relief, 80% relief, single block, or controlled diagnostic blocks. *Pain Physician*. Mar-Apr 2010; 13(2): 133-43. PMID 20309379
10. Janapala RN, Manchikanti L, Sanapati MR, et al. Efficacy of Radiofrequency Neurotomy in Chronic Low Back Pain: A Systematic Review and Meta-Analysis. *J Pain Res*. 2021; 14: 2859-2891. PMID 34531682
11. Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (Facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: a randomized double-blind trial. *Spine (Phila Pa 1976)*. May 20 2008; 33(12): 1291-7; discussion 1298. PMID 18496338
12. Tekin I, Mirzai H, Ok G, et al. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain*. Jul-Aug 2007; 23(6): 524-9. PMID 17575493
13. van Wijk RM, Geurts JW, Wynne HJ, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain*. Jul-Aug 2005; 21(4): 335-44. PMID 15951652
14. Lakemeier S, Lind M, Schultz W, et al. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: a randomized, controlled, double-blind trial. *Anesth Analg*. Jul 2013; 117(1): 228-35. PMID 23632051
15. Manchikanti L, Kaye AD, Boswell MV, et al. A Systematic Review and Best Evidence Synthesis of the Effectiveness of Therapeutic Facet Joint Interventions in Managing Chronic Spinal Pain. *Pain Physician*. Jul-Aug 2015; 18(4): E535-82. PMID 26218948
16. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg*. 2012; 22(2): 200-6. PMID 22437295
17. Lord SM, Barnsley L, Wallis BJ, et al. Percutaneous radio-frequency neurotomy for chronic cervical zygapophyseal-joint pain. *N Engl J Med*. Dec 05 1996; 335(23): 1721-6. PMID 8929263
18. van Eerd M, de Meij N, Kessels A, et al. Efficacy and Long-term Effect of Radiofrequency Denervation in Patients with Clinically Diagnosed Cervical Facet Joint Pain: A Double-blind Randomized Controlled Trial. *Spine (Phila Pa 1976)*. Mar 01 2021; 46(5): 285-293. PMID 33534439
19. Husted DS, Orton D, Schofferman J, et al. Effectiveness of repeated radiofrequency neurotomy for cervical facet joint pain. *J Spinal Disord Tech*. Aug 2008; 21(6): 406-8. PMID 18679094
20. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine (Phila Pa 1976)*. Nov 01 2004; 29(21): 2471-3. PMID 15507813
21. Smuck M, Crisostomo RA, Trivedi K, et al. Success of initial and repeated medial branch neurotomy for zygapophysial joint pain: a systematic review. *PM R*. Sep 2012; 4(9): 686-92. PMID 22980421
22. Rambaransingh B, Stanford G, Burnham R. The effect of repeated zygapophysial joint radiofrequency neurotomy on pain, disability, and improvement duration. *Pain Med*. Sep 2010; 11(9): 1343-7. PMID 20667024
23. Manchikanti L, Singh V, Falco FJ, et al. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: a randomized, double-blind controlled trial. *Pain Physician*. Sep-Oct 2010; 13(5): 437-50. PMID 20859313
24. Manchikanti L, Singh V, Falco FJ, et al. Evaluation of lumbar facet joint nerve blocks in managing chronic low back pain: a randomized, double-blind, controlled trial with a 2-year follow-up. *Int J Med Sci*. May 28 2010; 7(3): 124-35. PMID 20567613
25. Manchikanti L, Singh V, Falco FJ, et al. Comparative effectiveness of a one-year follow-up of thoracic medial branch blocks in management of chronic thoracic pain: a randomized, double-blind active controlled trial. *Pain Physician*. Nov-Dec 2010; 13(6): 535-48. PMID 21102966

26. Manchikanti L, Singh V, Falco FJ, et al. The role of thoracic medial branch blocks in managing chronic mid and upper back pain: a randomized, double-blind, active-control trial with a 2-year followup. *Anesthesiol Res Pract.* 2012; 2012: 585806. PMID 22851967
27. Hashemi M, Hashemian M, Mohajerani SA, et al. Effect of pulsed radiofrequency in treatment of facet-joint origin back pain in patients with degenerative spondylolisthesis. *Eur Spine J.* Sep 2014; 23(9): 1927-32. PMID 24997616
28. Kroll HR, Kim D, Danic MJ, et al. A randomized, double-blind, prospective study comparing the efficacy of continuous versus pulsed radiofrequency in the treatment of lumbar facet syndrome. *J Clin Anesth.* Nov 2008; 20(7): 534-7. PMID 19041042
29. Van Zundert J, Patijn J, Kessels A, et al. Pulsed radiofrequency adjacent to the cervical dorsal root ganglion in chronic cervical radicular pain: a double blind sham controlled randomized clinical trial. *Pain.* Jan 2007; 127(1-2): 173-82. PMID 17055165
30. Iwatsuki K, Yoshimine T, Awazu K. Alternative denervation using laser irradiation in lumbar facet syndrome. *Lasers Surg Med.* Mar 2007; 39(3): 225-9. PMID 17345622
31. Joo YC, Park JY, Kim KH. Comparison of alcohol ablation with repeated thermal radiofrequency ablation in medial branch neurotomy for the treatment of recurrent thoracolumbar facet joint pain. *J Anesth.* Jun 2013; 27(3): 390-5. PMID 23192698
32. Haufe SM, Mork AR. Endoscopic facet debridement for the treatment of facet arthritic pain--a novel new technique. *Int J Med Sci.* May 25 2010; 7(3): 120-3. PMID 20567612
33. Watters WC, Resnick DK, Eck JC, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 13: injection therapies, low-back pain, and lumbar fusion. *J Neurosurg Spine.* Jul 2014; 21(1): 79-90. PMID 24980590
34. Manchikanti L, Kaye AD, Soin A, et al. Comprehensive Evidence-Based Guidelines for Facet Joint Interventions in the Management of Chronic Spinal Pain: American Society of Interventional Pain Physicians (ASIPP) Guidelines Facet Joint Interventions 2020 Guidelines. *Pain Physician.* May 2020; 23(3S): S1-S127. PMID 32503359
35. Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. *Reg Anesth Pain Med.* Jun 2020; 45(6): 424-467. PMID 32245841
36. National Institute for Health and Clinical Excellence (NICE). NICE guideline [NG59]: Low back pain and sciatica in over 16s: assessment and management. 2016; <https://www.nice.org.uk/guidance/NG59>. Accessed September 29, 2021.
37. Evidence Based Clinical Guidelines for Multidisciplinary Spine Care: Diagnosis and Treatment of Low Back Pain. American Spine Society. Published 2020. Accessed September 29, 2021.