



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Heart Transplant

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 197

BCBSA Reference Number: 7.03.09 (For Plan internal use only)

NCD/LCD: N/A

Related Policies

- Heart-Lung Transplant, #[269](#)
- Total Artificial Hearts and Ventricular Assist Devices, #[280](#)
- Immune Cell Function Assay in Solid Organ Transplantation, #[182](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Human heart transplantation may be considered **MEDICALLY NECESSARY** for selected adults and children with end-stage heart failure when any one of the following criteria are met:

Adult Patients

1. Accepted Indications for Transplantation
 - a. Hemodynamic compromise due to heart failure demonstrated by any of the following 3 bulleted items:
 - Maximal Vo_2 (oxygen consumption) <10 mL/kg/min with achievement of anaerobic metabolism
 - Refractory cardiogenic shock
 - Documented dependence on intravenous inotropic support to maintain adequate organ perfusion
 - or
 - b. Severe ischemia consistently limiting routine activity not amenable to bypass surgery or angioplasty, or
 - c. Recurrent symptomatic ventricular arrhythmias refractory to ALL accepted therapeutic modalities.
2. Probable Indications for Cardiac Transplantation
 - a. Maximal Vo_2 <14 mL/kg/min and major limitation of the patient's activities, or
 - b. Recurrent unstable ischemia not amenable to bypass surgery or angioplasty, or

- c. Instability of fluid balance/renal function not due to patient noncompliance with regimen of weight monitoring, flexible use of diuretic drugs, and salt restriction.
- 3. The following conditions are inadequate indications for transplantation unless other factors as listed above are present:
 - a. Ejection fraction <20%
 - b. History of functional class III or IV symptoms of heart failure
 - c. Previous ventricular arrhythmias
 - d. Maximal $\text{Vo}_2 > 15 \text{ mL/kg/min}$.

Pediatric Patients

Patients with heart failure with persistent symptoms at rest who require one or more of the following:

- Continuous infusion of intravenous inotropic agents, or
- Mechanical ventilatory support, or
- Mechanical circulatory support, or

Patients with pediatric heart disease with symptoms of heart failure who do not meet the above criteria but who have:

- Severe limitation of exercise and activity (if measurable, such patients would have a peak maximum oxygen consumption <50% predicted for age and sex); or
- Cardiomyopathies or previously repaired or palliated congenital heart disease and growth failure attributable to the heart disease; or
- Near sudden death and/or life-threatening arrhythmias untreatable with medications or an implantable defibrillator; or
- Restrictive cardiomyopathy with reactive pulmonary hypertension; or
- Reactive pulmonary hypertension and risk of developing fixed, irreversible elevation of pulmonary vascular resistance that could preclude orthotopic heart transplantation in the future; or
- Anatomical and physiological conditions likely to worsen the natural history of congenital heart disease in infants with a functional single ventricle; or
- Anatomical and physiological conditions that lead to heart transplantation without systemic ventricular dysfunction.

Heart retransplantation after a failed primary heart transplant may be considered **MEDICALLY NECESSARY** in patients who meet criteria for heart transplantation.

Heart transplantation is **INVESTIGATIONAL** in all other situations.

In addition to the above information, we do not cover heart transplantation when any of the following conditions are present:

- Known current malignancy, including metastatic cancer
- Recent malignancy with high risk of recurrence
 - Note: the assessment of risk of recurrence for a previously treated malignancy is made by the transplant team; providers must submit a statement with an explanation of why the patient with a recently treated malignancy is an appropriate candidate for a transplant.
- Untreated systemic infection making immunosuppression unsafe, including chronic infection
- Other irreversible end-stage disease not attributed to heart or lung disease
- History of cancer with a moderate risk of recurrence
- Systemic disease that could be exacerbated by immunosuppression
- Psychosocial conditions or chemical dependency affecting ability to adhere to therapy
- Pulmonary hypertension that is fixed as evidenced by pulmonary vascular resistance (PVR) greater than 5 Wood units, or transpulmonary gradient (TPG) greater than or equal to 16 mm/Hg despite treatment*
- Severe pulmonary disease despite optimal medical therapy, not expected to improve with heart transplantation. *

*Some patients may be candidates for combined heart-lung transplantation (See policy #269).

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This procedure is performed in the inpatient setting.
Commercial PPO and Indemnity	This procedure is performed in the inpatient setting.
Medicare HMO BlueSM	This procedure is performed in the inpatient setting.
Medicare PPO BlueSM	This procedure is performed in the inpatient setting.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above **medical necessity criteria MUST** be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
33945	Heart transplant, with or without recipient cardiectomy

ICD-10 Procedure Codes

ICD-10-PCS procedure codes:	Code Description
02YA0Z0	Transplantation of Heart, Allogeneic, Open Approach
02YA0Z1	Transplantation of Heart, Syngeneic, Open Approach

Description

Solid Organ Transplantation

Solid organ transplantation offers a treatment option for patients with different types of end-stage organ failure that can be lifesaving or provide significant improvements to a patient's quality of life.¹ Many advances have been made in the last several decades to reduce perioperative complications. Available data supports improvement in long-term survival as well as improved quality of life, particularly for liver, kidney, pancreas, heart, and lung transplants. Allograft rejection remains a key early and late complication risk for any organ transplantation. Transplant recipients require life-long immunosuppression to prevent rejection. Patients are prioritized for transplant by mortality risk and severity of illness criteria developed by Organ Procurement and Transplantation Network and United Network for Organ Sharing.

Heart Transplant

In 2020, 39,036 transplants were performed in the United States procured from almost 33,300 deceased donors and 5,700 living donors.² Heart transplants were the third most common procedure with 3,658 transplants performed from both deceased and living donors in 2020. As of June 2021, there were 3,597 patients on the waiting list for a heart transplant.³

Most heart transplant recipients now are hospitalized as status 1 patients at the time of transplant. This shift has occurred due to the increasing demand for the scarce resource of donor organs resulting in an increased waiting time for recipients. Patients initially listed as status 2 candidates may deteriorate to a status 1 candidate before a donor organ becomes available. Alternatively, as medical and device therapy for advanced heart failure improves, some patients on the transplant list will recover enough function to be delisted. Lietz and Miller (2007) reported on survival for patients on the heart transplant waiting list, comparing the era between 1990 and 1994 with the era of 2000 to 2005.⁴ One-year survival for United Network for Organ Sharing status 1 candidate improved from 49.5% to 69.0%. Status 2 candidates fared even better, with 89.4% surviving 1 year compared with 81.8% in the earlier time period.

Johnson et al (2010) reported on waiting list trends in the U. S. between 1999 and 2008.⁵ The proportion of patients listed as status 1 increased, even as the waiting list and posttransplant mortality for this group have decreased. Meanwhile, status 2 patients have decreased as a proportion of all candidates. Completed transplants have trended toward the extremes of age, with more infants and patients older than age 65 years having transplants in recent years.

Alshawabkeh et al (2018) reported on the 1-year probability of the combined outcome of death or delisting due to clinical worsening for patients on the heart transplant waiting list, comparing the periods of April 1, 1986 to January 19, 1999, (early era) and January 20, 1999 to June 2, 2014 (current era).⁶ For adults without congenital heart disease (CHD), the probability of the combined outcome was lower in the current era compared with the early era, regardless of whether the patient was listed in status I (14.5% vs 22.7%; $p < 0.0001$) or 2 (9.0% vs 12.8%, $p < 0.0001$). When comparing the current and early eras in adults with CHD, a reduction in the probability of the combined outcome was demonstrated in those listed in status I (17.6% vs 43.3%, respectively; $p < 0.0001$), whereas the outcome remained unchanged for those listed in status 2 (10.6% vs 10.4%, respectively; $p = 0.94$).

In adults with CHD, factors associated with waitlist death or delisting due to clinical worsening within 1 year were also examined by Alshawabkeh et al (2016).⁷ A multivariate analysis identified that an estimated glomerular filtration rate less than 60 ml/min/1.73 m² (hazard ratio [HR], 1.4; 95% confidence interval [CI], 1.0 to 1.9; $p = 0.043$), albumin less than 3.2 g/dl (HR, 2.0; 95% CI, 1.3 to 2.9; $p < 0.001$), and hospitalization at the time of listing in the intensive care unit (HR, 2.3; 95% CI, 1.6 to 3.5; $p < 0.001$) or a non-intensive care hospital unit (HR, 1.9; 95% CI, 1.2 to 3.0; $p = 0.006$) were associated with waitlist death or delisting due to clinical worsening within 1 year.

Magnetta et al (2019) reported outcomes for children on the heart transplant waiting list, comparing the periods of December 16, 2011 to March 21, 2016 (era 1), and March 22, 2016 to June 30, 2018 (era 2).⁸ There was a significant decrease from era 1 to era 2 in the proportion of patients listed as status 1 (70% vs 56%; $p < 0.001$), while the proportion of patients with CHD significantly increased across eras (49% to 54%; $p = 0.018$). The median time on the waitlist increased from 68 days to 78 days ($p = 0.005$). There were no significant differences across eras in the cumulative incidence of death on the waitlist among all candidates (subdistribution hazard ratio, 0.96; 95% CI, 0.80 to 1.14; $p = 0.63$) and among those listed status 1A (subdistribution hazard ratio, 1.16; 95% CI, 0.95 to 1.41; $p = 0.14$). Graft survival at 90 days was also similar across eras in the overall population and in those with CHD ($p > 0.53$ for both).

As a consequence, aggressive treatment of heart failure has been emphasized in recent guidelines. Prognostic criteria have been investigated to identify patients who have truly exhausted medical therapy and thus are likely to derive the maximum benefit for heart transplantation. Maximal oxygen consumption

(VO₂max), which is measured during maximal exercise, is a measure suggested as a critical objective criterion of the functional reserve of the heart. The American College of Cardiology and American Heart Association have adopted VO₂max as a criterion for patient selection.⁹ Studies have suggested that transplantation can be safely deferred in those patients with a VO₂max greater than 14 mL/kg/min. The importance of VO₂max has also been emphasized by the American Heart Association when addressing heart transplant candidacy.¹⁰ In past years, a left ventricular ejection fraction of less than 20% or a New York Heart Association class III or IV status might have been used to determine transplant candidacy. However, as indicated by the American College of Cardiology criteria, these measurements are no longer considered adequate to identify transplant candidates. These measurements may be used to identify patients for further cardiovascular workup but should not be the sole criteria for transplant.

Methods other than VO₂max have been proposed as predictive models in adults.^{11,12,13,14} The Heart Failure Survival Scale and the Seattle Heart Failure Model (SHFM) are examples. In particular, the SHFM provides an estimate of 1-, 2-, and 3-year survival with the use of routinely obtained clinical and laboratory data. Information on pharmacologic and device usage is incorporated into the model, permitting some estimation on the effects of current, more aggressive heart failure treatment strategies. Levy et al (2006) introduced the model using a multivariate analysis of data from the Prospective Randomized Amlodipine Survival Evaluation-1 heart failure trial (n=1 125).¹⁵ Applied to the data of 5 other heart failure trials, SHFM correlated well with actual survival (*r*=0.98). SHFM has been validated in both ambulatory and hospitalized heart failure populations,^{16,17,18} but with a noted underestimation of mortality risk, particularly in blacks and device recipients.^{19,20} None of these models has been universally adopted by transplant centers.

Summary

A heart transplant and a retransplant consists of replacing a diseased heart with a healthy donor heart. Transplantation is used for patients with refractory end-stage cardiac disease.

For individuals who have end-stage heart failure who receive a heart transplant, the evidence includes retrospective studies and registry data. Relevant outcomes are OS, symptoms, and morbid events. Heart transplant remains a viable treatment for those with severe heart dysfunction despite appropriate medical management with medication, surgery, or medical devices. Given the exceedingly poor survival rates without transplantation for these patients, evidence of post-transplant survival is sufficient to demonstrate that heart transplantation provides a survival benefit. Heart transplantation is contraindicated for patients for whom the procedure is expected to be futile due to comorbid disease or for whom posttransplantation care is expected to worsen comorbid conditions significantly. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have had a prior heart transplant complicated by graft failure or severe dysfunction of the heart who receive a heart retransplant, the evidence includes systematic reviews, retrospective studies, and registry data. Relevant outcomes are OS, symptoms, and morbid events. Despite improvements in the prognosis for many patients with graft failure, cardiac allograft vasculopathy, and severe dysfunction of the transplanted heart, heart retransplant remains a viable treatment for those whose severe symptoms persist despite treatment with other medical or surgical remedies. Given the exceedingly poor survival rates without retransplantation for patients who have exhausted other treatments, evidence of post-transplant survival is sufficient to demonstrate that heart retransplantation provides a survival benefit in appropriately selected patients. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Policy History

Date	Action
9/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.

10/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
10/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
10/2018	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2018	Clarified coding information.
11/2017	Annual policy review. New references added.
1/2016	Annual policy review. New references added.
11/2015	Added coding language.
12/2014	Annual policy review. New references added.
10/2014	Medical policy remediation: New indications for non-coverage. Coding information clarified. Effective 10/1/2014.
6/2014	Updated Coding section with ICD10 procedure and diagnosis codes, effective 10/2015.
4/2014	Annual policy review. New medically necessary and investigational indications described. Effective 4/1/2014.
12/2013	Removed ICD-9 diagnosis codes as this policy requires prior authorization
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
10/2011	Medical Policy Group - GI, Nutrition and Organ Transplantation. No changes to policy statements.
3/22/2011	Clarified medical necessity criteria based on Annual policy review.
11/2010	Medical Policy Group - Gastroenterology, Nutrition and Organ Transplantation. No changes to policy statements.
5/20/2010	Updated to clarify and reword when services are not covered section. No changes to policy statement.
3/2010	National Policy Review # 7.03.09. Revision to policy statement.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Black CK, Termanini KM, Aguirre O, et al. Solid organ transplantation in the 21 st century. Ann Transl Med. Oct 2018; 6(20): 409. PMID 30498736
2. United Network for Organ Sharing (UNOS). Transplant trends. 2021; <https://unos.org/data/transplant-trends/>. Accessed June 14, 2021
3. Organ Procurement and Transplantation Network (OPTN). National data. 2021; <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/>. Accessed June 14, 2021
4. Lietz K, Miller LW. Improved survival of patients with end-stage heart failure listed for heart transplantation: analysis of organ procurement and transplantation network/U.S. United Network of Organ Sharing data, 1990 to 2005. J Am Coll Cardiol. Sep 25 2007; 50(13): 1282-90. PMID 17888847
5. Johnson MR, Meyer KH, Haft J, et al. Heart transplantation in the United States, 1999-2008. Am J Transplant. Apr 2010; 10(4 Pt 2): 1035-46. PMID 20420651

6. Bakhtiyar SS, Godfrey EL, Ahmed S, et al. Survival on the Heart Transplant Waiting List. *JAMA Cardiol.* Nov 01 2020; 5(11): 1227-1235. PMID 32785619
7. Alshawabkeh L, Opotowsky AR, Carter KD, et al. Disparities in Wait-List Outcomes for Adults With Congenital Heart Disease Listed for Heart Transplantation Before and Since Revision of Status I Listing. *Am J Cardiol.* Nov 15 2018; 122(10): 1761-1764. PMID 30236623
8. Alshawabkeh LI, Hu N, Carter KD, et al. Wait-List Outcomes for Adults With Congenital Heart Disease Listed for Heart Transplantation in the U.S. *J Am Coll Cardiol.* Aug 30 2016; 68(9): 908-17. PMID 27561764
9. Magnetta DA, Godown J, West S, et al. Impact of the 2016 revision of US Pediatric Heart Allocation Policy on waitlist characteristics and outcomes. *Am J Transplant.* Dec 2019; 19(12): 3276-3283. PMID 31544351
10. Hunt SA, Abraham WT, Chin MH, et al. ACC/AHA 2005 Guideline Update for the Diagnosis and Management of Chronic Heart Failure in the Adult: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Update the 2001 Guidelines for the Evaluation and Management of Heart Failure): developed in collaboration with the American College of Chest Physicians and the International Society for Heart and Lung Transplantation: endorsed by the Heart Rhythm Society. *Circulation.* Sep 20 2005; 112(12): e154-235. PMID 16160202
11. Costanzo MR, Augustine S, Bourge R, et al. Selection and treatment of candidates for heart transplantation. A statement for health professionals from the Committee on Heart Failure and Cardiac Transplantation of the Council on Clinical Cardiology, American Heart Association. *Circulation.* Dec 15 1995; 92(12): 3593-612. PMID 8521589
12. Aaronson KD, Schwartz JS, Chen TM, et al. Development and prospective validation of a clinical index to predict survival in ambulatory patients referred for cardiac transplant evaluation. *Circulation.* Jun 17 1997; 95(12): 2660-7. PMID 9193435
13. Alla F, Briancon S, Juilliere Y, et al. Differential clinical prognostic classifications in dilated and ischemic advanced heart failure: the EPICAL study. *Am Heart J.* May 2000; 139(5): 895-904. PMID 10783225
14. Hansen A, Haass M, Zugck C, et al. Prognostic value of Doppler echocardiographic mitral inflow patterns: implications for risk stratification in patients with chronic congestive heart failure. *J Am Coll Cardiol.* Mar 15 2001; 37(4): 1049-55. PMID 11263607
15. Lee DS, Austin PC, Rouleau JL, et al. Predicting mortality among patients hospitalized for heart failure: derivation and validation of a clinical model. *JAMA.* Nov 19 2003; 290(19): 2581-7. PMID 14625335
16. Levy WC, Mozaffarian D, Linker DT, et al. The Seattle Heart Failure Model: prediction of survival in heart failure. *Circulation.* Mar 21 2006; 113(11): 1424-33. PMID 16534009
17. Gorodeski EZ, Chu EC, Chow CH, et al. Application of the Seattle Heart Failure Model in ambulatory patients presented to an advanced heart failure therapeutics committee. *Circ Heart Fail.* Nov 2010; 3(6): 706-14. PMID 20798278
18. Ketchum ES, Moorman AJ, Fishbein DP, et al. Predictive value of the Seattle Heart Failure Model in patients undergoing left ventricular assist device placement. *J Heart Lung Transplant.* Sep 2010; 29(9): 1021-5. PMID 20558086
19. Nutter AL, Tanawuttiwat T, Silver MA. Evaluation of 6 prognostic models used to calculate mortality rates in elderly heart failure patients with a fatal heart failure admission. *Congest Heart Fail.* Sep-Oct 2010; 16(5): 196-201. PMID 20887615
20. Kalogeropoulos AP, Georgiopoulou VV, Giamouzis G, et al. Utility of the Seattle Heart Failure Model in patients with advanced heart failure. *J Am Coll Cardiol.* Jan 27 2009; 53(4): 334-42. PMID 19161882
21. May HT, Horne BD, Levy WC, et al. Validation of the Seattle Heart Failure Model in a community-based heart failure population and enhancement by adding B-type natriuretic peptide. *Am J Cardiol.* Aug 15 2007; 100(4): 697-700. PMID 17697831
22. Virani SS, Alonso A, Aparicio HJ, et al. Heart Disease and Stroke Statistics-2021 Update: A Report From the American Heart Association. *Circulation.* Feb 23 2021; 143(8): e254-e743. PMID 33501848

23. Lund LH, Edwards LB, Dipchand AI, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-third Adult Heart Transplantation Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016; 35(10): 1158-1169. PMID 27772668
24. Jaramillo N, Segovia J, Gomez-Bueno M, et al. Characteristics of patients with survival longer than 20 years following heart transplantation. *Rev Esp Cardiol*. Oct 2013;66(10):797-802. PMID 23932221
25. Nguyen VP, Mahr C, Mokadam NA, et al. The Benefit of Donor-Recipient Matching for Patients Undergoing Heart Transplantation. *J Am Coll Cardiol*. Apr 04 2017; 69(13): 1707-1714. PMID 28359517
26. Rana A, Gruessner A, Agopian VG, et al. Survival benefit of solid-organ transplant in the United States. *JAMA Surg*. Mar 01 2015; 150(3): 252-9. PMID 25629390
27. Kilic A, Weiss ES, George TJ, et al. What predicts long-term survival after heart transplantation? An analysis of 9,400 ten-year survivors. *Ann Thorac Surg*. Mar 2012; 93(3): 699-704. PMID 22226494
28. Dipchand AI, Kirk R, Mahle WT, et al. Ten yr of pediatric heart transplantation: a report from the Pediatric Heart Transplant Study. *Pediatr Transplant*. Mar 2013; 17(2): 99-111. PMID 23442098
29. Auerbach SR, Richmond ME, Chen JM, et al. Multiple risk factors before pediatric cardiac transplantation are associated with increased graft loss. *Pediatr Cardiol*. Jan 2012; 33(1): 49-54. PMID 21892650
30. Rossano JW, Dipchand AI, Edwards LB, et al. The Registry of the International Society for Heart and Lung Transplantation: Nineteenth Pediatric Heart Transplantation Report-2016; Focus Theme: Primary Diagnostic Indications for Transplant. *J Heart Lung Transplant*. Oct 2016; 35(10): 1185-1195. PMID 27772670
31. Savla J, Lin KY, Lefkowitz DS, et al. Adolescent age and heart transplantation outcomes in myocarditis or congenital heart disease. *J Heart Lung Transplant*. Sep 2014; 33(9): 943-9. PMID 24929645
32. Almond CSD, Thiagarajan RR, Piercey GE, et al. Waiting list mortality among children listed for heart transplantation in the United States. *Circulation*. Feb 10 2009; 119(5): 717-727. PMID 19171850
33. Tjang YS, Tenderich G, Hornik L, et al. Cardiac retransplantation in adults: an evidence-based systematic review. *Thorac Cardiovasc Surg*. Sep 2008; 56(6): 323-7. PMID 18704853
34. Saito A, Novick RJ, Kiaii B, et al. Early and late outcomes after cardiac retransplantation. *Can J Surg*. Feb 2013; 56(1): 21-6. PMID 23187039
35. Miller RJH, Clarke BA, Howlett JG, et al. Outcomes in patients undergoing cardiac retransplantation: A propensity matched cohort analysis of the UNOS Registry. *J Heart Lung Transplant*. Oct 2019; 38(10): 1067-1074. PMID 31378576
36. Goldraich LA, Stehlik J, Kucheryavaya AY, et al. Retransplant and Medical Therapy for Cardiac Allograft Vasculopathy: International Society for Heart and Lung Transplantation Registry Analysis. *Am J Transplant*. Jan 2016; 16(1): 301-9. PMID 26274617
37. Belli E, Leoni Moreno JC, Hosenpud J, et al. Preoperative risk factors predict survival following cardiac retransplantation: analysis of the United Network for Organ Sharing database. *J Thorac Cardiovasc Surg*. Jun 2014; 147(6): 1972-7. PMID 24636155
38. Friedland-Little JM, Gajarski RJ, Yu S, et al. Outcomes of third heart transplants in pediatric and young adult patients: analysis of the United Network for Organ Sharing database. *J Heart Lung Transplant*. Sep 2014; 33(9): 917-23. PMID 24861821
39. Bock MJ, Nguyen K, Malerba S, et al. Pediatric cardiac retransplantation: Waitlist mortality stratified by age and era. *J Heart Lung Transplant*. Apr 2015; 34(4): 530-7. PMID 25016920
40. Conway J, Manlhiot C, Kirk R, et al. Mortality and morbidity after retransplantation after primary heart transplant in childhood: an analysis from the registry of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. Mar 2014; 33(3): 241-51. PMID 24462559
41. Mistiaen WP. Heart transplantation in patients with previous malignancy. An overview. *Acta Cardiol*. Apr 2015; 70(2): 123-30. PMID 26148371
42. Yoosabai A, Mehta A, Kang W, et al. Pretransplant malignancy as a risk factor for posttransplant malignancy after heart transplantation. *Transplantation*. Feb 2015; 99(2): 345-50. PMID 25606783

43. Oliveira GH, Hardaway BW, Kucheryavaya AY, et al. Characteristics and survival of patients with chemotherapy-induced cardiomyopathy undergoing heart transplantation. *J Heart Lung Transplant.* Aug 2012; 31(8): 805-10. PMID 22551930
44. Sigurdardottir V, Bjortuft O, Eiskjaer H, et al. Long-term follow-up of lung and heart transplant recipients with pre-transplant malignancies. *J Heart Lung Transplant.* Dec 2012; 31(12): 1276-80. PMID 23089300
45. Agüero F, Castel MA, Cocchi S, et al. An Update on Heart Transplantation in Human Immunodeficiency Virus-Infected Patients. *Am J Transplant.* Jan 2016; 16(1): 21-8. PMID 26523614
46. Uriel N, Jorde UP, Cotarlan V, et al. Heart transplantation in human immunodeficiency virus-positive patients. *J Heart Lung Transplant.* Jul 2009; 28(7): 667-9. PMID 19560693
47. Doberne JW, Jawitz OK, Raman V, et al. Heart Transplantation Survival Outcomes of HIV Positive and Negative Recipients. *Ann Thorac Surg.* May 2021; 111(5): 1465-1471. PMID 32946847
48. Organ Procurement and Transplantation Network (OPTN). Organ Procurement and Transplantation Network Policies. 2021; <https://optn.transplant.hrsa.gov/governance/policies/>. Accessed June 15, 2021
49. Working Party of the British Transplantation Society. Kidney and Pancreas Transplantation in Patients with HIV. Second Edition (Revised). British Transplantation Society Guidelines. Macclesfield, UK: British Transplantation Society; 2017.
50. Jamil A, Qin H, Felius J, et al. Comparison of Clinical Characteristics, Complications, and Outcomes in Recipients Having Heart Transplants 65 Years of Age Versus 65 Years of Age. *Am J Cardiol.* Dec 15 2017; 120(12): 2207-2212. PMID 29056228
51. Cooper LB, Lu D, Mentz RJ, et al. Cardiac transplantation for older patients: Characteristics and outcomes in the septuagenarian population. *J Heart Lung Transplant.* Mar 2016; 35(3): 362-369. PMID 26632028
52. Awad M, Czer LS, Mirocha J, et al. Similar Mortality and Morbidity of Orthotopic Heart Transplantation for Patients 70 Years of Age and Older Compared With Younger Patients. *Transplant Proc.* Oct 2016; 48(8): 2782-2791. PMID 27788818
53. Kilic A, Weiss ES, Yuh DD, et al. Factors associated with 5-year survival in older heart transplant recipients. *J Thorac Cardiovasc Surg.* Feb 2012; 143(2): 468-74. PMID 22248684
54. De Santo LS, Romano G, Maiello C, et al. Pulmonary artery hypertension in heart transplant recipients: how much is too much?. *Eur J Cardiothorac Surg.* Nov 2012; 42(5): 864-9; discussion 869-70. PMID 22402452
55. Perez-Villa F, Farrero M, Cardona M, et al. Bosentan in heart transplantation candidates with severe pulmonary hypertension: efficacy, safety and outcome after transplantation. *Clin Transplant.* Jan-Feb 2013; 27(1): 25-31. PMID 22861120
56. Pons J, Leblanc MH, Bernier M, et al. Effects of chronic sildenafil use on pulmonary hemodynamics and clinical outcomes in heart transplantation. *J Heart Lung Transplant.* Dec 2012; 31(12): 1281-7. PMID 23127754
57. Bedanova H, Orban M, Vrsansky D, et al. Impact of pulmonary hypertension on early hemodynamics, morbidity and mortality after orthotopic heart transplantation. A single center study. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub.* Mar 2013; 157(1): 35-40. PMID 23073529
58. Tsukashita M, Takayama H, Takeda K, et al. Effect of pulmonary vascular resistance before left ventricular assist device implantation on short- and long-term post-transplant survival. *J Thorac Cardiovasc Surg.* Nov 2015; 150(5): 1352-60, 1361.e1-2. PMID 26253875
59. Arshad A, Kew EP, Lim S. Comparison of Renal Outcomes in Patients With Left Ventricular Assist Device and Heart Transplantation. *Transplant Proc.* Dec 2019; 51(10): 3395-3398. PMID 31810507
60. Kolsrud O, Karason K, Holmberg E, et al. Renal function and outcome after heart transplantation. *J Thorac Cardiovasc Surg.* Apr 2018; 155(4): 1593-1604.e1. PMID 29338859
61. Hong KN, Merlo A, Chauhan D, et al. Evidence supports severe renal insufficiency as a relative contraindication to heart transplantation. *J Heart Lung Transplant.* Jul 2016; 35(7): 893-900. PMID 27105687
62. Goel AN, Iyengar A, Schowengerdt K, et al. Heart transplantation in children with intellectual disability: An analysis of the UNOS database. *Pediatr Transplant.* Mar 2017; 21(2). PMID 27933693

63. Wightman A, Bartlett HL, Zhao Q, et al. Prevalence and outcomes of heart transplantation in children with intellectual disability. *Pediatr Transplant*. Mar 2017; 21(2). PMID 27801533
64. Wightman A, Bradford MC, Hsu E, et al. Prevalence and Long-Term Outcomes of Solid Organ Transplant in Children with Intellectual Disability. *J Pediatr*. Aug 2021; 235: 10-17.e4. PMID 33794218
65. Prendergast C, McKane M, Dodd DA, et al. The impact of cognitive delay on pediatric heart transplant outcomes. *Pediatr Transplant*. Mar 2017; 21(2). PMID 28191755
66. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. Aug 08 2017; 136(6): e137-e161. PMID 28455343
67. Rosenthal D, Chrisant MR, Edens E, et al. International Society for Heart and Lung Transplantation: Practice guidelines for management of heart failure in children. *J Heart Lung Transplant*. Dec 2004; 23(12): 1313-33. PMID 15607659
68. Mehra MR, Canter CE, Hannan MM, et al. The 2016 International Society for Heart Lung Transplantation listing criteria for heart transplantation: A 10-year update. *J Heart Lung Transplant*. Jan 2016; 35(1): 1-23. PMID 26776864
69. Costanzo MR, Dipchand A, Starling R, et al. The International Society of Heart and Lung Transplantation Guidelines for the care of heart transplant recipients. *J Heart Lung Transplant*. Aug 2010; 29(8): 914-56. PMID 20643330
70. Canter CE, Shaddy RE, Bernstein D, et al. Indications for heart transplantation in pediatric heart disease: a scientific statement from the American Heart Association Council on Cardiovascular Disease in the Young; the Councils on Clinical Cardiology, Cardiovascular Nursing, and Cardiovascular Surgery and Anesthesia; and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation*. Feb 06 2007; 115(5): 658-76. PMID 17261651
71. Centers for Medicare & Medicaid Services (CMS). National Coverage Determination (NCD) for HEART TRANSPLANTS (260.9). 2008; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=112&ncdver=3&CoverageSelection=National&Keyword=heart+transplant&KeywordLookUp=Title&KeywordSearchType=And&clickon=search&bc=gAAAABAAAA&>. Accessed June 15, 2021