



## MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

# Medical Policy Occipital Nerve Stimulation

### Table of Contents

- [Policy: Commercial](#)
- [Coding Information](#)
- [Information Pertaining to All Policies](#)
- [Policy: Medicare](#)
- [Description](#)
- [References](#)
- [Authorization Information](#)
- [Policy History](#)

### Policy Number: 237

BCBSA Reference Number: 7.01.125 (For Plans internal use only)  
NCD/LCD: NA

### Related Policies

- Percutaneous Electrical Nerve Stimulation and Percutaneous Neuromodulation Therapy, #[172](#)
- Spinal Cord and Dorsal Root Ganglion Stimulation, #[472](#)

### Policy

#### Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

Occipital nerve stimulation is considered [INVESTIGATIONAL](#) for all indications.

### Prior Authorization Information

#### Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

#### Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is <b>not</b> a covered service.
Commercial PPO and Indemnity	This is <b>not</b> a covered service.
Medicare HMO Blue <sup>SM</sup>	This is <b>not</b> a covered service.
Medicare PPO Blue <sup>SM</sup>	This is <b>not</b> a covered service.

### CPT Codes / HCPCS Codes / ICD Codes

The following codes are included below for informational purposes. Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's

*contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.*

*Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.*

## **CPT Codes**

There is no specific CPT code for this service.

## **ICD Diagnosis Codes**

Investigational for all diagnoses.

## **Description**

### **Headache**

There are 4 types of headache: vascular, muscle contraction (tension), traction, and inflammatory. Primary (not the result of another condition) chronic headache is defined as headache occurring more than 15 days of the month for at least 3 consecutive months. An estimated 45 million Americans experience chronic headaches. For at least half of these people, the problem is severe and sometimes disabling. Herein, we only discuss types of vascular headache, including migraine, hemicrania continua, and cluster.

### **Migraine**

Migraine is the most common type of vascular headache. Migraine headaches are usually characterized by severe pain on one or both sides of the head, an upset stomach, and, at times, disturbed vision. One-year prevalence of migraine ranges from 6% to 15% in adult men and from 14% to 35% in adult women. Migraine headaches may last a day or more and can strike as often as several times a week or as rarely as once every few years.

### **Treatment of Migraine**

Drug therapy for migraine is often combined with biofeedback and relaxation training. Sumatriptan and other triptans are commonly used for relief of symptoms. Drugs used to prevent migraine include amitriptyline, propranolol and other  $\beta$ -blockers, topiramate and other antiepileptic drugs, and verapamil.

### **Hemicrania Continua**

Hemicrania continua causes moderate and occasionally severe pain on only one side of the head. At least one of the following symptoms must also occur: conjunctival injection and/or lacrimation, nasal congestion and/or rhinorrhea, or ptosis, and/or miosis. Headache occurs daily and is continuous with no pain-free periods. Hemicrania continua occurs mainly in women, and its true prevalence is not known.

### **Treatment of Hemicrania Continua**

Indomethacin usually provides rapid relief of symptoms. Other nonsteroidal anti-inflammatory drugs, including ibuprofen, celecoxib, and naproxen, can provide some relief of symptoms. Amitriptyline and other tricyclic antidepressants are effective in some patients.

### **Cluster Headache**

Cluster headache occurs in cyclical patterns or clusters of severe or very severe unilateral orbital or supraorbital and/or temporal pain. The headache is accompanied by at least one of the following autonomic symptoms: ptosis, conjunctival injection, lacrimation, rhinorrhea, and, less commonly, facial blushing, swelling, or sweating. Bouts of 1 headache every other day up to 8 attacks per day may last from weeks to months, usually followed by remission periods when the headache attacks stop completely. The pattern varies by person, but most people have 1 or 2 cluster periods a year. During remission, no headaches occur for months, and sometimes even years. The intense pain is caused by the dilation of blood vessels, which creates pressure on the trigeminal nerve. While this process is the

immediate cause of the pain, the etiology is not fully understood. It is more common in men than in woman. One-year prevalence is estimated to be 0. to 1. in 1000.

### Treatment of Cluster Headache

Management of cluster headache consists of abortive and preventive treatment. Abortive treatments include subcutaneous injection of sumatriptan, topical anesthetics sprayed into the nasal cavity, and strong coffee. Some patients respond to rapidly inhaled pure oxygen. A variety of other pharmacologic and behavioral methods of aborting and preventing attacks have been reported with wide variation in patient response.

### Peripheral Nerve Stimulators

Implanted peripheral nerve stimulators have been used to treat refractory pain for many years but have only recently been proposed to manage craniofacial pain. Occipital, supraorbital, and infraorbital stimulation have been reported in the literature.

### Summary

Occipital nerve stimulation delivers a small electrical charge to the occipital nerve intended to prevent migraines and other headaches in patients who have not responded to medications. The device consists of a subcutaneously implanted pulse generator (in the chest wall or abdomen) attached to extension leads that are tunneled to join electrodes placed across one or both occipital nerves at the base of the skull. Continuous or intermittent stimulation may be used.

For individuals who have migraine headaches refractory to preventive medical management who receive occipital nerve stimulation, the evidence includes randomized controlled trials (RCTs), systematic reviews of RCTs, and observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Systematic reviews identified 5 sham-controlled randomized trials. Findings from pooled analyses of these RCTs were mixed. For example, compared with placebo, response rates to occipital nerve stimulation did not differ significantly but did reduce the number of days with prolonged moderate-to-severe headache. Occipital nerve stimulation was also associated with a substantial number of minor and serious adverse events. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have non-migraine headaches (eg, hemicrania continua, cluster headaches) who receive occipital nerve stimulation, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Many of the case series had small sample sizes; series with over 25 patients were available only for treatment of cluster headache. Although the case series tended to find that a substantial number of patients improved after occipital nerve stimulation, these studies lacked blinding and comparison groups. RCTs are needed to compare outcomes between occipital nerve stimulation and comparators (eg, to control for a potential placebo effect). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### Policy History

Date	Action
6/2022	Annual policy review. Policy statements unchanged.
5/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2017	Annual policy review. New references added.

5/2016	Annual policy review. New references added.
12/2014	Annual policy review. New references added.
2/2014	Annual policy review. New references added.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
4/21/2011	References updated.
1/2011	Medical Policy Group – Neurology and Neurosurgery, No changes to policy statements.
11/1/10	Medical Policy 237 created.

## Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

## References

1. Chen YF, Bramley G, Unwin G, et al. Occipital nerve stimulation for chronic migraine--a systematic review and meta-analysis. *PLoS One*. 2015; 10(3): e0116786. PMID 25793740
2. Yang Y, Song M, Fan Y, et al. Occipital Nerve Stimulation for Migraine: A Systematic Review. *Pain Pract*. Apr 2016; 16(4): 509-17. PMID 25865962
3. Saper JR, Dodick DW, Silberstein SD, et al. Occipital nerve stimulation for the treatment of intractable chronic migraine headache: ONSTIM feasibility study. *Cephalalgia*. Feb 2011; 31(3): 271-85. PMID 20861241
4. Silberstein SD, Dodick DW, Saper J, et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: results from a randomized, multicenter, double-blinded, controlled study. *Cephalalgia*. Dec 2012; 32(16): 1165-79. PMID 23034698
5. Dodick DW, Silberstein SD, Reed KL, et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: long-term results from a randomized, multicenter, double-blinded, controlled study. *Cephalalgia*. Apr 2015; 35(4): 344-58. PMID 25078718
6. Burns B, Watkins L, Goadsby PJ. Treatment of hemicrania continua by occipital nerve stimulation with a bion device: long-term follow-up of a crossover study. *Lancet Neurol*. Nov 2008; 7(11): 1001-12. PMID 18845482
7. Burns B, Watkins L, Goadsby PJ. Treatment of intractable chronic cluster headache by occipital nerve stimulation in 14 patients. *Neurology*. Jan 27 2009; 72(4): 341-5. PMID 19171831
8. Magis D, Gerardy PY, Remeacle JM, et al. Sustained effectiveness of occipital nerve stimulation in drug-resistant chronic cluster headache. *Headache*. Sep 2011; 51(8): 1191-201. PMID 21848953
9. Mueller OM, Gaul C, Katsarava Z, et al. Occipital nerve stimulation for the treatment of chronic cluster headache - lessons learned from 18 months experience. *Cent Eur Neurosurg*. May 2011; 72(2): 84-9. PMID 21448856
10. Fontaine D, Blond S, Lucas C, et al. Occipital nerve stimulation improves the quality of life in medically-intractable chronic cluster headache: Results of an observational prospective study. *Cephalalgia*. Oct 2017; 37(12): 1173-1179. PMID 27697849
11. Leone M, Proietti Cecchini A, Messina G, et al. Long-term occipital nerve stimulation for drug-resistant chronic cluster headache. *Cephalalgia*. Jul 2017; 37(8): 756-763. PMID 27250232
12. Miller S, Watkins L, Matharu M. Treatment of intractable chronic cluster headache by occipital nerve stimulation: a cohort of 51 patients. *Eur J Neurol*. Feb 2017; 24(2): 381-390. PMID 27995704
13. Leplus A, Fontaine D, Donnet A, et al. Long-Term Efficacy of Occipital Nerve Stimulation for Medically Intractable Cluster Headache. *Neurosurgery*. Jan 13 2021; 88(2): 375-383. PMID 32985662

14. Vadivelu S, Bolognese P, Milhorat TH, et al. Occipital nerve stimulation for refractory headache in the Chiari malformation population. *Neurosurgery*. Jun 2012; 70(6): 1430-6; discussion 1436-7. PMID 22418582
15. Sweet JA, Mitchell LS, Narouze S, et al. Occipital Nerve Stimulation for the Treatment of Patients With Medically Refractory Occipital Neuralgia: Congress of Neurological Surgeons Systematic Review and Evidence-Based Guideline. *Neurosurgery*. Sep 2015; 77(3): 332-41. PMID 26125672
16. National Institute for Health and Care Excellence. Occipital nerve stimulation for intractable chronic migraine [IPG452]. 2013; <https://www.nice.org.uk/guidance/ipg452>. Accessed March 10, 2021.