



Medical Policy

Cardiac Hemodynamic Monitoring for the Management of Heart Failure in the Outpatient Setting

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 287

BCBSA Reference Number: 2.02.24 (For Plan internal use only)

Related Policies

Biventricular Pacemakers (Cardiac Resynchronization Therapy) for the Treatment of Heart Failure, #[101](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Ambulatory Care and Outpatient Setting

In the ambulatory care and outpatient setting, cardiac hemodynamic monitoring for the management of heart failure using implantable direct pressure monitoring of the pulmonary artery, thoracic bioimpedance, inert gas rebreathing, and arterial pressure during the Valsalva maneuver, is considered [INVESTIGATIONAL](#). (This includes inpatient placement of the device.)

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service in the outpatient setting. This is not a covered service in the inpatient setting.
Commercial PPO and Indemnity	This is not a covered service in the outpatient setting. This is not a covered service in the inpatient setting.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine

coverage or non-coverage as it applies to an individual member. A draft of future ICD-10 Coding related to this document, as it might look today, is included below for your reference.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following CPT and HCPCS codes are considered investigational for Commercial Members: Managed Care (HMO and POS), PPO and Indemnity:

CPT Codes

CPT codes:	Code Description
33289	Transcatheter implantation of wireless pulmonary artery pressure sensor for long-term hemodynamic monitoring, including deployment and calibration of the sensor, right heart catheterization, selective pulmonary catheterization, radiological supervision and interpretation, and pulmonary artery angiography, when performed
93264	Remote monitoring of a wireless pulmonary artery pressure sensor for up to 30 days, including at least weekly downloads of pulmonary artery pressure recordings, interpretation(s), trend analysis, and report(s) by a physician or other qualified health care professional
93701	Bioimpedance-derived physiologic cardiovascular analysis.

HCPCS Codes

HCPCS codes:	Code Description
C2624	Implantable wireless pulmonary artery pressure sensor with delivery catheter, including all system components

Description

Chronic Heart Failure

Patients with chronic heart failure are at risk of developing acute decompensated heart failure, often requiring hospital admission. Patients with a history of acute decompensation have the additional risk of future episodes of decompensation and death. Reasons for the transition from a stable, chronic state to an acute, decompensated state include disease progression, as well as acute events such as coronary ischemia and dysrhythmias. While precipitating factors are frequently not identified, the most common preventable cause is noncompliance with medication and dietary regimens.¹

Management

Strategies for reducing decompensation, and thus the need for hospitalization, are aimed at early identification of patients at risk for imminent decompensation. Programs for early identification of heart failure are characterized by frequent contact with patients to review signs and symptoms with a health care provider, education, and medication adjustments as appropriate. These encounters may occur face-to-face in the office or at home, or via cellular or computed technology.²

Precise measurement of cardiac hemodynamics is often employed in the intensive care setting to carefully manage fluid status in acutely decompensated heart failure. Transthoracic echocardiography, transesophageal echocardiography, and Doppler ultrasound are noninvasive methods for monitoring cardiac output on an intermittent basis for the more stable patient but are not addressed herein. A variety of biomarkers and radiologic techniques may be used for dyspnea when the diagnosis of acute decompensated heart failure is uncertain.

The criterion standard for hemodynamic monitoring is pulmonary artery (PA) catheters and central venous pressure catheters. However, they are invasive, inaccurate, and inconsistent in predicting fluid responsiveness. Several studies have demonstrated that catheters fail to improve outcomes in critically ill

patients and may be associated with harm. To overcome these limitations, multiple techniques and devices have been developed that use complex imaging technology and computer algorithms to estimate fluid responsiveness, volume status, cardiac output and tissue perfusion. Many are intended for use in outpatient settings but can be used in the emergency department, intensive care unit, and operating room. Four methods are reviewed here: implantable pressure monitoring devices, thoracic bioimpedance, inert gas rebreathing, and arterial waveform during the Valsalva maneuver. Use of the last 3 is not widespread because of several limitations including use of proprietary technology making it difficult to confirm their validity and lack of large randomized controlled trials to evaluate treatment decisions guided by these hemodynamic monitors.

Summary

Description

A variety of outpatient cardiac hemodynamic monitoring devices are intended to improve quality of life and reduce morbidity for patients with heart failure by decreasing episodes of acute decompensation. Monitors can identify physiologic changes that precede clinical symptoms and thus allow preventive intervention. These devices operate through various mechanisms, including implantable pressure sensors, thoracic bioimpedance measurement, inert gas rebreathing, and estimation of left ventricular end-diastolic pressure by arterial pressure during the Valsalva maneuver.

Summary of Evidence

For individuals with New York Heart Association (NYHA) class II-IV heart failure in outpatient settings who have had a hospitalization in the past year and/or have elevated natriuretic peptides who receive hemodynamic monitoring with an implantable pulmonary artery pressure sensor device, the evidence includes 2 meta-analyses, randomized controlled trials (RCTs), and nonrandomized studies. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. One implantable pressure monitor, the CardioMEMS device, has U.S. Food and drug Administration (FDA) approval. The pivotal CHAMPION RCT reported a statistically significant 28% decrease in heart failure hospitalization (HFH) in patients implanted with the CardioMEMS device compared with usual care. However, trial results were potentially biased in favor of the treatment group due to the use of additional nurse communication to enhance protocol compliance with the device. The manufacturer conducted multiple analyses to address potential bias from the nurse interventions. Results were reviewed favorably by the FDA. While these analyses demonstrated the consistency of benefit of the CardioMEMS device, all such analyses have methodologic limitations. Early safety data have been suggestive of a higher rate of procedural complications, particularly related to pulmonary artery injury. While the U.S. CardioMEMS post-approval study and CardioMEMS European Monitoring Study for Heart Failure (MEMS-HF) study reported a significant decrease in HFH with few device- or system-related complications at 1 year, the impact of nursing interventions remains unclear. The subsequent GUIDE-HF RCT failed to meet its primary efficacy endpoint, the composite of HFH, urgent heart failure visits, and death at 1 year. With the approval of the FDA, the statistical analysis plan was updated to pre-specify sensitivity analyses to assess the impact of COVID-19 on the trial. For the 72% of patients who completed follow-up prior to the public health emergency declaration in March 2020, a statistically significant 19% reduction in the primary endpoint was reported, driven by a 28% reduction in HFH. However, lifestyle changes during the COVID-19 pandemic such as changes in physical activity, exposure to infections, willingness to seek medical care, and adherence to medications are unmeasured and add imprecision to treatment effect estimates, as do alterations in provider behaviors. Enrollment of NYHA Class II patients were significantly enriched in the first 500 patients, potentially impacting the pre-COVID-19 analysis. The MONITOR-HF trial, an open-label RCT conducted in the Netherlands, showed that hemodynamic monitoring significantly improved quality of life on the Kansas City Cardiomyopathy Questionnaire (KCCQ) and reduced HFH but did not impact mortality at 1 year follow-up. Overall, the beneficial effect of CardioMEMS, if any, appears to be on the hospitalization outcome of the composite. Both urgent heart failure visits and death outcomes had hazard ratios favoring the control group with wide confidence intervals including the null value in pre-COVID-19, during-COVID-19, and overall analyses of the GUIDE-HF trial. The MONITOR-HF trial found improvement in quality of life on the KCCQ for the CardioMEMS group relative to the control, but no significant differences were observed in secondary quality of life and functional status outcomes in the other included trials. While the HFH reduction of 28% found in

the pre-COVID-19 analysis is consistent with findings from the CHAMPION trial, it is unclear whether physician knowledge of treatment assignment biases the decision to hospitalize and administer intravenous diuretics. The two included meta-analyses showed a reduction in HFHs with hemodynamic monitoring in heart failure patients but had discordant findings regarding the impact on mortality. One meta-analysis found no pooled difference in mortality between hemodynamic monitoring and control groups; however, a patient-level meta-analysis revealed a significant 25% decrease in mortality associated with hemodynamic monitoring in patients with heart failure with reduced ejection fraction. Given that the intervention is invasive and intended to be used for a highly prevalent condition and, in light of the conflicting evidence of benefit on mortality and functional outcomes, the lack of periprocedural safety data, and unclear impact of COVID-19 on remote monitoring in the GUIDE-HF trial, the net benefit of the CardioMEMS device remains uncertain. Concerns may be clarified by the ongoing open access phase of the GUIDE-HF RCT and the German non-industry-sponsored PASSPORT-HF trial. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring by thoracic bioimpedance, the evidence includes uncontrolled prospective studies and case series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. There is a lack of RCT evidence evaluating whether the use of these technologies improves health outcomes over standard active management of heart failure patients. The case series have reported physiologic measurement-related outcomes and/or associations between monitoring information and heart failure exacerbations, but do not provide definitive evidence on device efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring with inert gas rebreathing, no studies have been identified on clinical validity or clinical utility. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring of arterial pressure during the Valsalva maneuver, a single study was identified. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. The study assessed the use of left ventricular end-diastolic pressure (LVEDP) monitoring and reported an 85% sensitivity and an 80% specificity to detect LVEDP greater than 15 mm Hg. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
8/2024	Annual policy review. Policy updated with literature review through May 3, 2024; references added. Policy statement unchanged.
8/2023	Annual policy review. References added. Policy statement unchanged.
9/2022	Annual policy review. References added. Policy statement unchanged.
12/2021	Policy clarified to include that placement of implantable cardiac hemodynamic devices in the inpatient setting is considered investigational.
6/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
7/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2019	Clarified coding information.

7/2018	Annual policy review. Policy statement clarified. 7/1/2018
6/2017	Annual policy review. New references added.
7/2016	Annual policy review. New references added.
8/2015	Annual policy review. New references added.
1/2015	Clarified coding information.
10/2014	Clarified coding information.
9/2014	Annual policy review. New references added.
6/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
5/2014	Annual policy review. New references added.
10/2013	Annual policy review. New references added.
5/2013	Annual policy review. New references added.
11/2011-4/2012	Medical policy ICD10 remediation: Formatting, editing and coding updates. No changes to policy statements.
4/2011	Reviewed - Medical Policy Group - Cardiology and Pulmonology. No changes to policy statements.
11/10/2011	Medical policy #287. Effective 11/10/2011.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Opasich C, Rapezzi C, Lucci D, et al. Precipitating factors and decision-making processes of short-term worsening heart failure despite "optimal" treatment (from the IN-CHF Registry). *Am J Cardiol.* Aug 15 2001; 88(4): 382-7. PMID 11545758
2. McAlister FA, Stewart S, Ferrua S, et al. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. *J Am Coll Cardiol.* Aug 18 2004; 44(4): 810-9. PMID 15312864
3. Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): CardioMEMS HF System. 2014; https://www.accessdata.fda.gov/cdrh_docs/pdf10/P100045b.pdf. Accessed May 5, 2024.
4. Loh JP, Barbash IM, Waksman R. Overview of the 2011 Food and Drug Administration Circulatory System Devices Panel of the Medical Devices Advisory Committee Meeting on the CardioMEMS Champion Heart Failure Monitoring System. *J Am Coll Cardiol.* Apr 16 2013; 61(15): 1571-6. PMID 23352783
5. Burns DJP, Arora J, Okunade O, et al. International Consortium for Health Outcomes Measurement (ICHOM): Standardized Patient-Centered Outcomes Measurement Set for Heart Failure Patients. *JACC Heart Fail.* Mar 2020; 8(3): 212-222. PMID 31838032
6. Zannad F, Garcia AA, Anker SD, et al. Clinical outcome endpoints in heart failure trials: a European Society of Cardiology Heart Failure Association consensus document. *Eur J Heart Fail.* Oct 2013; 15(10): 1082-94. PMID 23787718
7. Givertz MM, Stevenson LW, Costanzo MR, et al. Pulmonary Artery Pressure-Guided Management of Patients With Heart Failure and Reduced Ejection Fraction. *J Am Coll Cardiol.* Oct 10 2017; 70(15): 1875-1886. PMID 28982501
8. Curtain JP, Lee MMY, McMurray JJ, et al. Efficacy of implantable haemodynamic monitoring in heart failure across ranges of ejection fraction: a systematic review and meta-analysis. *Heart.* May 15 2023; 109(11): 823-831. PMID 36522146

9. Adamson PB, Abraham WT, Bourge RC, et al. Wireless pulmonary artery pressure monitoring guides management to reduce decompensation in heart failure with preserved ejection fraction. *Circ Heart Fail.* Nov 2014; 7(6): 935-44. PMID 25286913
10. Adamson PB, Abraham WT, Stevenson LW, et al. Pulmonary Artery Pressure-Guided Heart Failure Management Reduces 30-Day Readmissions. *Circ Heart Fail.* Jun 2016; 9(6). PMID 27220593
11. Krahnke JS, Abraham WT, Adamson PB, et al. Heart failure and respiratory hospitalizations are reduced in patients with heart failure and chronic obstructive pulmonary disease with the use of an implantable pulmonary artery pressure monitoring device. *J Card Fail.* Mar 2015; 21(3): 240-9. PMID 25541376
12. Assmus B, Angermann CE, Alkhlout B, et al. Effects of remote haemodynamic-guided heart failure management in patients with different subtypes of pulmonary hypertension: insights from the MEMS-HF study. *Eur J Heart Fail.* Dec 2022; 24(12): 2320-2330. PMID 36054647
13. Heywood JT, Jermyn R, Shavelle D, et al. Impact of Practice-Based Management of Pulmonary Artery Pressures in 2000 Patients Implanted With the CardioMEMS Sensor. *Circulation.* Apr 18 2017; 135(16): 1509-1517. PMID 28219895
14. Iaconelli A, Pellicori P, Caiazzo E, et al. Implanted haemodynamic telemonitoring devices to guide management of heart failure: a review and meta-analysis of randomised trials. *Clin Res Cardiol.* Aug 2023; 112(8): 1007-1019. PMID 36241896
15. Lindenfeld J, Costanzo MR, Zile MR, et al. Implantable Hemodynamic Monitors Improve Survival in Patients With Heart Failure and Reduced Ejection Fraction. *J Am Coll Cardiol.* Feb 13 2024; 83(6): 682-694. PMID 38325994
16. Abraham WT, Adamson PB, Bourge RC, et al. Wireless pulmonary artery haemodynamic monitoring in chronic heart failure: a randomised controlled trial. *Lancet.* Feb 19 2011; 377(9766): 658-66. PMID 21315441
17. Abraham WT, Stevenson LW, Bourge RC, et al. Sustained efficacy of pulmonary artery pressure to guide adjustment of chronic heart failure therapy: complete follow-up results from the CHAMPION randomised trial. *Lancet.* Jan 30 2016; 387(10017): 453-61. PMID 26560249
18. Adamson PB, Abraham WT, Aaron M, et al. CHAMPION trial rationale and design: the long-term safety and clinical efficacy of a wireless pulmonary artery pressure monitoring system. *J Card Fail.* Jan 2011; 17(1): 3-10. PMID 21187258
19. CardioMEMS Champion™ HF Monitoring System: FDA Review of P100045/A004FDA Presentation - CardioMEMS: Oct. 9, 2013. 2013; <https://wayback.archive-it.org/7993/20170111163259/http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM370955.pdf>. Accessed May 5, 2024.
20. CardioMEMS Champion™ Heart Failure Monitoring System: Presentation - CardioMEMS: Oct. 9, 2013. 2013; <https://wayback.archive-it.org/7993/20170111163201/http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/MedicalDevices/MedicalDevicesAdvisoryCommittee/CirculatorySystemDevicesPanel/UCM370951.pdf>. Accessed May 4, 2024.
21. Shavelle DM, Desai AS, Abraham WT, et al. Lower Rates of Heart Failure and All-Cause Hospitalizations During Pulmonary Artery Pressure-Guided Therapy for Ambulatory Heart Failure: One-Year Outcomes From the CardioMEMS Post-Approval Study. *Circ Heart Fail.* Aug 2020; 13(8): e006863. PMID 32757642
22. DeFilippis EM, Henderson J, Axsom KM, et al. Remote Hemodynamic Monitoring Equally Reduces Heart Failure Hospitalizations in Women and Men in Clinical Practice: A Sex-Specific Analysis of the CardioMEMS Post-Approval Study. *Circ Heart Fail.* Jun 2021; 14(6): e007892. PMID 34129363
23. Lindenfeld J, Zile MR, Desai AS, et al. Haemodynamic-guided management of heart failure (GUIDE-HF): a randomised controlled trial. *Lancet.* Sep 11 2021; 398(10304): 991-1001. PMID 34461042
24. Zile MR, Desai AS, Costanzo MR, et al. The GUIDE-HF trial of pulmonary artery pressure monitoring in heart failure: impact of the COVID-19 pandemic. *Eur Heart J.* Jul 14 2022; 43(27): 2603-2618. PMID 35266003

25. Brughts JJ, Radhoe SP, Clephas PRD, et al. Remote haemodynamic monitoring of pulmonary artery pressures in patients with chronic heart failure (MONITOR-HF): a randomised clinical trial. *Lancet*. Jun 24 2023; 401(10394): 2113-2123. PMID 37220768
26. Kishino Y, Kuno T, Malik AH, et al. Effect of pulmonary artery pressure-guided therapy on heart failure readmission in a nationally representative cohort. *ESC Heart Fail*. Aug 2022; 9(4): 2511-2517. PMID 35560987
27. Cowie MR, Flett A, Cowburn P, et al. Real-world evidence in a national health service: results of the UK CardioMEMS HF System Post-Market Study. *ESC Heart Fail*. Feb 2022; 9(1): 48-56. PMID 34882989
28. Heywood JT, Zalawadiya S, Bourge RC, et al. Sustained Reduction in Pulmonary Artery Pressures and Hospitalizations During 2 Years of Ambulatory Monitoring. *J Card Fail*. Jan 2023; 29(1): 56-66. PMID 36332900
29. Angermann CE, Assmus B, Anker SD, et al. Pulmonary artery pressure-guided therapy in ambulatory patients with symptomatic heart failure: the CardioMEMS European Monitoring Study for Heart Failure (MEMS-HF). *Eur J Heart Fail*. Oct 2020; 22(10): 1891-1901. PMID 32592227
30. Abraham J, Bharmi R, Jonsson O, et al. Association of Ambulatory Hemodynamic Monitoring of Heart Failure With Clinical Outcomes in a Concurrent Matched Cohort Analysis. *JAMA Cardiol*. Jun 01 2019; 4(6): 556-563. PMID 31090869
31. Desai AS, Bhimaraj A, Bharmi R, et al. Ambulatory Hemodynamic Monitoring Reduces Heart Failure Hospitalizations in "Real-World" Clinical Practice. *J Am Coll Cardiol*. May 16 2017; 69(19): 2357-2365. PMID 28330751
32. Lin AL, Hu G, Dhruva SS, et al. Quantification of Device-Related Event Reports Associated With the CardioMEMS Heart Failure System. *Circ Cardiovasc Qual Outcomes*. Oct 2022; 15(10): e009116. PMID 36252112
33. Vaduganathan M, DeFilippis EM, Fonarow GC, et al. Postmarketing Adverse Events Related to the CardioMEMS HF System. *JAMA Cardiol*. Nov 01 2017; 2(11): 1277-1279. PMID 28975249
34. Krzesiński P, Jankowska EA, Siebert J, et al. Effects of an outpatient intervention comprising nurse-led non-invasive assessments, telemedicine support and remote cardiologists' decisions in patients with heart failure (AMULET study): a randomised controlled trial. *Eur J Heart Fail*. Mar 2022; 24(3): 565-577. PMID 34617373
35. Kamath SA, Drazner MH, Tasissa G, et al. Correlation of impedance cardiography with invasive hemodynamic measurements in patients with advanced heart failure: the BioImpedance CardioGraphy (BIG) substudy of the Evaluation Study of Congestive Heart Failure and Pulmonary Artery Catheterization Effectiveness (ESCAPE) Trial. *Am Heart J*. Aug 2009; 158(2): 217-23. PMID 19619697
36. Anand IS, Greenberg BH, Fogoros RN, et al. Design of the Multi-Sensor Monitoring in Congestive Heart Failure (MUSIC) study: prospective trial to assess the utility of continuous wireless physiologic monitoring in heart failure. *J Card Fail*. Jan 2011; 17(1): 11-6. PMID 21187259
37. Anand IS, Tang WH, Greenberg BH, et al. Design and performance of a multisensor heart failure monitoring algorithm: results from the multisensor monitoring in congestive heart failure (MUSIC) study. *J Card Fail*. Apr 2012; 18(4): 289-95. PMID 22464769
38. Packer M, Abraham WT, Mehra MR, et al. Utility of impedance cardiography for the identification of short-term risk of clinical decompensation in stable patients with chronic heart failure. *J Am Coll Cardiol*. Jun 06 2006; 47(11): 2245-52. PMID 16750691
39. Amir O, Ben-Gal T, Weinstein JM, et al. Evaluation of remote dielectric sensing (ReDS) technology-guided therapy for decreasing heart failure re-hospitalizations. *Int J Cardiol*. Aug 01 2017; 240: 279-284. PMID 28341372
40. Silber HA, Trost JC, Johnston PV, et al. Finger photoplethysmography during the Valsalva maneuver reflects left ventricular filling pressure. *Am J Physiol Heart Circ Physiol*. May 15 2012; 302(10): H2043-7. PMID 22389389
41. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Am Coll Cardiol*. Aug 08 2017; 70(6): 776-803. PMID 28461007
42. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint

Committee on Clinical Practice Guidelines. *Circulation*. May 03 2022; 145(18): e895-e1032. PMID 35363499

43. National Institute for Health and Care Excellence (NICE). Interventional procedures guidance: Percutaneous implantation of pulmonary artery pressure sensors for monitoring treatment of chronic heart failure [IPG711]. November 2021; <https://www.nice.org.uk/guidance/ipg711>. Accessed May 5, 2024.
44. Dickinson MG, Allen LA, Albert NA, et al. Remote Monitoring of Patients With Heart Failure: A White Paper From the Heart Failure Society of America Scientific Statements Committee. *J Card Fail*. Oct 2018; 24(10): 682-694. PMID 30308242
45. Centers for Medicare & Medicaid Services (CMS). National coverage decision for cardiac output monitoring by thoracic electrical bioimpedance (TEB) (20.16). 2006; <http://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=267&ncdver=3&NCAId=82>. Accessed May 5, 2024.