



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent licensee of the Blue Cross and Blue Shield Association

Medical Policy

Serum Biomarker Human Epididymis Protein 4

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 290

BCBSA Reference Number: 2.04.66 (For Plan internal use only)

NCD/LCD: N/A

Related Policies

Multimarker Serum Testing Related to Ovarian Cancer #[249](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Measurement of human epididymis protein 4 is [INVESTIGATIONAL](#) for all indications.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue SM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following CPT code is considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
86305	Human epididymis protein 4 (HE4)

Description

Ovarian Cancer

Ovarian cancer is the fifth most common cause of cancer mortality among U.S. women. According to Surveillance Epidemiology and End Results data, in 2023, an estimated 19,710 women will be diagnosed with ovarian cancer and 13,270 women will die of the disease.¹ The stage at diagnosis is an important predictor of survival; however, most women are not diagnosed until the disease has spread. For the period 2012 to 2018, 57% of women with ovarian cancer were diagnosed when the disease had distant metastases (stage IV), and this was associated with a 5-year survival rate of 31%. In contrast, 17% of women diagnosed with localized cancer (stage I) had a 5-year survival rate of 93%. Epithelial ovarian tumors account for 85% to 90% of ovarian cancers.²

Research from the Ovarian Cancer in Women of African Ancestry (OCWAA) consortium reports that Black women with ovarian cancer have worse survival than White women.³ Contributors to this disparity may include education level, nulliparity, smoking status, body mass index, diabetes, and postmenopausal hormone therapy duration.

Treatment

The standard treatment for epithelial ovarian cancer is surgical staging and primary cytoreductive surgery followed by chemotherapy in most cases. There is a lack of consensus about an optimal approach to the follow-up of patients with ovarian cancer after or during primary treatment. Patients undergo regular physical examinations and may have imaging studies. In addition, managing patients with serial measurements of the biomarker cancer antigen 125 (CA 125) to detect early recurrence of disease is common. A rising CA 125 level has been found to correlate with disease recurrence and has been found to detect recurrent ovarian cancer earlier than clinical detection. However, a survival advantage of initiating treatment based on early detection with CA 125 has not been demonstrated to date. For example, a 2010 randomized controlled trial in women with ovarian cancer that was in complete remission did not find a significant difference in overall survival when treatment for remission was initiated after CA 125 concentration exceeded twice the limit of normal compared to delaying treatment initiation until symptom onset.⁴

Human epididymis protein 4 (HE4) is a protein that circulates in the serum and has been found to be overexpressed in epithelial ovarian cancer, lung adenocarcinoma, breast cancer, pancreatic cancer, endometrial cancer, and bladder cancer. HE4 is made up of 2 whey acidic proteins with a 4 disulfide core domain and has been proposed as a biomarker for monitoring patients with epithelial ovarian cancer.

Evaluation of Adnexal Masses

This evidence review also addresses the use of the HE4 as a stand-alone test for evaluating women with ovarian masses who have not been diagnosed with ovarian cancer. Such patients undergo a diagnostic workup to determine whether the risk of malignancy is sufficiently high to warrant surgical removal. In patients for whom surgery is indicated, further evaluation may be warranted to determine if a surgical referral to a specialist with expertise in ovarian cancer is warranted. The Risk of Ovarian Malignancy Algorithm (ROMA) test combines HE4, CA 125, and menopausal status into a numeric score. The ROMA

test has been cleared by U.S. Food and Drug Administration (FDA) for predicting the risk that an adnexal mass is malignant; this test and other combination biomarker tests, are considered separately in policy #249 (multimarker serum testing related to ovarian cancer).

Summary

Human epididymis protein 4 (HE4) is a novel biomarker that has been cleared by the U.S. Food and Drug Administration for monitoring patients with epithelial ovarian cancer. HE4 is proposed as a replacement for or a complement to cancer antigen 125 (CA 125) for monitoring disease progression and recurrence. HE4 has also been proposed as a test to evaluate women with ovarian masses and to screen for ovarian cancer in asymptomatic women.

Summary of Evidence

For individuals who have ovarian cancer who receive a measurement of serum biomarker HE4, the evidence includes 7 nonrandomized prospective and retrospective studies comparing the diagnostic accuracy of HE4 with CA 125 for predicting disease progression and/or recurrence. Relevant outcomes are overall survival (OS), disease-specific survival, test validity, other test performance measures, and change in disease status. Data submitted to the U.S. Food and Drug Administration for approval of commercial HE4 tests found that HE4 was not inferior to CA 125 for detecting ovarian cancer recurrence. Although a single prospective observational study found elevated levels of HE4, but not CA 125, at the time of cancer progression to be significantly associated with reduced OS, a direct comparison between biomarkers was not provided. Overall, the superiority of HE4 to CA 125 (alone or in combination), the key question in the evidence review, was not demonstrated in the available literature. In addition, there is no established cutoff in HE4 levels for monitoring disease progression, and cutoffs in studies varied. There is no direct evidence from prospective controlled studies on the impact of HE4 testing on health outcomes, and no clear chain of evidence that changes in management based on HE4 would lead to an improved health outcome. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have adnexal masses who receive a measurement of serum biomarker HE4, the evidence includes diagnostic accuracy studies and meta-analyses. Relevant outcomes are OS, disease-specific survival, test validity, and other test performance measures. Meta-analyses have generally found that HE4 and CA 125 have a similar overall diagnostic accuracy (ie, sensitivity, specificity) and several found that HE4 has significantly higher specificity than CA 125, but not sensitivity. Two meta-analyses had mixed findings on whether the combination of HE4 and CA 125 is superior to CA 125 alone for the initial diagnosis of ovarian cancer. The number of studies evaluating the combined test is relatively low, and publication bias in studies of HE4 has been identified. In addition, studies have not found that HE4 improves diagnostic accuracy beyond that of subjective assessment of transvaginal ultrasound. There is no direct evidence from prospective controlled studies on the impact of HE4 testing on health outcomes, and no clear chain of evidence that changes in management based on HE4 would lead to an improved health outcome. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic and not at high risk of ovarian cancer who receive screening with a serum biomarker HE4 test, the evidence includes several retrospective comparative studies and no prospective studies comparing health outcomes in asymptomatic women managed with and without HE4 screening. Relevant outcomes are OS, disease-specific survival, test validity, and other test performance measures. The retrospective studies found that HE4 levels increased over time in women ultimately diagnosed with ovarian cancer. Prospective comparative studies are needed to determine definitively whether HE4 testing is a useful screening tool. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
1/2024	Annual policy review. Description, summary, and references updated. Policy statements unchanged.

2/2023	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
3/2018	Annual policy review. New references added.
1/2017	Annual policy review. New references added.
1/2016	Annual policy review. New references added.
12/2015	Added coding language.
5/2015	Annual policy review. New references added.
5/2014	Annual policy review. New references added.
12/2013	Annual policy review. New references added.
3/2013	Annual policy review. No changes to policy statement.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
9/2011	Reviewed - Medical Policy Group – Urology, Obstetrics and Gynecology. No changes to policy statements.
7/2011	Reviewed - Medical Policy Group – Hematology and Oncology. No changes to policy statements.
12/3/2010	New policy describing ongoing non-coverage of serum biomarker human epididymis protein 4 (HE4), effective 12/3/2010.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Surveillance Epidemiology and End Results Program (SEER). SEER Stat Fact: Ovarian Cancer. n.d.; <http://seer.cancer.gov/statfacts/html/ovary.html>. Accessed October 31, 2023.
2. Ledermann JA, Raja FA, Fotopoulou C, et al. Newly diagnosed and relapsed epithelial ovarian carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* Oct 2013; 24 Suppl 6: vi24-32. PMID 24078660
3. Harris HR, Guertin KA, Camacho TF, et al. Racial disparities in epithelial ovarian cancer survival: An examination of contributing factors in the Ovarian Cancer in Women of African Ancestry consortium. *Int J Cancer.* Oct 15 2022; 151(8): 1228-1239. PMID 35633315
4. Rustin GJ, van der Burg ME, Griffin CL, et al. Early versus delayed treatment of relapsed ovarian cancer (MRC OV05/EORTC 55955): a randomised trial. *Lancet.* Oct 02 2010; 376(9747): 1155-63. PMID 20888993
5. Han Y, Jiang L, Liu K, et al. Predictive Value of HE4 in Platinum-Based Chemotherapy for Ovarian Cancer : A Systematic Review. *Front Oncol.* 2021; 11: 703949. PMID 34307173
6. Food and Drug Administration, 510(k) substantial equivalence determination decision summary: assay only (K072939). n.d.; http://www.accessdata.fda.gov/cdrh_docs/reviews/K072939.pdf. Accessed October 31, 2023.
7. Food and Drug Administration. 510(k) substantial equivalence determination decision summary: assay only (K093957). n.d.; http://www.accessdata.fda.gov/cdrh_docs/reviews/K093957.pdf. Accessed October 30, 2023.

8. Nassir M, Guan J, Luketina H, et al. The role of HE4 for prediction of recurrence in epithelial ovarian cancer patients-results from the OVCAD study. *Tumour Biol.* Mar 2016; 37(3): 3009-16. PMID 26419591
9. Vallius T, Hynninen J, Auranen A, et al. Postoperative human epididymis protein 4 predicts primary therapy outcome in advanced epithelial ovarian cancer. *Tumour Biol.* Feb 2017; 39(2): 1010428317691189. PMID 28218038
10. Potenza E, Parpinel G, Laudani ME, et al. Prognostic and predictive value of combined HE-4 and CA-125 biomarkers during chemotherapy in patients with epithelial ovarian cancer. *Int J Biol Markers.* Dec 2020; 35(4): 20-27. PMID 33126819
11. Salminen L, Gidwani K, Grønman S, et al. HE4 in the evaluation of tumor load and prognostic stratification of high grade serous ovarian carcinoma. *Acta Oncol.* Dec 2020; 59(12): 1461-1468. PMID 33030975
12. Rong Y, Li L. Early clearance of serum HE4 and CA125 in predicting platinum sensitivity and prognosis in epithelial ovarian cancer. *J Ovarian Res.* Jan 04 2021; 14(1): 2. PMID 33397458
13. Samborski A, Miller MC, Blackman A, et al. HE4 and CA125 serum biomarker monitoring in women with epithelial ovarian cancer. *Tumour Biol.* 2022; 44(1): 205-213. PMID 36189508
14. Olsen M, Lof P, Stiekema A, et al. The diagnostic accuracy of human epididymis protein 4 (HE4) for discriminating between benign and malignant pelvic masses: a systematic review and meta-analysis. *Acta Obstet Gynecol Scand.* Oct 2021; 100(10): 1788-1799. PMID 34212386
15. Suri A, Perumal V, Ammalli P, et al. Diagnostic measures comparison for ovarian malignancy risk in Epithelial ovarian cancer patients: a meta-analysis. *Sci Rep.* Aug 27 2021; 11(1): 17308. PMID 34453074
16. Huang J, Chen J, Huang Q. Diagnostic value of HE4 in ovarian cancer: A meta-analysis. *Eur J Obstet Gynecol Reprod Biol.* Dec 2018; 231: 35-42. PMID 30317143
17. Dayyani F, Uhlig S, Colson B, et al. Diagnostic Performance of Risk of Ovarian Malignancy Algorithm Against CA125 and HE4 in Connection With Ovarian Cancer: A Meta-analysis. *Int J Gynecol Cancer.* Nov 2016; 26(9): 1586-1593. PMID 27540691
18. Macedo AC, da Rosa MI, Lumertz S, et al. Accuracy of serum human epididymis protein 4 in ovarian cancer diagnosis: a systematic review and meta-analysis. *Int J Gynecol Cancer.* Sep 2014; 24(7): 1222-31. PMID 25078339
19. Wang J, Gao J, Yao H, et al. Diagnostic accuracy of serum HE4, CA125 and ROMA in patients with ovarian cancer: a meta-analysis. *Tumour Biol.* Jun 2014; 35(6): 6127-38. PMID 24627132
20. Zhen S, Bian LH, Chang LL, et al. Comparison of serum human epididymis protein 4 and carbohydrate antigen 125 as markers in ovarian cancer: A meta-analysis. *Mol Clin Oncol.* Jul 2014; 2(4): 559-566. PMID 24940495
21. Yang Z, Wei C, Luo Z, et al. Clinical value of serum human epididymis protein 4 assay in the diagnosis of ovarian cancer: a meta-analysis. *Onco Targets Ther.* 2013; 6: 957-66. PMID 23901285
22. Ferraro S, Braga F, Lanzoni M, et al. Serum human epididymis protein 4 vs carbohydrate antigen 125 for ovarian cancer diagnosis: a systematic review. *J Clin Pathol.* Apr 2013; 66(4): 273-81. PMID 23426716
23. Yu S, Yang HJ, Xie SQ, et al. Diagnostic value of HE4 for ovarian cancer: a meta-analysis. *Clin Chem Lab Med.* Feb 03 2012; 50(8): 1439-46. PMID 22868811
24. Kaijser J, Van Gorp T, Smet ME, et al. Are serum HE4 or ROMA scores useful to experienced examiners for improving characterization of adnexal masses after transvaginal ultrasonography?. *Ultrasound Obstet Gynecol.* Jan 2014; 43(1): 89-97. PMID 23828371
25. Moszynski R, Szubert S, Szpurek D, et al. Usefulness of the HE4 biomarker as a second-line test in the assessment of suspicious ovarian tumors. *Arch Gynecol Obstet.* Dec 2013; 288(6): 1377-83. PMID 23722285
26. Nikolova T, Zivadinovic R, Evtimovska N, et al. Diagnostic performance of human epididymis protein 4 compared to a combination of biophysical and biochemical markers to differentiate ovarian endometriosis from epithelial ovarian cancer in premenopausal women. *J Obstet Gynaecol Res.* Dec 2017; 43(12): 1870-1879. PMID 29027715
27. Gentry-Maharaj A, Burnell M, Dilley J, et al. Serum HE4 and diagnosis of ovarian cancer in postmenopausal women with adnexal masses. *Am J Obstet Gynecol.* Jan 2020; 222(1): 56.e1-56.e17. PMID 31351062

28. Carreras-Diequez N, Glickman A, Munmany M, et al. Comparison of HE4, CA125, ROMA and CPH-I for Preoperative Assessment of Adnexal Tumors. *Diagnostics (Basel)*. Jan 17 2022; 12(1). PMID 35054393
29. Lof P, van de Vrie R, Korse CM, et al. Can serum human epididymis protein 4 (HE4) support the decision to refer a patient with an ovarian mass to an oncology hospital?. *Gynecol Oncol*. Aug 2022; 166(2): 284-291. PMID 35688656
30. Anderson GL, McIntosh M, Wu L, et al. Assessing lead time of selected ovarian cancer biomarkers: a nested case-control study. *J Natl Cancer Inst*. Jan 06 2010; 102(1): 26-38. PMID 20042715
31. Urban N, Thorpe JD, Bergan LA, et al. Potential role of HE4 in multimodal screening for epithelial ovarian cancer. *J Natl Cancer Inst*. Nov 02 2011; 103(21): 1630-4. PMID 21917606
32. Terry KL, Schock H, Fortner RT, et al. A Prospective Evaluation of Early Detection Biomarkers for Ovarian Cancer in the European EPIC Cohort. *Clin Cancer Res*. Sep 15 2016; 22(18): 4664-75. PMID 27060155
33. Eskander R, Berman M, Keder L. Practice Bulletin No. 174: Evaluation and Management of Adnexal Masses. *Obstet Gynecol*. Nov 2016; 128(5): e210-e226. PMID 27776072
34. Committee Opinion No. 716: The Role of the Obstetrician-Gynecologist in the Early Detection of Epithelial Ovarian Cancer in Women at Average Risk. *Obstet Gynecol*. Sep 2017; 130(3): e146-e149. PMID 28832487
35. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer. Version 2.2023. https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf. Accessed October 31, 2023.
36. National Institute for Health and Care Excellence (NICE). Ovarian cancer: recognition and initial management [CG122]. 2011; <https://www.nice.org.uk/guidance/cg122>. Accessed October 31, 2023.
37. U.S. Preventive Services Task Force. Recommendation Statement: Screening for Ovarian Cancer. 2018; file:///C:/Users/alt/Downloads/ovarian-cancer-final-rec-statement.pdf. Accessed October 31, 2023.