



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Diagnosis and Treatment of Sacroiliac Joint Pain

Table of Contents

- [Policy: Commercial](#)
- [Description](#)
- [Information Pertaining to All Policies](#)
- [Authorization Information](#)
- [Policy History](#)
- [Coding Information](#)
- [References](#)

Policy Number: 320

BCBSA Reference Number: 6.01.23 (For Plan internal use only)

Related Policies

- Epidural Steroid Injections for Neck and Back Pain, [#690](#)
- Facet Joint Denervation, [#140](#)
- Percutaneous Vertebroplasty and Sacroplasty, [#484](#)
- Prolotherapy, Joint Sclerotherapy and Ligamentous Injections with Sclerosing Agents, [#183](#)
- Prior Authorization Request Form for Diagnosis and Treatment of Sacroiliac Joint Pain, [#927](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Arthrography of the sacroiliac joint is [INVESTIGATIONAL](#).

Minimally invasive fixation/fusion of the sacroiliac joint using transiliac placement of a titanium triangular implant (eg, iFuse) may be considered [MEDICALLY NECESSARY](#) when **ALL** of the following criteria have been met:

- Pain is at least 5 on a 0 to 10 rating scale that impacts quality of life or limits activities of daily living; **AND**
- There is an absence of generalized pain behavior (eg, somatoform disorder) or generalized pain disorders (eg, fibromyalgia); **AND**
- Individuals have undergone and failed a minimum 6 months of intensive nonoperative treatment that must include medication optimization, activity modification, bracing, and active therapeutic exercise targeted at the lumbar spine, pelvis, sacroiliac joint, and hip, including a home exercise program; **AND**
- Pain is caudal to the lumbar spine (L5 vertebra), localized over the posterior sacroiliac joint, and consistent with sacroiliac joint pain; **AND**
- A thorough physical examination demonstrates localized tenderness with palpation over the sacral sulcus (Fortin's point) in the absence of tenderness of similar severity elsewhere; **AND**
- There is a positive response to a cluster of 3 provocative tests (eg, thigh thrust test, compression test, Gaenslen sign, distraction test, Patrick test, posterior provocation test); **AND**
- Diagnostic imaging studies include **ALL** of the following:

- Imaging (plain radiographs and computed tomography or magnetic resonance imaging) of the sacroiliac joint excludes the presence of destructive lesions (eg, tumor, infection) or inflammatory arthropathy of the sacroiliac joint; **AND**
- Imaging of the pelvis (anteroposterior plain radiograph) rules out concomitant hip pathology; **AND**
- Imaging of the lumbar spine (computed tomography or magnetic resonance imaging) is performed to rule out neural compression or other degenerative conditions that can be causing low back or buttock pain; **AND**
- Imaging of the sacroiliac joint indicates evidence of injury and/or degeneration; **AND**
- There is at least a 75% reduction in pain for the expected duration of the anesthetic used following an image-guided, contrast-enhanced intra-articular sacroiliac joint injection on 2 separate occasions; **AND**
- A trial of a therapeutic sacroiliac joint injection (ie, corticosteroid injection) has been performed at least once.

Note: This technically demanding procedure **should only be done by surgeons** who have specific training and expertise in minimally invasive sacroiliac joint fusion surgery for chronic sacroiliac joint pain and who regularly use image-guidance for implant placement.

Fusion/stabilization of the sacroiliac joint for the treatment of back pain presumed to originate from the sacroiliac joint is considered **INVESTIGATIONAL** under all other conditions and with any other devices not listed above.

Radiofrequency denervation of the sacroiliac joint is considered **INVESTIGATIONAL**.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is required for minimally invasive fixation/fusion of the sacroiliac joint.
Commercial PPO	Prior authorization is required for minimally invasive fixation/fusion of the sacroiliac joint.

Requesting Prior Authorization Using Authorization Manager

Providers will need to use [Authorization Manager](#) to submit initial authorization requests for services. Authorization Manager, available 24/7, is the quickest way to review authorization requirements, request authorizations, submit clinical documentation, check existing case status, and view/print the decision letter. For commercial members, the requests must meet medical policy guidelines.

To ensure the request is processed accurately and quickly:

- Enter the facility's NPI or provider ID for where services are being performed.
- Enter the appropriate surgeon's NPI or provider ID as the servicing provider, *not* the billing group.

Authorization Manager Resources

- Refer to our [Authorization Manager](#) page for tips, guides, and video demonstrations.

Complete Prior Authorization Request Form for Diagnosis and Treatment of Sacroiliac Joint Pain [\(927\)](#) using [Authorization Manager](#).

For out of network providers: Requests should still be faxed to 888-282-0780.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO and Indemnity:

CPT Codes

CPT codes:	Code Description
27279	Arthrodesis, sacroiliac joint, percutaneous or minimally invasive (indirect visualization), with image guidance, includes obtaining bone graft when performed, and placement of transfixing device

The following HCPCS and CPT code is considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:

HCPCS Codes

HCPCS codes:	Code Description
G0259	Injection procedure for sacroiliac joint; arthrography

CPT Codes

CPT codes:	Code Description
27278	Arthrodesis, sacroiliac joint, percutaneous, with image guidance, including placement of intra-articular implant(s) (eg, bone allograft[s], synthetic device[s]), without placement of transfixation device
64625	Radiofrequency ablation, nerves innervating the sacroiliac joint, with image guidance (ie, fluoroscopy or computed tomography)

DESCRIPTION

Sacroiliac Joint Pain

Similar to other structures in the spine, it is assumed the sacroiliac joint (SIJ) may be a source of low back pain. In fact, before 1928, the SIJ was thought to be the most common cause of sciatica. In 1928, the role of the intervertebral disc was elucidated, and from that point forward, the SIJ received less research attention.

Diagnosis

Research into SIJ pain has been plagued by a lack of a criterion standard to measure its prevalence and against which various clinical examinations can be validated. For example, SIJ pain typically presents without any consistent, demonstrable radiographic or laboratory features and most commonly exists in the setting of morphologically normal joints. Clinical tests for SIJ pain may include various movement tests, palpation to detect tenderness, and pain descriptions by the patient. Further confounding the study of the SIJ is that multiple structures, (eg, posterior facet joints, lumbar discs) may refer pain to the area surrounding the SIJ.

Because of inconsistent information obtained from history and physical examination, some have proposed the use of image-guided anesthetic injection into the SIJ for the diagnosis of SIJ pain. Treatments being investigated for SIJ pain include prolotherapy (see evidence review 2.01.26), corticosteroid injection, radiofrequency ablation, stabilization, and arthrodesis. Some procedures have been referred to as SIJ fusion but may be more appropriately called fixation due to little to no bridging bone on radiographs. Devices for SIJ fixation/fusion that promote bone ingrowth to fixate the implants include a triangular implant (iFuse Implant System) and cylindrical threaded devices (eg, Rialto, SImmetry, Silex, SambaScrew, SI-LOK). Some devices also have a slot in the middle where autologous or allogeneic bone can be inserted. This added bone is intended to promote the fusion of the SIJ.

A 2021 review identified 33 different devices that could be implanted using either a lateral transiliac approach (n=21), posterior allograft approach (n=6), posterolateral approach (n=3), or a combination of the approaches (n=3).¹ The iliosacral and posterolateral approaches use up to 3 implants that pass through the ilium, while the posterior approach involves inserting implants directly into the SIJ. Many of the devices are intended to be used with allograft bone. Implants composed entirely of allograft bone are typically inserted through a posterior approach. The authors found no published evidence for 23 of the 33 devices identified.

Summary

Description

Sacroiliac joint (SIJ) arthrography using fluoroscopic guidance with an injection of an anesthetic has been explored as a diagnostic test for SIJ pain. Duplication of the patient's pain pattern with the injection of contrast medium suggests a sacroiliac etiology, as does relief of chronic back pain with an injection of local anesthetic. Treatment of SIJ pain with corticosteroids, radiofrequency ablation (RFA), stabilization, or minimally invasive SIJ fusion has also been explored.

Summary of Evidence

Diagnostic

For individuals who have suspected SIJ pain who receive a diagnostic sacroiliac block, the evidence includes systematic reviews. Relevant outcomes are test validity, symptoms, functional outcomes, quality of life (QOL), medication use, and treatment-related morbidity. Current evidence is conflicting on the diagnostic utility of SIJ blocks. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have SIJ pain who receive RFA, the evidence includes 5 RCTs using different radiofrequency applications and case series. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. Meta-analysis of available sham-controlled RCTs suggests that there may be a small effect of RFA on SIJ pain at short-term (1 to 3 months) follow-up. However, the RCTs of RFA have methodologic limitations, and there is limited data on the duration of the treatment effect. The single RCT with 6 and 12-month follow-up showed no significant benefit of RFA compared to an exercise control group at these time points. In addition, heterogeneity of RFA treatment techniques precludes generalizing results across different studies. For RFA with a cooled probe, 2 small RCTs reported short-term benefits, but these are insufficient to determine the overall effect on health outcomes. An RCT on palisade RFA of the SIJ did not include a sham control. Another sham-controlled RCT showed no benefit from RFA. Further high-quality controlled trials are needed to compare this procedure in defined populations with sham control and alternative treatments. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have SIJ pain who receive SIJ fixation/fusion with a transiliac triangular implant, the evidence includes 2 nonblinded RCTs of minimally invasive fusion, prospective cohorts with more than 85% follow-up, and case series. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. Both RCTs have reported outcomes past 6 months, after which crossover was allowed. Both studies reported significantly greater reductions in visual analog scale (VAS) pain scores and Oswestry Disability Index (ODI) scores in SIJ fusion patients than in control groups. The reductions in pain and disability observed in the SIJ fusion group at 6 months were

maintained out to 1 year compared with controls who had not crossed over. The RCTs were nonblinded without a placebo or an active control group. Prospective cohorts and case series with sample sizes ranging from 45 to 149 patients and low dropout rates (<15%) also showed reductions in pain and disability out to 5 years. The cohort studies and case series are consistent with the durability of treatment benefit. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have SIJ pain who receive SIJ fusion/fixation with an implant other than a transiliac triangular implant, the evidence includes 3 prospective cohort studies and retrospective case series. Relevant outcomes are symptoms, functional outcomes, QOL, medication use, and treatment-related morbidity. Two prospective cohort studies were conducted with transiliac screws and the third with a device inserted through a posterior approach. No controlled studies were identified. Meta-analyses of the available prospective and retrospective studies indicate improvement in subjective outcomes from before surgery to follow-up, but with a possible difference in outcomes between the more well studied triangular transiliac implant and other implant designs and approaches. There is uncertainty in the health benefit of SIJ fusion/fixation with these implant designs. Therefore, controlled studies with a larger number of patients and longer follow-up are needed to evaluate these devices. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
1/2024	Annual policy review. Policy updated with literature review through September 13, 2023; references added. Policy statements unchanged. Clarified coding information.
9/2023	Policy clarified to include prior authorization requests using Authorization Manager.
7/2023	Clarified coding information.
5/2023	Policy clarified. Policy reactivated in May 2023 to reinstate policy statements on minimally invasive fixation/fusion of the sacroiliac joint. Effective 5/1/2023. Policy statements on anesthetic injection for diagnosing SIJ pain and corticosteroid injection for treatment of SIJ pain remain retired.
1/2023	Annual policy review. Minor editorial refinements to policy statements; intent unchanged. Clarified coding information.
6/2022	Prior authorization information clarified for PPO plans. Effective 6/1/2022.
1/2022	Annual policy review. Policy clarified. "Transiliac placement" and "eg, iFuse" added to the medically necessary statement on sacroiliac joint fusion.
1/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
1/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2020	Clarified coding information.
1/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2018	Annual policy review. New medically necessary indications described for SIJ fusion/stabilization with a titanium triangular implant under the specific conditions. Clarified coding information. Effective 6/1/2018.
12/2016	Annual policy review. New references added.
11/2016	Annual policy review. New references added.
10/2016	Annual policy review. First medically necessary policy statement clarified.
1/2016	Annual policy review. New references added.
6/2015	Annual policy review. New references added.
1/2015	Clarified coding information.
10/2014	Annual policy review. New medically necessary and investigational indications described. Coding information clarified. Effective 10/1/2014.

6/2014	Annual policy review. New medical policy describing investigational indications. Effective 6/1/2014.
--------	------------------------------------------------------------------------------------------------------

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Himstead AS, Brown NJ, Shahrestani S, et al. Trends in Diagnosis and Treatment of Sacroiliac Joint Pathology Over the Past 10 Years: Review of Scientific Evidence for New Devices for Sacroiliac Joint Fusion. *Cureus*. Jun 2021; 13(6): e15415. PMID 34249562
2. Dreyfuss P, Michaelsen M, Pauza K, et al. The value of medical history and physical examination in diagnosing sacroiliac joint pain. *Spine (Phila Pa 1976)*. Nov 15 1996; 21(22): 2594-602. PMID 8961447
3. Simopoulos TT, Manchikanti L, Gupta S, et al. Systematic Review of the Diagnostic Accuracy and Therapeutic Effectiveness of Sacroiliac Joint Interventions. *Pain Physician*. Sep-Oct 2015; 18(5): E713-56. PMID 26431129
4. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. Apr 2013; 16(2 Suppl): S49-283. PMID 23615883
5. Manchikanti L, Datta S, Derby R, et al. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: part 1. Diagnostic interventions. *Pain Physician*. May-Jun 2010; 13(3): E141-74. PMID 20495596
6. Manchikanti L, Datta S, Gupta S, et al. A critical review of the American Pain Society clinical practice guidelines for interventional techniques: part 2. Therapeutic interventions. *Pain Physician*. Jul-Aug 2010; 13(4): E215-64. PMID 20648212
7. Rupert MP, Lee M, Manchikanti L, et al. Evaluation of sacroiliac joint interventions: a systematic appraisal of the literature. *Pain Physician*. Mar-Apr 2009; 12(2): 399-418. PMID 19305487
8. Chou R, Atlas SJ, Stanos SP, et al. Nonsurgical interventional therapies for low back pain: a review of the evidence for an American Pain Society clinical practice guideline. *Spine (Phila Pa 1976)*. May 01 2009; 34(10): 1078-93. PMID 19363456
9. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976)*. May 01 2009; 34(10): 1066-77. PMID 19363457
10. Hansen H, Manchikanti L, Simopoulos TT, et al. A systematic evaluation of the therapeutic effectiveness of sacroiliac joint interventions. *Pain Physician*. May-Jun 2012; 15(3): E247-78. PMID 22622913
11. Patel A, Kumar D, Singh S, et al. Effect of Fluoroscopic-Guided Corticosteroid Injection in Patients With Sacroiliac Joint Dysfunction. *Cureus*. Mar 2023; 15(3): e36406. PMID 37090293
12. Visser LH, Woudenberg NP, de Bont J, et al. Treatment of the sacroiliac joint in patients with leg pain: a randomized-controlled trial. *Eur Spine J*. Oct 2013; 22(10): 2310-7. PMID 23720124
13. Kim WM, Lee HG, Jeong CW, et al. A randomized controlled trial of intra-articular prolotherapy versus steroid injection for sacroiliac joint pain. *J Altern Complement Med*. Dec 2010; 16(12): 1285-90. PMID 21138388
14. Kennedy DJ, Engel A, Kreiner DS, et al. Fluoroscopically Guided Diagnostic and Therapeutic Intra-Articular Sacroiliac Joint Injections: A Systematic Review. *Pain Med*. Aug 2015; 16(8): 1500-18. PMID 26178855
15. Ab Aziz SNF, Zakaria Mohamad Z, Karupiah RK, et al. Efficacy of Sacroiliac Joint Injection With Anesthetic and Corticosteroid: A Prospective Observational Study. *Cureus*. Apr 2022; 14(4): e24039. PMID 35547453

16. Al Khayyat SG, Fogliame G, Barbagli S, et al. Ultrasound guided corticosteroids sacroiliac joint injections (SIJIs) in the management of active sacroiliitis: a real-life prospective experience. *J Ultrasound*. Jun 2023; 26(2): 479-486. PMID 36229757
17. Chandrupatla RS, Shahidi B, Bruno K, et al. A Retrospective Study on Patient-Specific Predictors for Non-Response to Sacroiliac Joint Injections. *Int J Environ Res Public Health*. Nov 23 2022; 19(23). PMID 36497595
18. Chou R, Fu R, Dana T, Pappas M, Hart E, Mauer KM. Interventional Treatments for Acute and Chronic Pain: Systematic Review. Comparative Effectiveness Review No. 247. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 75Q80120D00006.) AHRQ Publication No. 21-EHC030. Rockville, MD: Agency for Healthcare Research and Quality; September 2021. PMID: 34524764
19. Chappell ME, Lakshman R, Trotter P, et al. Radiofrequency denervation for chronic back pain: a systematic review and meta-analysis. *BMJ Open*. Jul 21 2020; 10(7): e035540. PMID 32699129
20. Juch JNS, Maas ET, Ostelo RWJG, et al. Effect of Radiofrequency Denervation on Pain Intensity Among Patients With Chronic Low Back Pain: The Mint Randomized Clinical Trials. *JAMA*. Jul 04 2017; 318(1): 68-81. PMID 28672319
21. Chen CH, Weng PW, Wu LC, et al. Radiofrequency neurotomy in chronic lumbar and sacroiliac joint pain: A meta-analysis. *Medicine (Baltimore)*. Jun 2019; 98(26): e16230. PMID 31261580
22. Cohen SP, Kapural L, Kohan L, et al. Cooled radiofrequency ablation versus standard medical management for chronic sacroiliac joint pain: a multicenter, randomized comparative effectiveness study. *Reg Anesth Pain Med*. Jul 05 2023. PMID 37407279
23. Mehta V, Poply K, Husband M, et al. The Effects of Radiofrequency Neurotomy Using a Strip-Lesioning Device on Patients with Sacroiliac Joint Pain: Results from a Single-Center, Randomized, Sham-Controlled Trial. *Pain Physician*. Nov 2018; 21(6): 607-618. PMID 30508988
24. van Tilburg CW, Schuurmans FA, Stronks DL, et al. Randomized Sham-controlled Double-Blind Multicenter Clinical Trial to Ascertain the Effect of Percutaneous Radiofrequency Treatment for Sacroiliac Joint Pain: Three-month Results. *Clin J Pain*. Nov 2016; 32(11): 921-926. PMID 26889616
25. Zheng Y, Gu M, Shi D, et al. Tomography-guided palisade sacroiliac joint radiofrequency neurotomy versus celecoxib for ankylosing spondylitis: a open-label, randomized, and controlled trial. *Rheumatol Int*. Sep 2014; 34(9): 1195-202. PMID 24518967
26. Patel N, Gross A, Brown L, et al. A randomized, placebo-controlled study to assess the efficacy of lateral branch neurotomy for chronic sacroiliac joint pain. *Pain Med*. Mar 2012; 13(3): 383-98. PMID 22299761
27. Patel N. Twelve-Month Follow-Up of a Randomized Trial Assessing Cooled Radiofrequency Denervation as a Treatment for Sacroiliac Region Pain. *Pain Pract*. Feb 2016; 16(2): 154-67. PMID 25565322
28. Whang P, Cher D, Polly D, et al. Sacroiliac Joint Fusion Using Triangular Titanium Implants vs. Non-Surgical Management: Six-Month Outcomes from a Prospective Randomized Controlled Trial. *Int J Spine Surg*. 2015; 9: 6. PMID 25785242
29. Polly DW, Cher DJ, Wine KD, et al. Randomized Controlled Trial of Minimally Invasive Sacroiliac Joint Fusion Using Triangular Titanium Implants vs Nonsurgical Management for Sacroiliac Joint Dysfunction: 12-Month Outcomes. *Neurosurgery*. Nov 2015; 77(5): 674-90; discussion 690-1. PMID 26291338
30. Polly DW, Swofford J, Whang PG, et al. Two-Year Outcomes from a Randomized Controlled Trial of Minimally Invasive Sacroiliac Joint Fusion vs. Non-Surgical Management for Sacroiliac Joint Dysfunction. *Int J Spine Surg*. 2016; 10: 28. PMID 27652199
31. Darr E, Meyer SC, Whang PG, et al. Long-term prospective outcomes after minimally invasive trans-iliac sacroiliac joint fusion using triangular titanium implants. *Med Devices (Auckl)*. 2018; 11: 113-121. PMID 29674852
32. Stuesson B, Kools D, Pflugmacher R, et al. Six-month outcomes from a randomized controlled trial of minimally invasive SI joint fusion with triangular titanium implants vs conservative management. *Eur Spine J*. Mar 2017; 26(3): 708-719. PMID 27179664
33. Dengler J, Stuesson B, Kools D, et al. Referred leg pain originating from the sacroiliac joint: 6-month outcomes from the prospective randomized controlled iMIA trial. *Acta Neurochir (Wien)*. Nov 2016; 158(11): 2219-2224. PMID 27629371

34. Dengler JD, Kools D, Pflugmacher R, et al. 1-Year Results of a Randomized Controlled Trial of Conservative Management vs. Minimally Invasive Surgical Treatment for Sacroiliac Joint Pain. *Pain Physician*. Sep 2017; 20(6): 537-550. PMID 28934785
35. Dengler J, Kools D, Pflugmacher R, et al. Randomized Trial of Sacroiliac Joint Arthrodesis Compared with Conservative Management for Chronic Low Back Pain Attributed to the Sacroiliac Joint. *J Bone Joint Surg Am*. Mar 06 2019; 101(5): 400-411. PMID 30845034
36. Duhon BS, Cher DJ, Wine KD, et al. Triangular Titanium Implants for Minimally Invasive Sacroiliac Joint Fusion: A Prospective Study. *Global Spine J*. May 2016; 6(3): 257-69. PMID 27099817
37. Duhon BS, Bitan F, Lockstadt H, et al. Triangular Titanium Implants for Minimally Invasive Sacroiliac Joint Fusion: 2-Year Follow-Up from a Prospective Multicenter Trial. *Int J Spine Surg*. 2016; 10: 13. PMID 27162715
38. Whang PG, Darr E, Meyer SC, et al. Long-Term Prospective Clinical And Radiographic Outcomes After Minimally Invasive Lateral Transiliac Sacroiliac Joint Fusion Using Triangular Titanium Implants. *Med Devices (Auckl)*. 2019; 12: 411-422. PMID 31576181
39. Patel V, Kovalsky D, Meyer SC, et al. Prospective Trial of Sacroiliac Joint Fusion Using 3D-Printed Triangular Titanium Implants. *Med Devices (Auckl)*. 2020; 13: 173-182. PMID 32607011
40. Vanaclocha V, Herrera JM, Saiz-Sapena N, et al. Minimally Invasive Sacroiliac Joint Fusion, Radiofrequency Denervation, and Conservative Management for Sacroiliac Joint Pain: 6-Year Comparative Case Series. *Neurosurgery*. Jan 01 2018; 82(1): 48-55. PMID 28431026
41. Spain K, Holt T. Surgical Revision after Sacroiliac Joint Fixation or Fusion. *Int J Spine Surg*. 2017; 11: 5. PMID 28377863
42. Schoell K, Buser Z, Jakoi A, et al. Postoperative complications in patients undergoing minimally invasive sacroiliac fusion. *Spine J*. Nov 2016; 16(11): 1324-1332. PMID 27349627
43. Tran ZV, Ivashchenko A, Brooks L. Sacroiliac Joint Fusion Methodology - Minimally Invasive Compared to Screw-Type Surgeries: A Systematic Review and Meta-Analysis. *Pain Physician*. Jan 2019; 22(1): 29-40. PMID 30700066
44. Lorio M, Kube R, Araghi A. International Society for the Advancement of Spine Surgery Policy 2020 Update-Minimally Invasive Surgical Sacroiliac Joint Fusion (for Chronic Sacroiliac Joint Pain): Coverage Indications, Limitations, and Medical Necessity. *Int J Spine Surg*. Dec 2020; 14(6): 860-895. PMID 33560247
45. Rappoport LH, Luna IY, Joshua G. Minimally Invasive Sacroiliac Joint Fusion Using a Novel Hydroxyapatite-Coated Screw: Preliminary 1-Year Clinical and Radiographic Results of a 2-Year Prospective Study. *World Neurosurg*. May 2017; 101: 493-497. PMID 28216399
46. Rappoport LH, Helsper K, Shirk T. Minimally invasive sacroiliac joint fusion using a novel hydroxyapatite-coated screw: final 2-year clinical and radiographic results. *J Spine Surg*. Jun 2021; 7(2): 155-161. PMID 34296027
47. Fuchs V, Ruhl B. Distraction arthrodesis of the sacroiliac joint: 2-year results of a descriptive prospective multi-center cohort study in 171 patients. *Eur Spine J*. Jan 2018; 27(1): 194-204. PMID 29058134
48. Calodney AK, Azeem N, Buchanan P, et al. Six Month Interim Outcomes from SECURE: A Single arm, Multicenter, Prospective, Clinical Study on a Novel Minimally Invasive Posterior Sacroiliac Fusion Device. *Expert Rev Med Devices*. May 2022; 19(5): 451-461. PMID 35724479
49. Kucharzyk D, Colle K, Boone C, et al. Clinical Outcomes Following Minimally Invasive Sacroiliac Joint Fusion With Decortication: The EVoluSlon Clinical Study. *Int J Spine Surg*. Feb 2022; 16(1): 168-175. PMID 35217586
50. Benzon HT, Connis RT, De Leon-Casasola OA, et al. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*. Apr 2010; 112(4): 810-33. PMID 20124882
51. Lee DW, Pritzlaff S, Jung MJ, et al. Latest Evidence-Based Application for Radiofrequency Neurotomy (LEARN): Best Practice Guidelines from the American Society of Pain and Neuroscience (ASPN). *J Pain Res*. 2021; 14: 2807-2831. PMID 34526815
52. Sayed D, Grider J, Strand N, et al. The American Society of Pain and Neuroscience (ASPN) Evidence-Based Clinical Guideline of Interventional Treatments for Low Back Pain. *J Pain Res*. 2022; 15: 3729-3832. PMID 36510616

53. National Institute for Health and Care Excellence. Minimally invasive sacroiliac joint fusion surgery for chronic sacroiliac pain [IPG578]. 2017; <https://www.nice.org.uk/guidance/ipg578>. Accessed October 4, 2022.
54. National Institute for Health and Care Excellence. iFuse for treating chronic sacroiliac joint pain [MTG39]. 2022; <https://www.nice.org.uk/guidance/mtg39>. Accessed October 4, 2023.