

Blue Cross Blue Shield of Massachusetts is an Independent Licenses of the Blue Cross and Blue Shield Association

# Medical Policy Open and Thoracoscopic Approaches to Treat Atrial Fibrillation and Atrial Flutter (Maze and Related Procedures)

# **Table of Contents**

- Policy: Commercial
- Policy: Medicare
- <u>Authorization Information</u>
- <u>Coding Information</u>

Policy History

- Description
- Information Pertaining to All Policies
- <u>References</u>

# Policy Number: 356

BCBSA Reference Number: 7.01.14 (For Plan internal use only) NCD/LCD: N/A

# **Related Policies**

- Catheter Ablation as Treatment of Atrial Fibrillation, #141
- Percutaneous Left Atrial Appendage Closure Devices for Stroke Prevention in Atrial Fibrillation, #334

# Policy

# Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO Blue<sup>SM</sup> and Medicare PPO Blue<sup>SM</sup> Members

The maze procedure or modified procedure, performed on a non-beating heart during cardiopulmonary bypass with concomitant cardiac surgery, is considered <u>MEDICALLY NECESSARY</u> for the treatment of symptomatic atrial fibrillation or flutter.

Stand-alone minimally invasive, off-pump maze procedures (ie, modified maze procedures), including those done via mini-thoracotomy, are considered **INVESTIGATIONAL** for treatment of atrial fibrillation or flutter.

Hybrid ablation (defined as a combined percutaneous and thoracoscopic approach) is considered **INVESTIGATIONAL** for the treatment of atrial fibrillation or flutter.

The use of an open maze or modified maze procedure performed on a non–beating heart during cardiopulmonary bypass without concomitant cardiac surgery is considered <u>NOT MEDICALLY</u> <u>NECESSARY</u> for treatment of symptomatic atrial fibrillation or flutter.

# **Prior Authorization Information**

### Inpatient

 For services described in this policy, precertification/preauthorization <u>IS REQUIRED</u> for all products if the procedure is performed <u>inpatient</u>.

Outpatient

• For services described in this policy, see below for products where prior authorization <u>might be</u> <u>required</u> if the procedure is performed <u>outpatient</u>.

	Outpatient
Commercial Managed Care (HMO and POS)	This procedure is performed in the inpatient setting
Commercial PPO and Indemnity	This procedure is performed in the inpatient setting
Medicare HMO Blue <sup>sM</sup>	This procedure is performed in the inpatient setting
Medicare PPO Blue <sup>sm</sup>	This procedure is performed in the inpatient setting

## **CPT Codes / HCPCS Codes / ICD Codes**

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above <u>medical necessity criteria MUST</u> be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

### **CPT Codes**

CPT codes:	Code Description
33256	Operative tissue ablation and reconstruction of atria, extensive (e.g., maze procedure); with cardiopulmonary bypass
33257	Operative tissue ablation and reconstruction of atria, performed at the time of other cardiac procedure(s), limited (e.g., modified maze procedure) (List separately in addition to code for primary procedure)
33259	Operative tissue ablation and reconstruction of atria, performed at the time of other cardiac procedure(s), extensive (e.g., maze procedure), with cardiopulmonary bypass (List separately in addition to code for primary procedure)

### **ICD-10 Procedure Codes**

ICD-10	
procedure	
codes:	Code Description
02560ZZ	Destruction of Right Atrium, Open Approach
02563ZZ	Destruction of Right Atrium, Percutaneous Approach
02570ZZ	Destruction of Left Atrium, Open Approach
02573ZZ	Destruction of Left Atrium, Percutaneous Approach

# The following CPT codes are considered investigational for <u>Commercial Members: Managed Care</u> (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

### **CPT Codes**

CPT codes:	Code Description
33254	Operative tissue ablation and reconstruction of atria, limited (e.g., modified maze procedure)
33255	Operative tissue ablation and reconstruction of atria, extensive (e.g., maze procedure); without cardiopulmonary bypass

33258	Operative tissue ablation and reconstruction of atria, performed at the time of other cardiac procedure(s), extensive (e.g., maze procedure), without cardiopulmonary bypass (List separately in addition to code for primary procedure)
33265	Endoscopy, surgical; operative tissue ablation and reconstruction of atria, limited (e.g., modified maze procedure), without cardiopulmonary bypass
33266	Endoscopy, surgical; operative tissue ablation and reconstruction of atria, limited (e.g., maze procedure), without cardiopulmonary bypass

## Description

### **Atrial Fibrillation**

Atrial fibrillation (AF) is a supraventricular tachyarrhythmia characterized by disorganized atrial activation with ineffective atrial ejection. The underlying mechanism of AF involves the interplay between electrical triggering events that initiate AF and the myocardial substrate that permits propagation and maintenance of the aberrant electrical circuit. The most common focal trigger of AF appears to be located within the cardiac muscle that extends into the pulmonary veins. The atria are frequently abnormal in patients with AF and demonstrate enlargement or increased conduction time. Atrial flutter is a variant of AF.

### Epidemiology

In the US, more than 3 to 6 million people have AF and it has been estimated that more than 12 million people will have AF by 2030.<sup>1,2,3,</sup> Age, body mass index, height, hypertension, diabetes mellitus, obstructive sleep apnea, myocardial infarction, heart failure, hyperthyroidism, chronic kidney disease, smoking, moderate to heavy alcohol consumption, and genetic predisposition are all risk factors for AF.<sup>4,3,</sup> Age-adjusted AF incidence and prevalence is higher among men than women, although the lifetime risk is similar at 24% for men and 22% for women<sup>5,</sup>. AF incidence and prevalence appear lower in individuals who are Black compared to White, despite a higher burden of comorbidities. However, this difference is likely largely explained by differential detection of AF by race/ethnicity.<sup>6,</sup>

#### Treatment

The first-line treatment for AF usually includes medications to maintain sinus rhythm and/or control the ventricular rate. Antiarrhythmic medications are only partially effective; therefore, medical treatment is not sufficient for many patients. Percutaneous catheter ablation, using endocardial ablation, is an accepted second-line treatment for patients who are not adequately controlled on medications and may also be used as first-line treatment. Catheter ablation (CA) is successful in maintaining sinus rhythm for most patients, but long-term recurrences are common and increase over time. Performed either by open surgical techniques or thoracoscopy, surgical ablation is an alternative approach to percutaneous CA.

### **Summary**

There are various surgical approaches to treat atrial fibrillation (AF) that work by interrupting abnormal electrical activity in the atria. Open surgical procedures, such as the Cox maze procedure were first developed for this purpose and are now generally performed in conjunction with valvular or coronary artery bypass graft surgery. Surgical techniques have evolved to include minimally invasive approaches that use epicardial radiofrequency ablation, a thoracoscopic or mediastinal approach, and hybrid catheter ablations/open procedures.

For individuals who have symptomatic AF or flutter who are undergoing cardiac surgery with bypass who received a Cox maze or a modified maze procedure, the evidence includes several randomized controlled trials (RCTs) and nonrandomized comparative studies, along with systematic reviews of these studies. Relevant outcomes are overall survival, medication use, and treatment-related morbidity. Several small RCTs have provided most of the direct evidence confirming the benefit of a modified maze procedure for patients with AF who are undergoing mitral valve surgery. These trials have established that the addition of a modified maze procedure results in a lower incidence of atrial arrhythmias following surgery, with minimal additional risks. Observational studies have supported these RCT findings. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have symptomatic, drug-resistant AF or flutter who are not undergoing cardiac surgery with bypass who receive minimally invasive, off-pump thoracoscopic maze procedures, the evidence includes RCTs and observational studies, some of which identify control groups. Relevant outcomes are overall survival, medication use, and treatment-related morbidity. Two RCTs reported significantly higher rates of freedom from AF at 1-year with surgical ablation but also reported significantly higher rates of serious adverse events. The remaining 2 RCTs found no significant differences between treatment groups in rates of freedom from AF and either did not assess or did not find significant differences in serious adverse events. The comparative observational studies consistently found significantly higher rates of freedom from atrial arrhythmias but lacked assessment of serious adverse events. The noncomparative studies generally only reported short-term outcomes and did not consistently report adverse events. Therefore, this evidence does not permit definitive conclusions about whether a specific approach is superior to the other. Factors, such as previous treatment, the probability of maintaining sinus rhythm, the risk of complications, contraindications to anticoagulation, and patient preference, may all affect the risk-benefit ratio for each procedure. Additionally, the studies do not permit conclusions about harm due to heterogeneous measurement across studies, with mixed results. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have symptomatic, drug-resistant AF or flutter who are not undergoing cardiac surgery with bypass who receive hybrid thoracoscopic and endocardial ablation procedures, the evidence includes 5 RCTs (sample sizes ranging from 41 to 154), comparative observational studies, single-arm case series, and systematic reviews of these studies. Evidence from randomized and nonrandomized studies found an increased rate of AF-free survival, reduced risk of cardioversion, and increased risk of periprocedural adverse events with hybrid procedures relative to standard ablation. The largest RCT (CEASE-AF) reported composite major complications at 1 year in 9% vs 6% with hybrid vs standard ablation. The surgical ablation lesion sets varied across studies and have not been standardized in practice. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Date	Action
5/2025	Annual policy review. Summary and references updated. Policy statements unchanged.
5/2024	Clarified prior authorization table
7/2023	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
7/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
10/2017	Annual policy review. "Drug resistant" removed from medically necessary statement and not medically necessary statement. Effective 10/1/2017.
7/2016	Annual policy review. New references added.
12/2015	Annual policy review. New not medically necessary indications described. The phrase "without concomitant cardiac surgery" was removed from the medically necessary policy statement. Title revised to include Atrial Flutter. Clarified coding information. Effective 12/1/2015.
9/2014	Annual policy review. New references added.
6/2014	Updated Coding section with ICD10 procedure and diagnosis codes. Effective 10/2015.
1/2014	Annual policy review. New investigational indications described. Effective 1/1/2014.

### **Policy History**

12/2013	Removed ICD-9 diagnosis codes 427.31, 427.32 as the policy is requires prior
7/2013	Annual policy review. Added "or modified" and "(i.e., modified maze procedures)" to
	the policy statements. Effective 7/1/2013.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No
	changes to policy statements.
4/2011	Reviewed - Medical Policy Group - Cardiology and Pulmonology. No changes to
	policy statements.
4/2010	Reviewed - Medical Policy Group - Cardiology and Pulmonology. No changes to
	policy statements.
4/2009	Reviewed - Medical Policy Group - Cardiology and Pulmonology. No changes to
	policy statements.
4/2008	Reviewed - Medical Policy Group - Cardiology and Pulmonology. No changes to
	policy statements.
8/2007	Annual policy review. Changes to policy statements.

# Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information: <u>Medical Policy Terms of Use</u> <u>Managed Care Guidelines</u> <u>Indemnity/PPO Guidelines</u> <u>Clinical Exception Process</u> <u>Medical Technology Assessment Guidelines</u>

### References

- Miyasaka Y, Barnes ME, Gersh BJ, et al. Secular trends in incidence of atrial fibrillation in Olmsted County, Minnesota, 1980 to 2000, and implications on the projections for future prevalence. Circulation. Jul 11 2006; 114(2): 119-25. PMID 16818816
- 2. Colilla S, Crow A, Petkun W, et al. Estimates of current and future incidence and prevalence of atrial fibrillation in the U.S. adult population. Am J Cardiol. Oct 15 2013; 112(8): 1142-7. PMID 23831166
- 3. Kornej J, Börschel CS, Benjamin EJ, et al. Epidemiology of Atrial Fibrillation in the 21st Century: Novel Methods and New Insights. Circ Res. Jun 19 2020; 127(1): 4-20. PMID 32716709
- 4. Benjamin EJ, Levy D, Vaziri SM, et al. Independent risk factors for atrial fibrillation in a populationbased cohort. The Framingham Heart Study. JAMA. Mar 16 1994; 271(11): 840-4. PMID 8114238
- 5. Heeringa J, van der Kuip DA, Hofman A, et al. Prevalence, incidence and lifetime risk of atrial fibrillation: the Rotterdam study. Eur Heart J. Apr 2006; 27(8): 949-53. PMID 16527828
- Heckbert SR, Austin TR, Jensen PN, et al. Differences by Race/Ethnicity in the Prevalence of Clinically Detected and Monitor-Detected Atrial Fibrillation: MESA. Circ Arrhythm Electrophysiol. Jan 2020; 13(1): e007698. PMID 31934795
- Calkins H, Hindricks G, Cappato R, et al. 2017 HRS/EHRA/ECAS/APHRS/SOLAECE expert consensus statement on catheter and surgical ablation of atrial fibrillation. Europace. Jan 01 2018; 20(1): e1-e160. PMID 29016840
- Sakurai Y, Kuno T, Yokoyama Y, et al. Late Survival Benefits of Concomitant Surgical Ablation for Atrial Fibrillation During Cardiac Surgery: A Systematic Review and Meta-Analysis. Am J Cardiol. Oct 28 2024; 235: 16-29. PMID 39471966
- Huffman MD, Karmali KN, Berendsen MA, et al. Concomitant atrial fibrillation surgery for people undergoing cardiac surgery. Cochrane Database Syst Rev. Aug 22 2016; 2016(8): CD011814. PMID 27551927
- 10. Phan K, Xie A, Tian DH, et al. Systematic review and meta-analysis of surgical ablation for atrial fibrillation during mitral valve surgery. Ann Cardiothorac Surg. Jan 2014; 3(1): 3-14. PMID 24516793
- 11. Gillinov AM, Gelijns AC, Parides MK, et al. Surgical ablation of atrial fibrillation during mitral-valve surgery. N Engl J Med. Apr 09 2015; 372(15): 1399-409. PMID 25853744
- 12. Budera P, Straka Z, Osmančík P, et al. Comparison of cardiac surgery with left atrial surgical ablation vs. cardiac surgery without atrial ablation in patients with coronary and/or valvular heart disease plus

atrial fibrillation: final results of the PRAGUE-12 randomized multicentre study. Eur Heart J. Nov 2012; 33(21): 2644-52. PMID 22930458

- Van Breugel HN, Nieman FH, Accord RE, et al. A prospective randomized multicenter comparison on health-related quality of life: the value of add-on arrhythmia surgery in patients with paroxysmal, permanent or persistent atrial fibrillation undergoing valvular and/or coronary bypass surgery. J Cardiovasc Electrophysiol. May 2010; 21(5): 511-20. PMID 19925605
- Kim KC, Cho KR, Kim YJ, et al. Long-term results of the Cox-Maze III procedure for persistent atrial fibrillation associated with rheumatic mitral valve disease: 10-year experience. Eur J Cardiothorac Surg. Feb 2007; 31(2): 261-6. PMID 17158057
- 15. Gerdisch M, Lehr E, Dunnington G, et al. Mid-term outcomes of concomitant Cox-Maze IV: Results from a multicenter prospective registry. J Card Surg. Oct 2022; 37(10): 3006-3013. PMID 35870185
- Damiano RJ, Badhwar V, Acker MA, et al. The CURE-AF trial: a prospective, multicenter trial of irrigated radiofrequency ablation for the treatment of persistent atrial fibrillation during concomitant cardiac surgery. Heart Rhythm. Jan 2014; 11(1): 39-45. PMID 24184028
- 17. Gaita F, Ebrille E, Scaglione M, et al. Very long-term results of surgical and transcatheter ablation of long-standing persistent atrial fibrillation. Ann Thorac Surg. Oct 2013; 96(4): 1273-1278. PMID 23915587
- Watkins AC, Young CA, Ghoreishi M, et al. Prospective assessment of the CryoMaze procedure with continuous outpatient telemetry in 136 patients. Ann Thorac Surg. Apr 2014; 97(4): 1191-8; discussion 1198. PMID 24582049
- McCarthy PM, Gerdisch M, Philpott J, et al. Three-year outcomes of the postapproval study of the AtriCure Bipolar Radiofrequency Ablation of Permanent Atrial Fibrillation Trial. J Thorac Cardiovasc Surg. Aug 2022; 164(2): 519-527.e4. PMID 33129501
- 20. van Laar C, Kelder J, van Putte BP. The totally thoracoscopic maze procedure for the treatment of atrial fibrillation. Interact Cardiovasc Thorac Surg. Jan 2017; 24(1): 102-111. PMID 27664426
- Yi S, Liu X, Wang W, et al. Thoracoscopic surgical ablation or catheter ablation for patients with atrial fibrillation? A systematic review and meta-analysis of randomized controlled trials. Interact Cardiovasc Thorac Surg. Dec 07 2020; 31(6): 763-773. PMID 33166993
- 22. Phan K, Phan S, Thiagalingam A, et al. Thoracoscopic surgical ablation versus catheter ablation for atrial fibrillation. Eur J Cardiothorac Surg. Apr 2016; 49(4): 1044-51. PMID 26003961
- Boersma LV, Castella M, van Boven W, et al. Atrial fibrillation catheter ablation versus surgical ablation treatment (FAST): a 2-center randomized clinical trial. Circulation. Jan 03 2012; 125(1): 23-30. PMID 22082673
- Castellá M, Kotecha D, van Laar C, et al. Thoracoscopic vs. catheter ablation for atrial fibrillation: long-term follow-up of the FAST randomized trial. Europace. May 01 2019; 21(5): 746-753. PMID 30715255
- 25. Pokushalov E, Romanov A, Elesin D, et al. Catheter versus surgical ablation of atrial fibrillation after a failed initial pulmonary vein isolation procedure: a randomized controlled trial. J Cardiovasc Electrophysiol. Dec 2013; 24(12): 1338-43. PMID 24016147
- Adiyaman A, Buist TJ, Beukema RJ, et al. Randomized Controlled Trial of Surgical Versus Catheter Ablation for Paroxysmal and Early Persistent Atrial Fibrillation. Circ Arrhythm Electrophysiol. Oct 2018; 11(10): e006182. PMID 30354411
- Haldar S, Khan HR, Boyalla V, et al. Catheter ablation vs. thoracoscopic surgical ablation in longstanding persistent atrial fibrillation: CASA-AF randomized controlled trial. Eur Heart J. Dec 14 2020; 41(47): 4471-4480. PMID 32860414
- Boyalla V, Haldar S, Khan H, et al. Long-term clinical outcomes and cost-effectiveness of catheter vs thoracoscopic surgical ablation in long-standing persistent atrial fibrillation using continuous cardiac monitoring: CASA-AF randomized controlled trial. Heart Rhythm. Sep 2024; 21(9): 1562-1569. PMID 38763376
- Kwon HJ, Choi JH, Kim HR, et al. Radiofrequency vs. Cryoballoon vs. Thoracoscopic Surgical Ablation for Atrial Fibrillation: A Single-Center Experience. Medicina (Kaunas). Sep 26 2021; 57(10). PMID 34684060
- 30. Mahapatra S, LaPar DJ, Kamath S, et al. Initial experience of sequential surgical epicardial-catheter endocardial ablation for persistent and long-standing persistent atrial fibrillation with long-term followup. Ann Thorac Surg. Jun 2011; 91(6): 1890-8. PMID 21619988

- Stulak JM, Dearani JA, Sundt TM, et al. Ablation of atrial fibrillation: comparison of catheter-based techniques and the Cox-Maze III operation. Ann Thorac Surg. Jun 2011; 91(6): 1882-8; discussion 1888-9. PMID 21619987
- 32. Wang J, Li Y, Shi J, et al. Minimally invasive surgical versus catheter ablation for the long-lasting persistent atrial fibrillation. PLoS One. 2011; 6(7): e22122. PMID 21765943
- Lawrance CP, Henn MC, Miller JR, et al. A minimally invasive Cox maze IV procedure is as effective as sternotomy while decreasing major morbidity and hospital stay. J Thorac Cardiovasc Surg. Sep 2014; 148(3): 955-61; discussion 962-2. PMID 25048635
- De Maat GE, Pozzoli A, Scholten MF, et al. Surgical minimally invasive pulmonary vein isolation for lone atrial fibrillation: midterm results of a multicenter study. Innovations (Phila). 2013; 8(6): 410-5. PMID 24356430
- 35. Massimiano PS, Yanagawa B, Henry L, et al. Minimally invasive fibrillating heart surgery: a safe and effective approach for mitral valve and surgical ablation for atrial fibrillation. Ann Thorac Surg. Aug 2013; 96(2): 520-7. PMID 23773732
- 36. Cui YQ, Li Y, Gao F, et al. Video-assisted minimally invasive surgery for lone atrial fibrillation: a clinical report of 81 cases. J Thorac Cardiovasc Surg. Feb 2010; 139(2): 326-32. PMID 19660413
- Edgerton JR, Brinkman WT, Weaver T, et al. Pulmonary vein isolation and autonomic denervation for the management of paroxysmal atrial fibrillation by a minimally invasive surgical approach. J Thorac Cardiovasc Surg. Oct 2010; 140(4): 823-8. PMID 20299028
- 38. Han FT, Kasirajan V, Kowalski M, et al. Results of a minimally invasive surgical pulmonary vein isolation and ganglionic plexi ablation for atrial fibrillation: single-center experience with 12-month follow-up. Circ Arrhythm Electrophysiol. Aug 2009; 2(4): 370-7. PMID 19808492
- Pruitt JC, Lazzara RR, Ebra G. Minimally invasive surgical ablation of atrial fibrillation: the thoracoscopic box lesion approach. J Interv Card Electrophysiol. Dec 2007; 20(3): 83-7. PMID 18214660
- 40. Sirak J, Jones D, Sun B, et al. Toward a definitive, totally thoracoscopic procedure for atrial fibrillation. Ann Thorac Surg. Dec 2008; 86(6): 1960-4. PMID 19022018
- 41. Speziale G, Bonifazi R, Nasso G, et al. Minimally invasive radiofrequency ablation of lone atrial fibrillation by monolateral right minithoracotomy: operative and early follow-up results. Ann Thorac Surg. Jul 2010; 90(1): 161-7. PMID 20609767
- 42. Wudel JH, Chaudhuri P, Hiller JJ. Video-assisted epicardial ablation and left atrial appendage exclusion for atrial fibrillation: extended follow-up. Ann Thorac Surg. Jan 2008; 85(1): 34-8. PMID 18154774
- 43. Yilmaz A, Geuzebroek GS, Van Putte BP, et al. Completely thoracoscopic pulmonary vein isolation with ganglionic plexus ablation and left atrial appendage amputation for treatment of atrial fibrillation. Eur J Cardiothorac Surg. Sep 2010; 38(3): 356-60. PMID 20227287
- Yilmaz A, Van Putte BP, Van Boven WJ. Completely thoracoscopic bilateral pulmonary vein isolation and left atrial appendage exclusion for atrial fibrillation. J Thorac Cardiovasc Surg. Aug 2008; 136(2): 521-2. PMID 18692667
- 45. Geuzebroek GS, Bentala M, Molhoek SG, et al. Totally thoracoscopic left atrial Maze: standardized, effective and safe. Interact Cardiovasc Thorac Surg. Mar 2016; 22(3): 259-64. PMID 26705300
- 46. Vos LM, Bentala M, Geuzebroek GS, et al. Long-term outcome after totally thoracoscopic ablation for atrial fibrillation. J Cardiovasc Electrophysiol. Jan 2020; 31(1): 40-45. PMID 31691391
- Harlaar N, Oudeman MA, Trines SA, et al. Long-term follow-up of thoracoscopic ablation in longstanding persistent atrial fibrillation. Interact Cardiovasc Thorac Surg. Jun 01 2022; 34(6): 990-998. PMID 34957518
- Ad N, Henry L, Hunt S, et al. The outcome of the Cox Maze procedure in patients with previous percutaneous catheter ablation to treat atrial fibrillation. Ann Thorac Surg. May 2011; 91(5): 1371-7; discussion 1377. PMID 21457939
- Castellá M, Pereda D, Mestres CA, et al. Thoracoscopic pulmonary vein isolation in patients with atrial fibrillation and failed percutaneous ablation. J Thorac Cardiovasc Surg. Sep 2010; 140(3): 633-8. PMID 20117799
- 50. MacGregor RM, Bakir NH, Pedamallu H, et al. Late results after stand-alone surgical ablation for atrial fibrillation. J Thorac Cardiovasc Surg. Nov 2022; 164(5): 1515-1528.e8. PMID 34045056

- Bisleri G, Rosati F, Bontempi L, et al. Hybrid approach for the treatment of long-standing persistent atrial fibrillation: electrophysiological findings and clinical results. Eur J Cardiothorac Surg. Nov 2013; 44(5): 919-23. PMID 23475587
- 52. Winkle RA, Mead RH, Engel G, et al. Very long term outcomes of atrial fibrillation ablation. Heart Rhythm. May 2023; 20(5): 680-688. PMID 36764350
- Eranki A, Wilson-Smith A, Flynn C, et al. Mid term freedom from atrial fibrillation following hybrid ablation, a systematic review and meta analysis. J Cardiothorac Surg. Apr 19 2023; 18(1): 155. PMID 37076929
- Mhanna M, Beran A, Al-Abdouh A, et al. Hybrid convergent ablation versus endocardial catheter ablation for atrial fibrillation: A systematic review and meta-analysis. J Arrhythm. Dec 2021; 37(6): 1459-1467. PMID 34887950
- 55. Eranki A, Wilson-Smith AR, Williams ML, et al. Hybrid convergent ablation versus endocardial catheter ablation for atrial fibrillation: a systematic review and meta-analysis of randomised control trials and propensity matched studies. J Cardiothorac Surg. Aug 13 2022; 17(1): 181. PMID 35964093
- DeLurgio DB, Crossen KJ, Gill J, et al. Hybrid Convergent Procedure for the Treatment of Persistent and Long-Standing Persistent Atrial Fibrillation: Results of CONVERGE Clinical Trial. Circ Arrhythm Electrophysiol. Dec 2020; 13(12): e009288. PMID 33185144
- 57. Lee KN, Kim DY, Boo KY, et al. Combined epicardial and endocardial approach for redo radiofrequency catheter ablation in patients with persistent atrial fibrillation: a randomized clinical trial. Europace. Oct 13 2022; 24(9): 1412-1419. PMID 35640923
- van der Heijden CAJ, Weberndörfer V, Vroomen M, et al. Hybrid Ablation Versus Repeated Catheter Ablation in Persistent Atrial Fibrillation: A Randomized Controlled Trial. JACC Clin Electrophysiol. Jul 2023; 9(7 Pt 2): 1013-1023. PMID 36752455
- 59. Doll N, Weimar T, Kosior DA, et al. Efficacy and safety of hybrid epicardial and endocardial ablation versus endocardial ablation in patients with persistent and longstanding persistent atrial fibrillation: a randomised, controlled trial. EClinicalMedicine. Jul 2023; 61: 102052. PMID 37425372
- 60. Jan M, Žižek D, Geršak ŽM, et al. Comparison of treatment outcomes between convergent procedure and catheter ablation for paroxysmal atrial fibrillation evaluated with implantable loop recorder monitoring. J Cardiovasc Electrophysiol. Aug 2018; 29(8): 1073-1080. PMID 29722468
- DeLurgio DB, Blauth C, Halkos ME, et al. Hybrid epicardial-endocardial ablation for long-standing persistent atrial fibrillation: A subanalysis of the CONVERGE Trial. Heart Rhythm O2. Feb 2023; 4(2): 111-118. PMID 36873309
- Kress DC, Erickson L, Choudhuri I, et al. Comparative Effectiveness of Hybrid Ablation Versus Endocardial Catheter Ablation Alone in Patients With Persistent Atrial Fibrillation. JACC Clin Electrophysiol. Apr 2017; 3(4): 341-349. PMID 29759446
- Maclean E, Yap J, Saberwal B, et al. The convergent procedure versus catheter ablation alone in longstanding persistent atrial fibrillation: A single centre, propensity-matched cohort study. Int J Cardiol. Mar 15 2020; 303: 49-53. PMID 32063280
- 64. Pannone L, Mouram S, Della Rocca DG, et al. Hybrid atrial fibrillation ablation: long-term outcomes from a single-centre 10-year experience. Europace. May 19 2023; 25(5). PMID 37246904
- Mannakkara NN, Porter B, Child N, et al. Convergent ablation for persistent atrial fibrillation: outcomes from a single-centre real-world experience. Eur J Cardiothorac Surg. Dec 02 2022; 63(1). PMID 36346176
- Kiankhooy A, Pierce C, Burk S, et al. Hybrid ablation of persistent and long-standing persistent atrial fibrillation with depressed ejection fraction: A single-center observational study. JTCVS Open. Dec 2022; 12: 137-146. PMID 36590727
- 67. Gehi AK, Mounsey JP, Pursell I, et al. Hybrid epicardial-endocardial ablation using a pericardioscopic technique for the treatment of atrial fibrillation. Heart Rhythm. Jan 2013; 10(1): 22-8. PMID 23064043
- 68. Gersak B, Pernat A, Robic B, et al. Low rate of atrial fibrillation recurrence verified by implantable loop recorder monitoring following a convergent epicardial and endocardial ablation of atrial fibrillation. J Cardiovasc Electrophysiol. Oct 2012; 23(10): 1059-66. PMID 22587585
- La Meir M, Gelsomino S, Lorusso R, et al. The hybrid approach for the surgical treatment of lone atrial fibrillation: one-year results employing a monopolar radiofrequency source. J Cardiothorac Surg. Jul 19 2012; 7: 71. PMID 22812613

- Muneretto C, Bisleri G, Bontempi L, et al. Successful treatment of lone persistent atrial fibrillation by means of a hybrid thoracoscopic-transcatheter approach. Innovations (Phila). 2012; 7(4): 254-8. PMID 23123991
- Muneretto C, Bisleri G, Bontempi L, et al. Durable staged hybrid ablation with thoracoscopic and percutaneous approach for treatment of long-standing atrial fibrillation: a 30-month assessment with continuous monitoring. J Thorac Cardiovasc Surg. Dec 2012; 144(6): 1460-5; discussion 1465. PMID 23062968
- 72. Pison L, La Meir M, van Opstal J, et al. Hybrid thoracoscopic surgical and transvenous catheter ablation of atrial fibrillation. J Am Coll Cardiol. Jul 03 2012; 60(1): 54-61. PMID 22742400
- Zembala M, Filipiak K, Kowalski O, et al. Minimally invasive hybrid ablation procedure for the treatment of persistent atrial fibrillation: one year results. Kardiol Pol. 2012; 70(8): 819-28. PMID 22933215
- Geršak B, Zembala MO, Müller D, et al. European experience of the convergent atrial fibrillation procedure: multicenter outcomes in consecutive patients. J Thorac Cardiovasc Surg. Apr 2014; 147(4): 1411-6. PMID 23988287
- 75. Civello KC, Smith CA, Boedefeld W. Combined endocardial and epicardial ablation for symptomatic atrial fibrillation: single center experience in 100+ consecutive patients. J Innovations Cardiac Rhythm Manage. 2013;August.
- Tonks R, Lantz G, Mahlow J, et al. Short and Intermediate Term Outcomes of the Convergent Procedure: Initial Experience in a Tertiary Referral Center. Ann Thorac Cardiovasc Surg. Feb 20 2020; 26(1): 13-21. PMID 31495813
- Carpenter A, Pannell LMK, Rizvi SIA, et al. Convergent approach to persistent atrial fibrillation ablation: long-term single-centre safety and efficacy. Front Cardiovasc Med. 2023; 10: 1336801. PMID 38390303
- Wyler von Ballmoos MC, Hui DS, Mehaffey JH, et al. The Society of Thoracic Surgeons 2023 Clinical Practice Guidelines for the Surgical Treatment of Atrial Fibrillation. Ann Thorac Surg. Jan 27 2024. PMID 38286206
- 79. Joglar JA, Chung MK, Armbruster AL, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Circulation. Jan 02 2024; 149(1): e1-e156. PMID 38033089
- Tzeis S, Gerstenfeld EP, Kalman J, et al. 2024 European Heart Rhythm Association/Heart Rhythm Society/Asia Pacific Heart Rhythm Society/Latin American Heart Rhythm Society expert consensus statement on catheter and surgical ablation of atrial fibrillation. Europace. Mar 30 2024; 26(4). PMID 38587017
- 81. Ad N, Damiano RJ, Badhwar V, et al. Expert consensus guidelines: Examining surgical ablation for atrial fibrillation. J Thorac Cardiovasc Surg. Jun 2017; 153(6): 1330-1354.e1. PMID 28390766