



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Computed Tomography Perfusion Imaging of the Brain

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 448

BCBSA Reference Number: 6.01.49

NCD/LCD: N/A

Related Policies

Endovascular Procedures for Intracranial Arterial Disease (Atherosclerosis and Aneurysms), #323

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Computed tomography (CT)-based perfusion imaging may be considered **MEDICALLY NECESSARY** to select patients with anterior large-vessel stroke for mechanical embolectomy.

CT-based perfusion imaging of the brain is **INVESTIGATIONAL** for all other indications.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is not required .
Commercial PPO and Indemnity	Prior authorization is not required .
Medicare HMO Blue SM	Prior authorization is not required .
Medicare PPO Blue SM	Prior authorization is not required .

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above **medical necessity criteria MUST** be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
0042T	Cerebral perfusion analysis using computed tomography with contrast administration, including post-processing of parametric maps with determination of cerebral blood flow, cerebral blood volume, and mean transit time

The following ICD Diagnosis Codes are considered medically necessary when submitted with the CPT codes above if **medical necessity criteria** are met:

ICD-10 Diagnosis Codes

ICD-10-CM diagnosis codes:	Code Description
I63.00	Cerebral infarction due to thrombosis of unspecified precerebral artery
I63.011	Cerebral infarction due to thrombosis of right vertebral artery
I63.012	Cerebral infarction due to thrombosis of left vertebral artery
I63.013	Cerebral infarction due to thrombosis of bilateral vertebral arteries
I63.019	Cerebral infarction due to thrombosis of unspecified vertebral artery
I63.02	Cerebral infarction due to thrombosis of basilar artery
I63.031	Cerebral infarction due to thrombosis of right carotid artery
I63.032	Cerebral infarction due to thrombosis of left carotid artery
I63.033	Cerebral infarction due to thrombosis of bilateral carotid arteries
I63.039	Cerebral infarction due to thrombosis of unspecified carotid artery
I63.09	Cerebral infarction due to thrombosis of other precerebral artery
I63.10	Cerebral infarction due to embolism of unspecified precerebral artery
I63.111	Cerebral infarction due to embolism of right vertebral artery
I63.112	Cerebral infarction due to embolism of left vertebral artery
I63.113	Cerebral infarction due to embolism of bilateral vertebral arteries
I63.119	Cerebral infarction due to embolism of unspecified vertebral artery
I63.12	Cerebral infarction due to embolism of basilar artery
I63.131	Cerebral infarction due to embolism of right carotid artery
I63.132	Cerebral infarction due to embolism of left carotid artery
I63.133	Cerebral infarction due to embolism of bilateral carotid arteries
I63.139	Cerebral infarction due to embolism of unspecified carotid artery
I63.19	Cerebral infarction due to embolism of other precerebral artery
I63.20	Cerebral infarction due to unspecified occlusion or stenosis of unspecified precerebral arteries
I63.211	Cerebral infarction due to unspecified occlusion or stenosis of right vertebral artery
I63.212	Cerebral infarction due to unspecified occlusion or stenosis of left vertebral artery

I63.213	Cerebral infarction due to unspecified occlusion or stenosis of bilateral vertebral arteries
I63.219	Cerebral infarction due to unspecified occlusion or stenosis of unspecified vertebral artery
I63.22	Cerebral infarction due to unspecified occlusion or stenosis of basilar artery
I63.231	Cerebral infarction due to unspecified occlusion or stenosis of right carotid arteries
I63.232	Cerebral infarction due to unspecified occlusion or stenosis of left carotid arteries
I63.233	Cerebral infarction due to unspecified occlusion or stenosis of bilateral carotid arteries
I63.239	Cerebral infarction due to unspecified occlusion or stenosis of unspecified carotid artery
I63.29	Cerebral infarction due to unspecified occlusion or stenosis of other precerebral arteries
I63.30	Cerebral infarction due to thrombosis of unspecified cerebral artery
I63.311	Cerebral infarction due to thrombosis of right middle cerebral artery
I63.312	Cerebral infarction due to thrombosis of left middle cerebral artery
I63.313	Cerebral infarction due to thrombosis of bilateral middle cerebral arteries
I63.319	Cerebral infarction due to thrombosis of unspecified middle cerebral artery
I63.321	Cerebral infarction due to thrombosis of right anterior cerebral artery
I63.322	Cerebral infarction due to thrombosis of left anterior cerebral artery
I63.323	Cerebral infarction due to thrombosis of bilateral anterior cerebral arteries
I63.329	Cerebral infarction due to thrombosis of unspecified anterior cerebral artery
I63.331	Cerebral infarction due to thrombosis of right posterior cerebral artery
I63.332	Cerebral infarction due to thrombosis of left posterior cerebral artery
I63.333	Cerebral infarction due to thrombosis of bilateral posterior cerebral arteries
I63.339	Cerebral infarction due to thrombosis of unspecified posterior cerebral artery
I63.341	Cerebral infarction due to thrombosis of right cerebellar artery
I63.342	Cerebral infarction due to thrombosis of left cerebellar artery
I63.343	Cerebral infarction due to thrombosis of bilateral cerebellar arteries
I63.349	Cerebral infarction due to thrombosis of unspecified cerebellar artery
I63.39	Cerebral infarction due to thrombosis of other cerebral artery
I63.40	Cerebral infarction due to embolism of unspecified cerebral artery
I63.411	Cerebral infarction due to embolism of right middle cerebral artery
I63.412	Cerebral infarction due to embolism of left middle cerebral artery
I63.413	Cerebral infarction due to embolism of bilateral middle cerebral arteries
I63.419	Cerebral infarction due to embolism of unspecified middle cerebral artery
I63.421	Cerebral infarction due to embolism of right anterior cerebral artery
I63.422	Cerebral infarction due to embolism of left anterior cerebral artery
I63.423	Cerebral infarction due to embolism of bilateral anterior cerebral arteries
I63.429	Cerebral infarction due to embolism of unspecified anterior cerebral artery
I63.431	Cerebral infarction due to embolism of right posterior cerebral artery
I63.432	Cerebral infarction due to embolism of left posterior cerebral artery
I63.433	Cerebral infarction due to embolism of bilateral posterior cerebral arteries
I63.439	Cerebral infarction due to embolism of unspecified posterior cerebral artery
I63.441	Cerebral infarction due to embolism of right cerebellar artery
I63.442	Cerebral infarction due to embolism of left cerebellar artery
I63.443	Cerebral infarction due to embolism of bilateral cerebellar arteries

I63.449	Cerebral infarction due to embolism of unspecified cerebellar artery
I63.49	Cerebral infarction due to embolism of other cerebral artery
I63.50	Cerebral infarction due to unspecified occlusion or stenosis of unspecified cerebral artery
I63.511	Cerebral infarction due to unspecified occlusion or stenosis of right middle cerebral artery
I63.512	Cerebral infarction due to unspecified occlusion or stenosis of left middle cerebral artery
I63.513	Cerebral infarction due to unspecified occlusion or stenosis of bilateral middle cerebral arteries
I63.519	Cerebral infarction due to unspecified occlusion or stenosis of unspecified middle cerebral artery
I63.521	Cerebral infarction due to unspecified occlusion or stenosis of right anterior cerebral artery
I63.522	Cerebral infarction due to unspecified occlusion or stenosis of left anterior cerebral artery
I63.523	Cerebral infarction due to unspecified occlusion or stenosis of bilateral anterior cerebral arteries
I63.529	Cerebral infarction due to unspecified occlusion or stenosis of unspecified anterior cerebral artery
I63.531	Cerebral infarction due to unspecified occlusion or stenosis of right posterior cerebral artery
I63.532	Cerebral infarction due to unspecified occlusion or stenosis of left posterior cerebral artery
I63.533	Cerebral infarction due to unspecified occlusion or stenosis of bilateral posterior cerebral arteries
I63.539	Cerebral infarction due to unspecified occlusion or stenosis of unspecified posterior cerebral artery
I63.541	Cerebral infarction due to unspecified occlusion or stenosis of right cerebellar artery
I63.542	Cerebral infarction due to unspecified occlusion or stenosis of left cerebellar artery
I63.543	Cerebral infarction due to unspecified occlusion or stenosis of bilateral cerebellar arteries
I63.549	Cerebral infarction due to unspecified occlusion or stenosis of unspecified cerebellar artery
I63.59	Cerebral infarction due to unspecified occlusion or stenosis of other cerebral artery
I63.6	Cerebral infarction due to cerebral venous thrombosis, nonpyogenic
I63.81	Other cerebral infarction due to occlusion or stenosis of small artery
I63.89	Other cerebral infarction
I63.9	Cerebral infarction, unspecified

DESCRIPTION

Acute Stroke

The goal of acute stroke thrombolytic treatment is to rescue the ischemic penumbra, an area of the brain that surrounds the infarct core and is hypoperfused but does not die quickly. Multimodal computed tomography (CT) and magnetic resonance imaging (MRI) can be used to assess the cerebral parenchyma, vasculature, and tissue viability in the acute ischemic stroke setting and are used to detect ischemic tissue and exclude hemorrhage and other conditions that mimic acute cerebral ischemia. Non-contrast CT is used to rule out intracranial hemorrhage, tumor, or infection. Diffusion-weighted MRI is used to identify acute infarction, and a gradient-recalled echo sequence is used to exclude intracerebral hemorrhage.

CT angiography and magnetic resonance angiography are used to evaluate intra- and extracranial vasculature to detect the vascular occlusion and potentially guide therapy (eg, intravenous thrombolysis or mechanical thrombectomy).

The approved therapy, use of an intravenous tissue plasminogen activator, requires only a non-contrast CT scan to exclude the presence of hemorrhage (a contraindication to use of the drug). Current guidelines are to administer tissue plasminogen activator within the first three hours after an ischemic event, preceded by a CT scan. Many patients, however, do not present to the emergency department within the three-hour window, and thrombolysis carries a risk of intracranial hemorrhage. Thus, more sophisticated imaging may be needed to select the proper use of intra-arterial thrombolysis or mechanical thrombectomy in patients who present more than three hours after an ischemic stroke. Perfusion imaging is also being evaluated in the management of other neurologic conditions, such as subarachnoid hemorrhage and head trauma.

The potential utility of perfusion imaging for acute stroke is as follows:

- identification of brain regions with extremely low cerebral blood flow, which represent the core
- identification of patients with at-risk brain regions (acutely ischemic but viable penumbra) that may be salvageable with successful intra-arterial thrombolysis beyond the standard three-hour window
- triage of patients with at-risk brain regions to other available therapies, such as induced hypertension or mechanical clot retrieval
- decisions regarding intensive monitoring of patients with large, abnormally perfused brain regions
- biologically based management of patients who awaken with a stroke for which the precise time of onset is unknown.

Additional potential uses of CT perfusion (CTP) in acute stroke may include the following:

- detection and differential diagnosis (eg, excluding stroke mimics such as a transient ischemic attack, complex migraine, seizure, conversion disorders, hypoglycemia, brain tumors)
- determination of stroke subtype
- determination of stroke extent, including additional vascular territories at risk
- identification of patients at high early risk of stroke following a transient ischemic attack
- determining the need for blood pressure management
- establishing prognosis.

Similar information can be provided by CT and MRI regarding infarct core and penumbra. However, multimodal CT has a short protocol time (5-6 minutes) and, because it can be performed with any modern CT equipment, is more widely available in the emergency department setting. CTP is performed by capturing images as an iodinated contrast agent bolus passes through the cerebral circulation and accumulates in the cerebral tissues. (Older perfusion methodologies such as single-photon emission CT and xenon-enhanced CT scanning use a diffusible tracer.) The quantitative perfusion parameters are calculated from density changes for each pixel over time with the commercially available deconvolution-based software, in which cerebral blood flow is equal to regional cerebral blood volume divided by mean transit time. CT angiography and CTP imaging require ionizing radiation and iodinated contrast. It is estimated that typical CTP imaging deposits a slightly greater radiation dose than a routine unenhanced head CT (≈ 3.3 mSv).

Subarachnoid Hemorrhage and Cerebral Vasospasm

Cerebral vasospasm is a major cause of morbidity and mortality following aneurysmal SAH in patients who survive the initial hemorrhage and can be seen in about two-thirds of patients with aneurysmal SAH. The typical onset of cerebral vasospasm occurs 3 to 5 days after hemorrhage, with maximal narrowing on digital subtraction angiography at 5 to 14 days. Currently, the diagnosis of vasospasm and the management decisions rely on clinical examination, transcranial Doppler sonography, and digital subtraction angiography. Although symptomatic vasospasm affects 20% to 30% of patients with aneurysmal SAH, not all patients with angiographic vasospasm manifest clinical symptoms, and the symptoms can be nonspecific. Also, patients do not always have both clinical and imaging findings of

vasospasm. Due to these limitations, more accurate and reliable methods to detect cerebral vasospasm are being investigated.

Brain Tumors

The current standard for tumor grading is a histopathologic assessment of tissue. Limitations of histologic assessment include sampling error due to regional heterogeneity and interobserver variation. These limitations can result in inaccurate classification and grading of gliomas. Because malignant brain tumors are characterized by neovascularity and increased angiogenic activity, perfusion imaging has been proposed as a method to assess tumor grade and prognosis. Also, perfusion imaging can be repeated and may help to assess the evolution of tumors and the treatment response. Traditionally, perfusion imaging of brain tumors has been performed with MRI, which can estimate tumor blood volume, blood flow, and permeability. More recently, CTP imaging has been investigated for glioma grading. Potential advantages, compared with magnetic resonance perfusion, include the wider availability, faster scanning times, and lower cost. CTP imaging may also be used to distinguish recurrent tumor from radiation necrosis.

Summary

Computed tomography perfusion (CTP) imaging provides an assessment of cerebral blood flow that may help identify ischemic regions of the brain. This technology is proposed to aid treatment decisions in patients being evaluated for acute ischemic stroke, subarachnoid hemorrhage, cerebral vasospasm, brain tumors, and head trauma.

Acute Stroke

For individuals who have acute stroke who are being evaluated for thrombolysis who receive CTP imaging, the evidence includes a systematic review with meta-analysis, a randomized controlled trial (RCT), and cohort studies. The relevant outcomes are overall survival (OS), test accuracy, symptoms, morbid events, and functional outcomes. One potential area of benefit is greater individualization of therapy for acute stroke by better defining at-risk ischemic areas that may benefit from thrombolysis. Evidence from nonrandomized comparative studies has suggested that outcomes after thrombolysis are better in patients who have target mismatch on perfusion imaging than in patients without target mismatch and that patients with target mismatch treated after a three-hour time window have outcomes similar to patients treated within three hours. However, the therapeutic changes that would be associated with identifying specific target mismatch pattern on CTP are not well-defined. Additionally, although available evidence from the RCT suggests some modest benefit for acute stroke patients who receive CTP or magnetic resonance imaging and receive alteplase up to nine hours post-stroke, the overall net health outcome is unclear because there was also a lack of significant benefit on the secondary outcome of functional improvement and a trend toward increased risk of symptomatic intracranial hemorrhage and there were important limitations in relevance and potential limitations in statistical power. Therefore, RCTs are needed to determine with greater certainty whether a strategy employing CTP imaging improves health outcomes compared with traditional strategies for the treatment of acute stroke. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have acute anterior large-vessel stroke who are being evaluated for mechanical embolectomy who receive CTP imaging, the evidence includes RCTs and cohort studies. The relevant outcomes are OS, test accuracy, symptoms, morbid events, and functional outcomes. CTP is one of the several approaches used in acute stroke to define viable ischemic tissue better and therefore identify patients who might benefit from mechanical endovascular intervention. Alternative methods of patient selection for mechanical embolectomy have included time from stroke onset, multiphase computed tomography angiography, or Alberta Stroke Program Early CT Score. Three RCTs showed improved outcomes with mechanical embolectomy when patients were selected based on CTP results within 6 hours, at 6 to 16 hours, and at 6 to 24 hours. The evidence is sufficient to quantitatively determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have acute stroke who are being evaluated for prognosis who receive CTP imaging, the evidence includes retrospective analyses of large randomized trials. The relevant outcomes are OS,

test accuracy, symptoms, morbid events, and functional outcomes. Retrospective analysis of data from the MR CLEAN and DUST trials have found that the ischemic core detected on CTP imaging was predictive of functional outcomes. However, analysis of data from the DUST study found no improvement in a prediction model when CTP imaging was added to a basic model that used only patient characteristics and non-contrast computed tomography. The evidence is insufficient to determine the effects of the technology on health outcomes.

Subarachnoid Hemorrhage

For individuals who have SAH and cerebral vasospasm who receive CTP imaging, the evidence includes a systematic review with meta-analysis and a cohort study. The relevant outcomes are OS, test accuracy, symptoms, morbid events, and functional outcomes. CTP imaging is being evaluated for the diagnosis of vasospasm and delayed cerebral ischemia following aneurysmal SAH. One prospective study showed a qualitative measure of cerebral blood flow to have 93% accuracy for the detection of delayed cerebral ischemia, with lower accuracy for cerebral blood volume. Prospective trials are needed to determine whether CTP imaging in patients with aneurysmal SAH leads to the early identification of patients at high-risk for vasospasm or delayed cerebral ischemia, alters treatment decisions, and improves health outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

Brain Tumors

For individuals who have brain tumors who receive CTP imaging, the evidence includes studies on diagnostic accuracy. The relevant outcomes are test accuracy, symptoms, morbid events, and functional outcomes. For indications such as brain tumors and head trauma, the data on CTP imaging is limited. One study assessed the diagnostic accuracy of CTP imaging to differentiate high-grade from low-grade gliomas. Prospective studies in an appropriate population of patients are needed to evaluate the sensitivity and specificity of CTP glioma grading, with a histopathologic assessment of tumors as the independent reference standard. One prospective study performed a receiver operating characteristic curve analysis to evaluate the diagnostic accuracy of volume perfusion computed tomography. This is the first report using volume perfusion computed tomography to differentiate gliomas; therefore, replication of these findings in an independent sample of patients is needed as well as clarification of the clinical utility of this information. Studies showing the consistency in the thresholds used are needed as are studies showing improvement in health outcomes with CTP imaging. No recent reports on the use of CTP imaging for the evaluation of brain tumors have been identified. The evidence is insufficient to determine the effects of the technology on health outcomes.

Policy History

Date	Action
10/2019	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
10/2018	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
10/2018	Clarified coding information.
10/2017	New references added from BCBSA National medical policy.
10/2017	Clarified coding information.
11/2016	New references added from BCBSA National medical policy.
10/2016	Clarified coding information.
3/2016	BCBSA National medical policy review. CT perfusion considered medically necessary in patients with anterior large-vessel stroke being evaluated for mechanical embolectomy. CT perfusion in other situations remains investigational. Clarified coding information. Effective 3/1/2016.
11/2015	New references added from BCBSA National medical policy.
10/2014	New references added from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
1/2012	BCBSA National medical policy review.

	No changes to policy statements.
7/2011	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
1/2011	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
9/2010	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
7/2010	BCBSA National medical policy review. No changes to policy statements.
1/2010	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
9/2009	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
7/2009	BCBSA National medical policy review. No changes to policy statements.
1/2009	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
10/2008	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
5/2008	BCBSA National medical policy review. No changes to policy statements.
1/2007	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
9/2007	Reviewed - Medical Policy Group - Hematology and Oncology. No changes to policy statements.
4/2007	BCBSA National medical policy review. No changes to policy statements.
1/2007	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Burton KR, Dhanoa D, Aviv RI, et al. Perfusion CT for selecting patients with acute ischemic stroke for intravenous thrombolytic therapy. *Radiology*. Jan 2015;274(1):103-114. PMID 25243539.
2. Bivard A, Levi C, Krishnamurthy V, et al. Perfusion computed tomography to assist decision making for stroke thrombolysis. *Brain*. Jul 2015;138(Pt 7):1919-1931. PMID 25808369.
3. Schaefer PW, Souza L, Kamalian S, et al. Limited reliability of computed tomographic perfusion acute infarct volume measurements compared with diffusion-weighted imaging in anterior circulation stroke. *Stroke*. Feb 2015;46(2):419-424. PMID 25550366.
4. Liebeskind DS, Parsons MW, Wintermark M, et al. Computed tomography perfusion in acute ischemic stroke: is it ready for prime time? *Stroke*. Aug 2015;46(8):2364-2367. PMID 26159791.
5. Ma H, Campbell BCV, Parsons MW et al. Thrombolysis Guided by Perfusion Imaging up to 9 Hours after Onset of Stroke. *N. Engl. J. Med.*, 2019 May 9;380(19). PMID 31067369.
6. Garcia-Bermejo P, Calleja AI, Perez-Fernandez S, et al. Perfusion computed tomography-guided intravenous thrombolysis for acute ischemic stroke beyond 4.5 hours: a case-control study. *Cerebrovasc Dis*. Jul 2012;34(1):31-37. PMID 22759450.

7. Sztrihai LK, Manawadu D, Jarosz J, et al. Safety and clinical outcome of thrombolysis in ischaemic stroke using a perfusion CT mismatch between 3 and 6 hours. *PLoS One*. Oct 2011;6(10):e25796. PMID 22016775.
8. Campbell BC, Mitchell PJ, Kleinig TJ, et al. Endovascular therapy for ischemic stroke with perfusion-imaging selection. *N Engl J Med*. Mar 12 2015;372(11):1009-1018. PMID 25671797.
9. Albers GW, Marks MP, Kemp S et al. Thrombectomy for Stroke at 6 to 16 Hours with Selection by Perfusion Imaging. *N. Engl. J. Med.*, 2018 Jan 25;378(8). PMID 29364767.
10. Nogueira RG, Jadhav AP, Haussen DC et al. Thrombectomy 6 to 24 Hours after Stroke with a Mismatch between Deficit and Infarct. *N. Engl. J. Med.*, 2017 Nov 14;378(1). PMID 29129157.
11. Berkhemer OA, Fransen PS, Beumer D, et al. A randomized trial of intraarterial treatment for acute ischemic stroke. *N Engl J Med*. Jan 1 2015;372(1):11-20. PMID 25517348.
12. Goyal M, Demchuk AM, Menon BK, et al. Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med*. Mar 12 2015;372(11):1019-1030. PMID 25671798.
13. Saver JL, Goyal M, Bonafe A, et al. Solitaire with the Intention for Thrombectomy as Primary Endovascular Treatment for Acute Ischemic Stroke (SWIFT PRIME) trial: protocol for a randomized, controlled, multicenter study comparing the Solitaire revascularization device with IV tPA alone in acute ischemic stroke. *Int J Stroke*. Apr 2015;10(3):439-448. PMID 25777831.
14. Borst J, Berkhemer OA, Roos YB, et al. Value of computed tomographic perfusion-based patient selection for intra-arterial acute ischemic stroke treatment. *Stroke*. Dec 2015;46(12):3375-3382. PMID 26542698.
15. Rai AT, Raghuram K, Domico J, et al. Pre-intervention triage incorporating perfusion imaging improves outcomes in patients undergoing endovascular stroke therapy: a comparison with the device trials. *J Neurointerv Surg*. Mar 2013;5(2):121-127. PMID 22345110.
16. Vanicek J, Cimflova P, Bulik M et al. Single-Centre Experience with Patients Selection for Mechanical Thrombectomy Based on Automated Computed Tomography Perfusion Analysis-A Comparison with Computed TomographyCT Perfusion Thrombectomy Trials. *J Stroke Cerebrovasc Dis*, 2019 Jan 19;28(4). PMID 30655040.
17. Lansberg MG, Christensen S, Kemp S, et al. Computed tomographic perfusion to Predict Response to Recanalization in ischemic stroke. *Ann Neurol*. Jun 2017;81(6):849-856. PMID 28486789.
18. Dankbaar JW, Horsch AD, van den Hoven AF, et al. Prediction of clinical outcome after acute ischemic stroke: the value of repeated noncontrast computed tomography, computed tomographic angiography, and computed tomographic perfusion. *Stroke*. Sep 2017;48(9):2593-2596. PMID 28716981.
19. van Seeters T, Biessels GJ, Kappelle LJ, et al. The prognostic value of CT angiography and CT perfusion in acute ischemic stroke. *Cerebrovasc Dis*. Oct 2015;40(5-6):258-269. PMID 26484857.
20. Greenberg ED, Gold R, Reichman M, et al. Diagnostic accuracy of CT angiography and CT perfusion for cerebral vasospasm: a meta-analysis. *AJNR Am J Neuroradiol*. Nov 2010;31(10):1853-1860. PMID 20884748.
21. Stecco A, Fabbiano F, Amatuzzo P et al. Computed tomography perfusion and computed tomography angiography in vasospasm after subarachnoid hemorrhage. *J Neurosurg Sci*, 2016 Apr 29;62(4). PMID 27124176.
22. Sanelli PC, Jou A, Gold R, et al. Using CT perfusion during the early baseline period in aneurysmal subarachnoid hemorrhage to assess for development of vasospasm. *Neuroradiology*. Jun 2011;53(6):425-434. PMID 20694461.
23. Cremers CH, van der Schaaf IC, Wensink E, et al. CT perfusion and delayed cerebral ischemia in aneurysmal subarachnoid hemorrhage: a systematic review and meta-analysis. *J Cereb Blood Flow Metab*. Feb 2014;34(2):200-207. PMID 24281744.
24. Sanelli PC, Ugorec I, Johnson CE, et al. Using quantitative CT perfusion for evaluation of delayed cerebral ischemia following aneurysmal subarachnoid hemorrhage. *AJNR Am J Neuroradiol*. Dec 2011;32(11):2047- 2053. PMID 21960495.
25. Xyda A, Haberland U, Klotz E, et al. Brain volume perfusion CT performed with 128-detector row CT system in patients with cerebral gliomas: a feasibility study. *Eur Radiol*. Sep 2011;21(9):1811-1819. PMID 21573969.
26. Jain R. Perfusion CT imaging of brain tumors: an overview. *AJNR Am J Neuroradiol*. Oct 2011;32(9):1570-1577. PMID 21051510.

27. Ellika SK, Jain R, Patel SC, et al. Role of perfusion CT in glioma grading and comparison with conventional MR imaging features. *AJNR Am J Neuroradiol*. Nov-Dec 2007;28(10):1981-1987. PMID 17893216.
28. Jain R, Ellika SK, Scarpace L, et al. Quantitative estimation of permeability surface-area product in astroglial brain tumors using perfusion CT and correlation with histopathologic grade. *AJNR Am J Neuroradiol*. Apr 2008;29(4):694-700. PMID 18202239.
29. Connolly ES, Jr., Rabinstein AA, Carhuapoma JR, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. Jun 2012;43(6):1711-1737. PMID 22556195.
30. Jauch EC, Saver JL, Adams HP, Jr., et al. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. Mar 2013;44(3):870-947. PMID 23370205.
31. Powers WJ, Rabinstein AA, Ackerson T, et al. 2018 guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. Mar 2018;49(3):e46-e110. PMID 29367334.
32. Wintermark M, Sanelli PC, Albers GW, et al. Imaging recommendations for acute stroke and transient ischemic attack patients: A joint statement by the American Society of Neuroradiology, the American College of Radiology, and the Society of NeuroInterventional Surgery. *AJNR Am J Neuroradiol*. Nov-Dec 2013;34(11):E117-127. PMID 23907247.
33. American College of Radiology, American Society of Neuroradiology, Society for Pediatric Radiology. ACR-ASNR-SPR practice parameter for the performance of computed tomography (CT) perfusion in neuroradiologic imaging. 2017; <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/ct-perfusion.pdf?la=en>. Accessed August 5, 2019.
34. American College of Radiology (ACR). ACR Appropriateness Criteria: Cerebrovascular Disease. 2016; <https://acsearch.acr.org/docs/69478/Narrative/>. Accessed August 5, 2019.