



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Spinal Cord and Dorsal Root Ganglion Stimulation

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 472

BCBSA Reference Number: 7.01.25 (For Plan internal use only)

Related Policies

Deep Brain Stimulation, [#473](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Spinal cord stimulation with standard or high-frequency stimulation may be considered **MEDICALLY NECESSARY** for treatment of severe and chronic pain of the trunk or limbs that is refractory to all other pain therapies when ALL of the following patient criteria are met:

- The treatment is used only as a last resort; other treatment modalities (pharmacologic, surgical, psychological, physical, if applicable) have failed or are judged to be unsuitable or contraindicated;
- Pain is neuropathic in nature (ie, resulting from actual damage to the peripheral nerves). Common indications include, but are not limited to, failed back surgery syndrome, complex regional pain syndrome (ie, reflex sympathetic dystrophy), arachnoiditis, radiculopathies, phantom limb/stump pain, and peripheral neuropathy (including diabetic peripheral neuropathy). Spinal cord stimulation is generally not effective in treating nociceptive pain (resulting from irritation, not damage to the nerves) and central deafferentation pain (related to central nervous system damage from a stroke or spinal cord injury).
- No serious untreated drug habituation exists;
- Demonstration of at least 50% pain relief with a temporarily implanted electrode precedes permanent implantation;
- All the facilities, equipment, and professional and support personnel required for the proper diagnosis, treatment, and follow-up of the patient are available.

Dorsal root ganglion neurostimulation is considered **MEDICALLY NECESSARY** for the treatment of severe and chronic pain of the trunk or limbs that is refractory to all other pain therapies when ALL of the following patient criteria are met:

- The treatment is used only as a last resort; other treatment modalities (pharmacologic, surgical, psychological, physical, if applicable) have failed or are judged to be unsuitable or contraindicated;
- Pain is neuropathic in nature (ie, resulting from actual damage to the peripheral nerves). Common indications include, but are not limited to, failed back surgery syndrome, complex regional pain syndrome (ie, reflex sympathetic dystrophy), arachnoiditis, radiculopathies, phantom limb/stump pain,

and peripheral neuropathy (including diabetic peripheral neuropathy). Spinal cord stimulation is generally not effective in treating nociceptive pain (resulting from irritation, not damage to the nerves) and central deafferentation pain (related to central nervous system damage from a stroke or spinal cord injury).

- No serious untreated drug habituation exists;
- Demonstration of at least 50% pain relief with a temporarily implanted electrode precedes permanent implantation;
- All the facilities, equipment, and professional and support personnel required for the proper diagnosis, treatment, and follow-up of the patient are available.

Spinal cord stimulation is considered **INVESTIGATIONAL** in all other situations including, but not limited to, treatment of critical limb ischemia to forestall amputation and treatment of refractory angina pectoris, heart failure, and cancer-related pain.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is required for 63655 & 63685
Commercial PPO	Prior authorization is required for 63655 & 63685

Note: For Commercial Managed Care (HMO and POS) and Medicare HMO BlueSM, the temporary implanted electrode (CPT code 63650) that precedes permanent implantation does not require prior authorization. However, prior authorization is required for the permanent placement of implanted electrodes (CPT codes 63655 and 63685).

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:

CPT Codes

CPT codes:	Code Description
63655	Laminectomy for implantation of neurostimulator electrode plate/paddle; epidural
63685	Insertion or replacement of spinal neurostimulator pulse generator or receiver, direct or inductive coupling

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:

CPT Codes

CPT codes:	Code Description
63650	Percutaneous implantation of neurostimulator electrode array; epidural

HCPCS Codes

HCPCS codes:	Code Description
C1822	Generator, neurostimulator (implantable), high frequency, with rechargeable battery and charging system

The following ICD Diagnosis Codes are considered medically necessary when submitted with the CPT and HCPCS codes above if medical necessity criteria are met:

ICD-10 Diagnosis Codes

ICD-10-CM Diagnosis codes:	Code Description
G03.9	Meningitis, unspecified
G56.40	Causalgia of unspecified upper limb
G56.41	Causalgia of right upper limb
G56.42	Causalgia of left upper limb
G57.70	Causalgia of unspecified lower limb
G57.71	Causalgia of right lower limb
G57.72	Causalgia of left lower limb
G60.3	Idiopathic progressive neuropathy
G60.8	Other hereditary and idiopathic neuropathies
G60.9	Hereditary and idiopathic neuropathy, unspecified
G62.89	Other specified polyneuropathies
G62.9	Polyneuropathy, unspecified
G89.21	Chronic pain due to trauma
G89.22	Chronic post-thoracotomy pain
G89.28	Other chronic postprocedural pain
G89.29	Other chronic pain
G89.3	Neoplasm related pain (acute) (chronic)
G89.4	Chronic pain syndrome
G90.511	Complex regional pain syndrome I of right upper limb
G90.512	Complex regional pain syndrome I of left upper limb
G90.513	Complex regional pain syndrome I of upper limb, bilateral
G90.519	Complex regional pain syndrome I of unspecified upper limb
G90.521	Complex regional pain syndrome I of right lower limb
G90.522	Complex regional pain syndrome I of left lower limb
G90.523	Complex regional pain syndrome I of lower limb, bilateral
G90.529	Complex regional pain syndrome I of unspecified lower limb
G90.59	Complex regional pain syndrome I of other specified site
M54.10	Radiculopathy, Site Unspecified
M54.11	Radiculopathy, occipito-atlanto-axial region
M54.12	Radiculopathy, cervical region
M54.13	Radiculopathy, cervicothoracic region
M54.14	Radiculopathy, thoracic region
M54.15	Radiculopathy, thoracolumbar region
M54.16	Radiculopathy, lumbar region
M54.17	Radiculopathy, lumbosacral region

M54.18	Radiculopathy, sacral and sacrococcygeal region
M79.2	Neuralgia And Neuritis, Unspecified
M96.1	Postlaminectomy syndrome, not elsewhere classified

Description

Chronic Pain

Spinal cord stimulation has been used in a wide variety of chronic refractory pain conditions, including pain associated with cancer, failed back pain syndromes, arachnoiditis, and complex regional pain syndrome (CPRS; ie, chronic reflex sympathetic dystrophy). There has also been interest in spinal cord stimulation as a treatment of critical limb ischemia, primarily in patients who are poor candidates for revascularization and in patients with refractory chest pain.

Spinal Cord Stimulation

Spinal cord stimulation (also called dorsal column stimulation) involves the use of low-level epidural electrical stimulation of the spinal cord dorsal columns. The neurophysiology of pain relief after spinal cord stimulation is uncertain, but may be related to either activation of an inhibitory system or blockage of facilitative circuits.

Spinal cord stimulation devices consist of several components: (1) the lead that delivers the electrical stimulation to the spinal cord; (2) an extension wire that conducts the electrical stimulation from the power source to the lead; and (3) a power source that generates the electricity. The lead may incorporate from 4 to 8 electrodes, with 8 electrodes more commonly used for complex pain patterns. There are 2 basic types of power source: 1 type, the power source (battery), can be surgically implanted or worn externally with an antenna over the receiver; the other, a radiofrequency receiver, is implanted. Totally implantable systems are most commonly used.

The patient's pain distribution pattern dictates at what level of the spinal cord the stimulation lead is placed. The pain pattern may influence the type of device used. For example, a lead with 8 electrodes may be selected for those with complex pain patterns or bilateral pain. Implantation of the spinal cord stimulator is typically a 2-step process. Initially, the electrode is temporarily implanted in the epidural space, allowing a trial period of stimulation. Once treatment effectiveness is confirmed (defined as at least 50% reduction in pain), the electrodes and radio-receiver/transducer are permanently implanted. Successful spinal cord stimulation may require extensive programming of the neurostimulators to identify the optimal electrode combinations and stimulation channels.

Traditional spinal cord stimulation devices use electrical stimulation with a frequency of 100 to 1000 Hz. In 2015, a spinal cord stimulation device, using a higher frequency (10,000 Hz) than predicate devices, was approved by the U.S. Food and Drug Administration (FDA) through the premarket approval process. High-frequency stimulation is proposed to be associated with fewer paresthesias, which are a recognized effect of spinal cord stimulation. In 2016, the FDA approved a clinician programmer application that allows a spinal cord stimulation device to provide stimulation in bursts rather than at a constant rate. Burst stimulation is proposed to relieve pain with fewer paresthesias. The burst stimulation device works in conjunction with standard spinal cord stimulation devices. With the newly approved app, stimulation is provided in five, 500-Hz burst spikes at a rate of 40 Hz, with a pulse width of 1 ms.

The incidence of adverse events related to spinal cord stimulation has been reported to occur in 30% to 40% of cases.¹ Adverse events can either be hardware-related or biological. Hardware-related complications include lead migration or failure or fracture. Biological complications include infection and pain. More severe biological complications are rare, including dural puncture headache (estimated incidence, up to 0.3%) and neurological damage (estimated incidence, 0.25%).

Other neurostimulators target the dorsal root ganglion. Dorsal root ganglia consist of sensory cell bodies that transmit input from the peripheral nervous system to the central nervous system and play a role in neuropathic pain perception. Dorsal root ganglia are located in the epidural space between spinal nerves

and the spinal cord on the posterior root in a minimal amount of cerebrospinal fluid, amenable to epidural access. Two systems targeting the dorsal root ganglion have received approval or clearance from the FDA.

A retrospective analysis of the FDA's Manufacturer and User Facility Device Experience (MAUDE) database provided information on complications related to the use of dorsal root ganglion stimulation.² The MAUDE database was queried for dorsal root ganglion stimulation reports through 2017, identifying 979 episodes. Complications were predominantly device-related (47%; lead migration and lead damage), with the remaining comprised of procedural complications (28%; infection, new neurologic symptoms, and dural puncture), patient complaints (12%; site pain and unwanted stimulation), serious adverse events (2.4%), and "other" complications (4.6%). The prevalence of complications cannot be estimated using the MAUDE database; while facilities are mandated to report events, patients and health care providers may report events, but are not mandated to do so.

In September 2020, the FDA released a letter to healthcare providers reminding them to conduct a trial stimulation period before implanting a spinal cord stimulator as the agency continues to receive reports of serious adverse effects associated with these devices.³ Between July 27, 2016 and July 27, 2020, the FDA received 107,728 medical device reports related to spinal cord stimulators intended for pain including 497 associated with patient death, 77,937 with patient injury, and 29,924 with device malfunction. The most frequently reported patient problem codes were inadequate pain relief (28.1%), pain (15.2%), unexpected therapeutic effects (10.9%), infection (7.5%), and discomfort (5.9%). Additionally, the most frequently reported device problem codes were charging problems (11.2%), impedance (10.6%), migration (7.2%), battery problem (6.4%), and premature discharge of battery (4.2%). The FDA made the following recommendations for clinicians to consider:

- Conduct a trial stimulation as described in the device labeling to identify and confirm satisfactory pain relief before permanent implantation.
- Permanent spinal cord stimulation should only be implanted in patients who have undergone and passed a stimulation trial.
- Providers typically perform a stimulation trial on a patient for 3 to 7 days, and success is usually defined by a 50% reduction in pain symptoms. Inform patients about the risks of serious side effects and what to expect during the trial stimulation.
- Before implantation of any spinal cord stimulation, discuss the benefits and risks of the different types of implants and other treatment options, including magnetic resonance imaging (MRI) compatibility of the devices.
- Before implantation, provide patients with the manufacturer's patient labeling and any other education materials for the device that will be implanted.
- Develop an individualized programming, treatment, and follow-up plan for spinal cord stimulation therapy delivery with each patient.
- Provide each patient with the name of the device manufacturer, model, and the unique device identifier of the implant received.

Summary

Spinal cord stimulation delivers low-voltage electrical stimulation to the dorsal columns of the spinal cord to block the sensation of pain; this is achieved through a surgically implanted spinal cord stimulation device, which comes equipped with a radiofrequency receiver. The neurostimulator device is also issued with a standard power source (battery) that can be implanted or worn externally. Other neurostimulators target the dorsal root ganglion.

Treatment-Refractory Chronic Pain

For individuals who have treatment-refractory chronic pain of the trunk or limbs who receive standard spinal cord stimulation, the evidence includes systematic reviews and randomized controlled trials (RCTs). Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. Available RCTs are heterogeneous regarding underlying diagnoses in select patient populations. However, the trials including patients with underlying neuropathic pain processes

have shown a significant benefit with spinal cord stimulation. Systematic reviews have supported the use of spinal cord stimulation to treat refractory trunk or limb pain, and patients who have failed all other treatment modalities have few options. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have treatment-refractory chronic pain of the trunk or limbs who receive high-frequency spinal cord stimulation, the evidence includes a systematic review and 4 RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. Two RCTs that enrolled participants not previously treated with spinal cord stimulation reported clinically and statistically significant benefits associated with high-frequency spinal cord stimulation. Another RCT in patients who had chronic pain despite previous treatment with standard spinal cord stimulation found no benefit for those receiving high-frequency stimulation compared with sham-control; however, it is difficult to compare these findings with other trials of spinal cord stimulation due to the different patient populations, short treatment periods, and the crossover period effect. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have treatment-refractory chronic pain of the trunk or limbs who receive dorsal root ganglion neurostimulation, the evidence includes a systematic review, an RCT, and case series. Relevant outcomes are symptoms, functional outcomes, quality of life, medication use, and treatment-related morbidity. The unblinded RCT found that patients receiving dorsal root ganglion neurostimulation had significantly higher rates of treatment success (physical functioning score and quality of life measures), at 3 and 12 months compared with those receiving standard spinal cord stimulation devices. Dorsal root ganglion neurostimulation was found to be noninferior to spinal cord stimulation in the percentage achieving $\geq 50\%$ pain reduction, emotional functioning score, and 36-Item Short-Form Health Survey scores. Both groups experienced paresthesias but patients in the dorsal root ganglion group reported less postural variation in paresthesia and reduced extraneous stimulation in nonpainful areas. Rates of serious adverse events were similar between the 2 study arms. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Critical Limb Ischemia

For individuals who have critical limb ischemia who receive spinal cord stimulation, the evidence includes systematic reviews of several small RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. In pooled analyses, spinal cord stimulation was associated with a lower risk of amputation versus control, but results were not consistently statistically significant due to differences in methodologies. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Treatment-Refractory Angina Pectoris

For individuals who have treatment-refractory angina pectoris who receive spinal cord stimulation, the evidence includes systematic reviews and RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. Numerous small RCTs have evaluated spinal cord stimulation as a treatment for refractory angina. While some have reported benefits, most have not. In 2 recent RCTs, there was no significant benefit in the primary outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Heart Failure

For individuals who have heart failure who receive spinal cord stimulation, the evidence includes RCTs. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. An RCT (N=66) comparing spinal cord stimulation using active stimulation with sham-control in patients who had New York Heart Association functional class III heart failure and a left ventricular ejection fraction of 35% or less did not find significant differences between groups, but might have been underpowered to do so. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Cancer-Related Pain

For individuals who have cancer-related pain who receive spinal cord stimulation, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, medication use, and treatment-related morbidity. No RCTs evaluating spinal cord stimulation in this population were identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
10/2022	Policy clarified. Peripheral neuropathy updated to include diabetic peripheral neuropathy.
6/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2022	Prior authorization information clarified for PPO plans. Effective 6/1/2022.
5/2021	BCBSA National medical policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
6/2020	BCBSA National medical policy review. Description, summary and references updated. Policy statements unchanged.
9/2019	BCBSA National medical policy review. New medically necessary indications described. Dorsal root ganglion neurostimulation is considered medically necessary for the treatment of severe and chronic pain of the trunk or limbs. Clarified coding information. Effective 9/1/2019.
1/2019	Clarified coding information.
6/2018	BCBSA National medical policy review. Policy clarified to include burst neurostimulation as an alternate programming of a standard SCS device.
3/2018	Clarified coding information.
12/2017	BCBSA National medical policy review. "Wireless injectable" removed from policy statement on dorsal root ganglion neurostimulation. "Dorsal root ganglion" added to policy title. Effective 12/1/2017.
6/2017	BCBSA National medical policy review. New medically necessary and investigational indications described. Effective 6/1/2017.
9/2016	BCBSA National medical policy review. New investigational indications described. Clarified coding information. Effective 9/1/2016
6/2016	Clarified coding information.
1/2016	Clarified coding information.
6/2015	BCBSA National medical policy review. New investigational indications described. Effective 6/1/2015.
2/2015	Prior authorization clarified for CPT codes 63650 and 63685. Clarified coding information.
6/2014	Updated Coding section with ICD10 procedure and diagnosis codes, effective 10/2015.
6/2014	BCBSA National medical policy review. New investigational indications described; existing investigational statement modified to state all other situations. Effective 6/1/2014.
12/2013	Removed HCPCS codes C1767, C1816, L8680, and L8682-L8688 as they do not meet the intent of the policy.
10/2013	Removed CPT codes 63661, 63662, 63663, 63664, 63688 and diagnosis codes 337.20 and 722.80 as they do not apply to the policy.
2/2013	New references from BCBSA National medical policy.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
4/2011	BCBSA National medical policy review. No changes to policy statements.

1/2011	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
4/2010	BCBSA National medical policy review. Changes to policy statements.
1/2010	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
1/2009	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
11/2008	Policy 083 effective 11/2008 stating covered and non-covered indications.
1/2008	BCBSA National medical policy review. No changes to policy statements.
1/2008	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
7/2007	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.
4/2007	BCBSA National medical policy review. No changes to policy statements.
2/2007	Reviewed - Medical Policy Group - Psychiatry and Ophthalmology. No changes to policy statements.
1/2007	Reviewed - Medical Policy Group - Neurology and Neurosurgery. No changes to policy statements.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Eldabe S, Buchser E, Duarte RV. Complications of Spinal Cord Stimulation and Peripheral Nerve Stimulation Techniques: A Review of the Literature. *Pain Med.* Feb 2016; 17(2): 325-36. PMID 26814260
2. Sivanesan E, Bicket MC, Cohen SP. Retrospective analysis of complications associated with dorsal root ganglion stimulation for pain relief in the FDA MAUDE database. *Reg Anesth Pain Med.* Jan 2019; 44(1): 100-106. PMID 30640660
3. U.S. Food and Drug Administration. Conduct a trial stimulation period before implanting a spinal cord stimulator (SCS) - letter to health care providers. September 3, 2020. <https://www.fda.gov/medical-devices/letters-health-care-providers/conduct-trial-stimulation-period-implanting-spinal-cord-stimulator-scs-letter-health-care-providers>. Accessed March 3, 2022.
4. Food and Drug Administration. Cordis Programmable Neural Stimulator: Premarket Approval. Accessed February 28, 2022.
5. Turk DC, Dworkin RH, Allen RR, et al. Core outcome domains for chronic pain clinical trials: IMMPACT recommendations. *Pain.* Dec 2003; 106(3): 337-345. PMID 14659516
6. Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain.* Jan 2005; 113(1-2): 9-19. PMID 15621359
7. Dworkin RH, Turk DC, Wyrwich KW, et al. Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain.* Feb 2008; 9(2): 105-21. PMID 18055266
8. Kerns RD, Turk DC, Rudy TE. The West Haven-Yale Multidimensional Pain Inventory (WHYMPI). *Pain.* Dec 1985; 23(4): 345-356. PMID 4088697
9. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singap.* Mar 1994; 23(2): 129-38. PMID 8080219
10. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine (Phila Pa 1976).* Nov 15 2000; 25(22): 2940-52; discussion 2952. PMID 11074683
11. Ostelo RW, Deyo RA, Stratford P, et al. Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine (Phila Pa 1976).* Jan 01 2008; 33(1): 90-4. PMID 18165753

12. Wells GA, Tugwell P, Kraag GR, et al. Minimum important difference between patients with rheumatoid arthritis: the patient's perspective. *J Rheumatol*. Mar 1993; 20(3): 557-60. PMID 8478873
13. Kosinski M, Zhao SZ, Dedhiya S, et al. Determining minimally important changes in generic and disease-specific health-related quality of life questionnaires in clinical trials of rheumatoid arthritis. *Arthritis Rheum*. Jul 2000; 43(7): 1478-87. PMID 10902749
14. Angst F, Aeschlimann A, Stucki G. Smallest detectable and minimal clinically important differences of rehabilitation intervention with their implications for required sample sizes using WOMAC and SF-36 quality of life measurement instruments in patients with osteoarthritis of the lower extremities. *Arthritis Rheum*. Aug 2001; 45(4): 384-91. PMID 11501727
15. Beck ATS, R.A. Beck Depression Inventory. San Antonio, TX: Psychological Corporation; 1993.
16. Curran SL, Andrykowski MA, Studts JL. Short Form of the Profile of Mood States (POMS-SF): Psychometric information. *Psychol Assess* 1995;7:80-83.
17. Visnjevac O, Costandi S, Patel BA, et al. A Comprehensive Outcome-Specific Review of the Use of Spinal Cord Stimulation for Complex Regional Pain Syndrome. *Pain Pract*. Apr 2017; 17(4): 533-545. PMID 27739179
18. O'Connell NE, Wand BM, McAuley J, et al. Interventions for treating pain and disability in adults with complex regional pain syndrome. *Cochrane Database Syst Rev*. Apr 30 2013; (4): CD009416. PMID 23633371
19. Grider JS, Manchikanti L, Carayannopoulos A, et al. Effectiveness of Spinal Cord Stimulation in Chronic Spinal Pain: A Systematic Review. *Pain Physician*. Jan 2016; 19(1): E33-54. PMID 26752493
20. Head J, Mazza J, Sabourin V, et al. Waves of Pain Relief: A Systematic Review of Clinical Trials in Spinal Cord Stimulation Waveforms for the Treatment of Chronic Neuropathic Low Back and Leg Pain. *World Neurosurg*. Nov 2019; 131: 264-274.e3. PMID 31369885
21. Duarte RV, Nevitt S, Maden M, et al. Spinal cord stimulation for the management of painful diabetic neuropathy: a systematic review and meta-analysis of individual patient and aggregate data. *Pain*. Nov 01 2021; 162(11): 2635-2643. PMID 33872236
22. Henson JV, Varhabhatla NC, Bebic Z, et al. Spinal Cord Stimulation for Painful Diabetic Peripheral Neuropathy: A Systematic Review. *Pain Ther*. Dec 2021; 10(2): 895-908. PMID 34244979
23. Raghu ALB, Parker T, Aziz TZ, et al. Invasive Electrical Neuromodulation for the Treatment of Painful Diabetic Neuropathy: Systematic Review and Meta-Analysis. *Neuromodulation*. Jan 2021; 24(1): 13-21. PMID 32588933
24. Strand NH, Burkey AR. Neuromodulation in the Treatment of Painful Diabetic Neuropathy: A Review of Evidence for Spinal Cord Stimulation. *J Diabetes Sci Technol*. Mar 2022; 16(2): 332-340. PMID 34842478
25. O'Connell NE, Ferraro MC, Gibson W, et al. Implanted spinal neuromodulation interventions for chronic pain in adults. *Cochrane Database Syst Rev*. Dec 02 2021; 12: CD013756. PMID 34854473
26. North RB, Kidd DH, Farrokhi F, et al. Spinal cord stimulation versus repeated lumbosacral spine surgery for chronic pain: a randomized, controlled trial. *Neurosurgery*. 2005; 56(1): 98-106; discussion 106-7. PMID 15617591
27. Kumar K, Taylor RS, Jacques L, et al. Spinal cord stimulation versus conventional medical management for neuropathic pain: a multicentre randomised controlled trial in patients with failed back surgery syndrome. *Pain*. Nov 2007; 132(1-2): 179-88. PMID 17845835
28. Kumar K, Taylor RS, Jacques L, et al. The effects of spinal cord stimulation in neuropathic pain are sustained: a 24-month follow-up of the prospective randomized controlled multicenter trial of the effectiveness of spinal cord stimulation. *Neurosurgery*. Oct 2008; 63(4): 762-70; discussion 770. PMID 18981888
29. Kemler MA, Barendse GA, van Kleef M, et al. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. *N Engl J Med*. Aug 31 2000; 343(9): 618-24. PMID 10965008
30. Kemler MA, De Vet HC, Barendse GA, et al. The effect of spinal cord stimulation in patients with chronic reflex sympathetic dystrophy: two years' follow-up of the randomized controlled trial. *Ann Neurol*. Jan 2004; 55(1): 13-8. PMID 14705107
31. Kemler MA, de Vet HC, Barendse GA, et al. Effect of spinal cord stimulation for chronic complex regional pain syndrome Type I: five-year final follow-up of patients in a randomized controlled trial. *J Neurosurg*. Feb 2008; 108(2): 292-8. PMID 18240925
32. Slangen R, Schaper NC, Faber CG, et al. Spinal cord stimulation and pain relief in painful diabetic peripheral neuropathy: a prospective two-center randomized controlled trial. *Diabetes Care*. Nov 2014; 37(11): 3016-24. PMID 25216508

33. de Vos CC, Meier K, Zaalberg PB, et al. Spinal cord stimulation in patients with painful diabetic neuropathy: a multicentre randomized clinical trial. *Pain*. Nov 2014; 155(11): 2426-31. PMID 25180016
34. Duarte RV, Andronis L, Lenders MW, et al. Quality of life increases in patients with painful diabetic neuropathy following treatment with spinal cord stimulation. *Qual Life Res*. Jul 2016; 25(7): 1771-7. PMID 26694963
35. Rigoard P, Basu S, Desai M, et al. Multicolumn spinal cord stimulation for predominant back pain in failed back surgery syndrome patients: a multicenter randomized controlled trial. *Pain*. Jun 2019; 160(6): 1410-1420. PMID 30720582
36. Hou S, Kemp K, Grabois M. A Systematic Evaluation of Burst Spinal Cord Stimulation for Chronic Back and Limb Pain. *Neuromodulation*. Jun 2016; 19(4): 398-405. PMID 27139915
37. De Ridder D, Plazier M, Kamerling N, et al. Burst spinal cord stimulation for limb and back pain. *World Neurosurg*. Nov 2013; 80(5): 642-649.e1. PMID 23321375
38. De Ridder D, Vanneste S, Plazier M, et al. Burst spinal cord stimulation: toward paresthesia-free pain suppression. *Neurosurgery*. May 2010; 66(5): 986-90. PMID 20404705
39. Schu S, Sloty PJ, Bara G, et al. A prospective, randomised, double-blind, placebo-controlled study to examine the effectiveness of burst spinal cord stimulation patterns for the treatment of failed back surgery syndrome. *Neuromodulation*. Jul 2014; 17(5): 443-50. PMID 24945621
40. Kriek N, Groeneweg JG, Stronks DL, et al. Preferred frequencies and waveforms for spinal cord stimulation in patients with complex regional pain syndrome: A multicentre, double-blind, randomized and placebo-controlled crossover trial. *Eur J Pain*. Mar 2017; 21(3): 507-519. PMID 27714945
41. Deer T, Slavin KV, Amirdelfan K, et al. Success Using Neuromodulation With BURST (SUNBURST) Study: Results From a Prospective, Randomized Controlled Trial Using a Novel Burst Waveform. *Neuromodulation*. Jan 2018; 21(1): 56-66. PMID 28961366
42. Eldabe S, Duarte R, Gulve A, et al. Analgesic Efficacy of "Burst" and Tonic (500 Hz) Spinal Cord Stimulation Patterns: A Randomized Placebo-Controlled Crossover Study. *Neuromodulation*. Apr 2021; 24(3): 471-478. PMID 33251662
43. Bicket MC, Dunn RY, Ahmed SU. High-Frequency Spinal Cord Stimulation for Chronic Pain: Pre-Clinical Overview and Systematic Review of Controlled Trials. *Pain Med*. Dec 2016; 17(12): 2326-2336. PMID 28025366
44. Perruchoud C, Eldabe S, Batterham AM, et al. Analgesic efficacy of high-frequency spinal cord stimulation: a randomized double-blind placebo-controlled study. *Neuromodulation*. Jul-Aug 2013; 16(4): 363-9; discussion 369. PMID 23425338
45. Kapural L, Yu C, Doust MW, et al. Novel 10-kHz High-frequency Therapy (HF10 Therapy) Is Superior to Traditional Low-frequency Spinal Cord Stimulation for the Treatment of Chronic Back and Leg Pain: The SENZA-RCT Randomized Controlled Trial. *Anesthesiology*. Oct 2015; 123(4): 851-60. PMID 26218762
46. Petersen EA, Stauss TG, Scowcroft JA, et al. Effect of High-frequency (10-kHz) Spinal Cord Stimulation in Patients With Painful Diabetic Neuropathy: A Randomized Clinical Trial. *JAMA Neurol*. Jun 01 2021; 78(6): 687-698. PMID 33818600
47. De Andres J, Monsalve-Dolz V, Fabregat-Cid G, et al. Prospective, Randomized Blind Effect-on-Outcome Study of Conventional vs High-Frequency Spinal Cord Stimulation in Patients with Pain and Disability Due to Failed Back Surgery Syndrome. *Pain Med*. Dec 01 2017; 18(12): 2401-2421. PMID 29126228
48. Petersen EA, Stauss TG, Scowcroft JA, et al. Durability of High-Frequency 10-kHz Spinal Cord Stimulation for Patients With Painful Diabetic Neuropathy Refractory to Conventional Treatments: 12-Month Results From a Randomized Controlled Trial. *Diabetes Care*. Jan 01 2022; 45(1): e3-e6. PMID 34844993
49. Kapural L, Yu C, Doust MW, et al. Comparison of 10-kHz High-Frequency and Traditional Low-Frequency Spinal Cord Stimulation for the Treatment of Chronic Back and Leg Pain: 24-Month Results From a Multicenter, Randomized, Controlled Pivotal Trial. *Neurosurgery*. Nov 2016; 79(5): 667-677. PMID 27584814
50. Al-Kaisy A, Palmisani S, Smith TE, et al. Long-Term Improvements in Chronic Axial Low Back Pain Patients Without Previous Spinal Surgery: A Cohort Analysis of 10-kHz High-Frequency Spinal Cord Stimulation over 36 Months. *Pain Med*. Jun 01 2018; 19(6): 1219-1226. PMID 29077889

51. Vuka I, Marcius T, Dosenovic S, et al. Neuromodulation with electrical field stimulation of dorsal root ganglion in various pain syndromes: a systematic review with focus on participant selection. *J Pain Res.* 2019; 12: 803-830. PMID 30881093
52. Deer TR, Hunter CW, Mehta P, et al. A Systematic Literature Review of Dorsal Root Ganglion Neurostimulation for the Treatment of Pain. *Pain Med.* Aug 01 2020; 21(8): 1581-1589. PMID 32803221
53. Moman RN, Peterson AA, Maher DP, et al. Infectious Complications of Dorsal Root Ganglion Stimulation: A Systematic Review and Pooled Analysis of Incidence. *Neuromodulation.* Jun 06 2021. PMID 34096135
54. Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): Axium Neurostimulator System. 2016; https://www.accessdata.fda.gov/cdrh_docs/pdf15/P150004b.pdf. Accessed March 3, 2022.
55. Deer TR, Levy RM, Kramer J, et al. Dorsal root ganglion stimulation yielded higher treatment success rate for complex regional pain syndrome and causalgia at 3 and 12 months: a randomized comparative trial. *Pain.* Apr 2017; 158(4): 669-681. PMID 28030470
56. Mekhail N, Deer TR, Kramer J, et al. Paresthesia-Free Dorsal Root Ganglion Stimulation: An ACCURATE Study Sub-Analysis. *Neuromodulation.* Feb 2020; 23(2): 185-195. PMID 30861286
57. Weiner RL, Yeung A, Montes Garcia C, et al. Treatment of FBSS Low Back Pain with a Novel Percutaneous DRG Wireless Stimulator: Pilot and Feasibility Study. *Pain Med.* Oct 2016; 17(10): 1911-1916. PMID 27125284
58. Ubbink DT, Vermeulen H. Spinal cord stimulation for non-reconstructable chronic critical leg ischaemia. *Cochrane Database Syst Rev.* Feb 28 2013; (2): CD004001. PMID 23450547
59. Klomp HM, Steyerberg EW, Habbema JD, et al. What is the evidence on efficacy of spinal cord stimulation in (subgroups of) patients with critical limb ischemia?. *Ann Vasc Surg.* May-Jun 2009; 23(3): 355-63. PMID 19128928
60. Klomp HM, Spincemaille GH, Steyerberg EW, et al. Spinal-cord stimulation in critical limb ischaemia: a randomised trial. ESES Study Group. *Lancet.* Mar 27 1999; 353(9158): 1040-4. PMID 10199350
61. Abu Dabrh AM, Steffen MW, Asi N, et al. Nonrevascularization-based treatments in patients with severe or critical limb ischemia. *J Vasc Surg.* Nov 2015; 62(5): 1330-9.e13. PMID 26409842
62. Pan X, Bao H, Si Y, et al. Spinal Cord Stimulation for Refractory Angina Pectoris: A Systematic Review and Meta-analysis. *Clin J Pain.* Jun 2017; 33(6): 543-551. PMID 27875377
63. Tsigaridas N, Naka K, Tsapogas P, et al. Spinal cord stimulation in refractory angina. A systematic review of randomized controlled trials. *Acta Cardiol.* Apr 2015; 70(2): 233-43. PMID 26148385
64. Zipes DP, Svorkdal N, Berman D, et al. Spinal cord stimulation therapy for patients with refractory angina who are not candidates for revascularization. *Neuromodulation.* Nov-Dec 2012; 15(6): 550-8; discussion 558-9. PMID 22494013
65. Lanza GA, Grimaldi R, Greco S, et al. Spinal cord stimulation for the treatment of refractory angina pectoris: a multicenter randomized single-blind study (the SCS-ITA trial). *Pain.* Jan 2011; 152(1): 45-52. PMID 21084162
66. Torre-Amione G, Alo K, Estep JD, et al. Spinal cord stimulation is safe and feasible in patients with advanced heart failure: early clinical experience. *Eur J Heart Fail.* Jul 2014; 16(7): 788-95. PMID 24961194
67. Zipes DP, Neuzil P, Theres H, et al. Determining the Feasibility of Spinal Cord Neuromodulation for the Treatment of Chronic Systolic Heart Failure: The DEFEAT-HF Study. *JACC Heart Fail.* Feb 2016; 4(2): 129-136. PMID 26682789
68. Lihua P, Su M, Zejun Z, et al. Spinal cord stimulation for cancer-related pain in adults. *Cochrane Database Syst Rev.* Feb 28 2013; (2): CD009389. PMID 23450600
69. Peng L, Min S, Zejun Z, et al. Spinal cord stimulation for cancer-related pain in adults. *Cochrane Database Syst Rev.* Jun 29 2015; (6): CD009389. PMID 26121600
70. Mekhail NA, Mathews M, Nageeb F, et al. Retrospective review of 707 cases of spinal cord stimulation: indications and complications. *Pain Pract.* Mar-Apr 2011; 11(2): 148-53. PMID 21371254
71. Lanza GA, Barone L, Di Monaco A. Effect of spinal cord stimulation in patients with refractory angina: evidence from observational studies. *Neuromodulation.* Nov-Dec 2012; 15(6): 542-9; discussion 549. PMID 22364309
72. Deer T, Pope J, Hunter C, et al. Safety Analysis of Dorsal Root Ganglion Stimulation in the Treatment of Chronic Pain. *Neuromodulation.* Feb 2020; 23(2): 239-244. PMID 30861617

73. Dworkin RH, O'Connor AB, Kent J, et al. Interventional management of neuropathic pain: NeuPSIG recommendations. *Pain*. Nov 2013; 154(11): 2249-2261. PMID 23748119
74. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. Apr 2013; 16(2 Suppl): S49-283. PMID 23615883
75. Aman MM, Mahmoud A, Deer T, et al. The American Society of Pain and Neuroscience (ASPN) Best Practices and Guidelines for the Interventional Management of Cancer-Associated Pain. *J Pain Res*. 2021; 14: 2139-2164. PMID 34295184
76. Deer TR, Pope JE, Lamer TJ, et al. The Neuromodulation Appropriateness Consensus Committee on Best Practices for Dorsal Root Ganglion Stimulation. *Neuromodulation*. Jan 2019; 22(1): 1-35. PMID 30246899
77. National Institute for Health and Care Excellence (NICE). Spinal cord stimulation for chronic pain of neuropathic or ischaemic origin [TA159]. 2008; <https://www.nice.org.uk/guidance/ta159>. Accessed March 2, 2022.
78. Centers for Medicare and Medicaid Services. National Coverage Determination (NCD) for Electrical Nerve Stimulators (160.7). 1995; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=240&ncdver=1&bc=AAAAGAAAAAA&>. Accessed March 3, 2022.