



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy Orthotics for Progressive Scoliosis

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)
- [Endnotes](#)

Policy Number: 550

BCBSA Reference Number: 2.01.83 (For Plan internal use only)

NCD/LCD: N/A

Related Policies

Vertical Expandable Prosthetic Titanium Rib, #[305](#)

Policy¹

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

A rigid cervical-thoracic-lumbar-sacral or thoracic-lumbar-sacral orthosis may be considered **MEDICALLY NECESSARY** for the treatment of scoliosis in juvenile and adolescent individuals at high-risk of progression which meets the following criteria:

- Idiopathic spinal curve angle between 25° and 40°; AND
- Spinal growth has not been completed (Risser grade 0-3; no more than 1 year post menarche in females)

OR

- Idiopathic spinal curve angle greater than 20°; AND
- There is documented increase in the curve angle; AND
- At least 2 years' growth remain (Risser grade 0 or 1; pre-menarche in females).

Use of an orthosis for the treatment of scoliosis that does not meet the criteria above is considered **INVESTIGATIONAL**.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	Prior authorization is not required .

Commercial PPO and Indemnity	Prior authorization is not required .
Medicare HMO BlueSM	Prior authorization is not required .
Medicare PPO BlueSM	Prior authorization is not required .

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The above medical necessity criteria MUST be met for the following codes to be covered for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

HCPCS Codes

HCPCS codes:	Code Description
L1000	Cervical-thoracic-lumbar-sacral orthotic (CTLSO) (Milwaukee), inclusive of furnishing initial orthotic, including model
L1001	Cervical-thoracic-lumbar-sacral orthotic (CTLSO), immobilizer, infant size, prefabricated, includes fitting and adjustment
L1006	Scoliosis orthosis, sagittal-coronal control provided by a rigid lateral frame, extends from axilla to trochanter, includes all accessory pads, straps and interface, prefabricated item that has been trimmed, bent, molded, assembled, or otherwise customized to fit a specific patient by an individual with expertise
L1200	Thoracic-lumbar-sacral orthotic (TLSO), inclusive of furnishing initial orthotic only
L1300	Other scoliosis procedure, body jacket molded to patient model
L1310	Other scoliosis procedure, postoperative body jacket

Description

Scoliosis

Scoliosis is an abnormal lateral and rotational curvature of the vertebral column. Adolescent idiopathic scoliosis is the most common form of idiopathic scoliosis, defined by the U.S. Preventive Services Task Force as "a lateral curvature of the spine with onset at ≥ 10 years of age, no underlying etiology, and risk for progression during puberty."¹ Progression of the curvature during periods of rapid growth can result in deformity, accompanied by cardiopulmonary complications. Diagnosis is made clinically and radiographically. The curve is measured by the Cobb angle, which is the angle formed between intersecting lines drawn perpendicular to the top of the vertebrae of the curve and the bottom vertebrae of the curve. Patients with adolescent idiopathic scoliosis are also assessed for skeletal maturity, using the Risser sign, which describes the level of ossification of the iliac apophysis.

The Risser sign measures remaining spinal growth by progressive anterolateral to posteromedial ossification. Risser sign ranges from 0 (no ossification) to 5 (full bony fusion of the apophysis). Immature patients will have 0% to 25% ossification (Risser grade 0 or 1), while 100% ossification (Risser grade 5) indicates maturity with no spinal growth remaining. Children may progress from a Risser grade 1 to grade 5 over a brief (eg, 2-year), period.

Males and females are equally affected by scoliosis, but curve progression is up to 10 times more common in females than males.² Patients who are overweight or obese have a greater risk of presenting with larger Cobb angles and more advanced skeletal maturity, possibly due to delayed detection.³ A retrospective review of 341 patients with adolescent idiopathic scoliosis who underwent surgery at a single tertiary

pediatric hospital between 2013 and 2018 found that the major curve magnitude at presentation was significantly higher in patients with public compared to private insurance (50.0° versus 45.1°; $p=.0040$ and in Black compared to White patients (51.8° versus 47.0°; $p=.042$). Additionally, the odds of having an initial major curve magnitude $<40^\circ$ within the range of nonoperative treatment were 67% lower among Black patients with public insurance compared to Black patients with private insurance (odds ratio [OR], 0.33; 95% CI, 0.13 to 0.83; $p=.019$).⁴

Treatment

Treatment of scoliosis currently depends on 3 factors: the cause of the condition (idiopathic, congenital, secondary), the severity of the condition (degrees of the curve), and the growth of the patient remaining at the time of presentation. Children who have vertebral curves measuring between 25° and 40° with at least 2 years of growth remaining are considered to be at high risk of curve progression. Genetic markers to evaluate the risk of progression are also being evaluated. Because severe deformity may lead to compromised respiratory function and is associated with back pain in adulthood, surgical intervention with spinal fusion is typically recommended for curves that progress to 45° or more.

Bracing

Bracing is used to reduce the need for spinal fusion by slowing or preventing further progression of the curve during rapid growth. Commonly used brace designs include the Milwaukee, Wilmington, Boston, Charleston, and Providence orthoses. The longest clinical experience is with the Milwaukee cervical-thoracic-lumbar-sacral orthosis. Thoracic-lumbar-sacral orthoses, such as the Wilmington and Boston braces, are intended to improve tolerability and compliance for extended (>18 -hour) wear and are composed of lighter weight plastics with a low profile (underarm) design. The design of the nighttime Charleston and Providence braces is based on the theory that increased corrective forces will reduce the needed wear time (ie, daytime), thereby lessening social anxiety and improving compliance. The smart brace consists of a standard rigid brace with a microcomputer system, a force transducer, and an air-bladder control system to control the interface pressure. Braces that are more flexible than thoracic-lumbar-sacral orthoses or nighttime braces, such as the SpineCor® Scoliosis System, are also being evaluated. The SpineCor is composed of a thermoplastic pelvic base with stabilizing and corrective bands across the upper body.

Surgery

Fusionless surgical procedures, such as vertebral body stapling and vertebral body tethering, are being evaluated as alternatives to bracing. Both procedures use orthopedic devices off-label. The goal of these procedures is to reduce the rate of spine growth unilaterally, thus allowing the other side of the spine to “catch up.” The mechanism of action is believed to be down-regulation of the growth plate on the convex (outer) side by compression and stimulation of growth on the endplate of the concave side by distraction. In the current stapling procedure, nickel-titanium alloy staples with shape memory are applied to the convex side of the curve. The shape memory allows the prongs to be straight when cooled and clamp down into the bone when the staple returns to body temperature. Anterolateral tethering uses polyethylene ligaments that are attached to the convex side of the vertebral bodies by pedicle screws or staples. The ligament can be tightened to provide greater tension than the staple. The optimum degree of tension is not known. The polyethylene ligaments are more flexible than staples and are predicted to allow more spinal mobility. The goal of a fusionless growth modulating procedure is to reduce the curve and prevent progression, maintain spine mobility following correction, and provide an effective treatment option for patients who are noncompliant or who have a large curve but substantial growth is remaining. Observational data suggest that overweight patients may be at higher risk for scoliosis progression after surgery.⁵

Research Recommendations

The Scoliosis Research Society provided evidence-based recommendations in 2005,⁶ which were updated in 2015,⁷ for bracing studies to standardize inclusion criteria, methodologies, and outcome measures to facilitate comparison of brace trials. Janicki et al (2007),⁸ the first study to use the Scoliosis Research Society criteria, concluded that a brace should prevent progression in 70% of patients to be considered effective.⁸ The Scoliosis Research Society evidence review and recommendations may also aid in the evaluation of fusionless surgical treatments for scoliosis progression in children.

The Scoliosis Research Society review of the natural history of scoliosis indicated that skeletally immature patients and patients with larger curves (between 20° and 29°) are significantly more likely to have more than 5° curve progression.⁶ Brace treatment for idiopathic scoliosis is usually recommended for juveniles and adolescents with curves measuring between 25° and 40° who have not completed spinal growth, with maturity defined as Risser grade 4, or at least 2 years after menarche for girls.^{9,10} Bracing may also be recommended for curves greater than 20° in a patient who has a rapidly progressing curve with more than 2 years of growth remaining.

Success from brace treatment is most frequently defined as progression of less than 5° before skeletal maturity, although alternative definitions may include progression of less than 10° before skeletal maturity or preventing the curve from reaching the threshold for surgical intervention. Surgery is usually recommended when the curve magnitude exceeds 45° to 50° (before or at skeletal maturity), although many patients will not undergo surgery at this point. Based on this information, Scoliosis Research Society provided the following recommendations for brace studies on adolescent idiopathic scoliosis:

- “Optimal inclusion criteria for brace studies consist of: age is 10 years or older when the brace is prescribed, Risser [grade] 0-2, curve 25°-40°, and no prior treatment.”
- Outcomes of brace effectiveness should include all of the following:
 - “The percentage of patients with 5° or less curve progression and the percentage of patients who have 6° or more progression at skeletal maturity.”
 - The number of patients at the start and end of treatment exceeding 10°, 30°, and 50° Cobb angles, as these risk thresholds have potential health consequences in adulthood, such as back pain and curve progression.
 - “A minimum of 2-year follow-up beyond skeletal maturity for each patient who was ‘successfully’ treated with a brace to determine the percentage who subsequently required or had surgery recommended. The surgical indications must be documented.”
 - Clinically significant outcomes such as aesthetics, deformity progression, disability, pain, and quality of life.
- “Skeletal maturity should be considered achieved when <1 cm change in standing height has occurred on measurements made on 2 consecutive visits 6 months apart.... when Risser 4 is present and, in females, when the patient is 2 years after menarche.”
- “All patients, regardless of subjective reports of compliance, should be included in the results. This process makes ‘intent to treat’ analysis possible.... An ‘efficacy analysis’ ... should also be considered.”

Summary

Orthotic bracing attempts to slow spinal curve progression and reduce the need for fusion surgery in patients with juvenile or adolescent idiopathic scoliosis who are at high-risk of progression. Vertebral body stapling and vertebral body tethering, both fusionless surgical procedures, have been evaluated to determine whether the procedures could be used as alternatives to traditional orthotic bracing. This review does not address patients who are not at high-risk of progression or conventional fusion surgery for scoliosis, such as patients with Cobb angles measuring 45° or more.

For individuals who have juvenile or adolescent idiopathic scoliosis at high-risk of progression who receive a conventional rigid brace, the evidence includes a systematic review, a high-quality nonrandomized controlled trial, and 3 retrospective studies. Relevant outcomes are change in disease status, morbid events, quality of life, and treatment-related morbidity. Bracing has been considered the only option to prevent curve progression in juvenile or adolescent idiopathic scoliosis. The highest quality study on bracing is a sizable 2013 National Institutes of Health-sponsored trial that, using both randomized and observational arms, compared bracing with watchful waiting. This trial was stopped after interim analysis because of a significant benefit of bracing for the prevention of spinal fusion. Two retrospective studies with long-term follow-up (mean, 13 to 15 years) has also shown that curvature corrections with bracing were maintained. Another retrospective study demonstrated that nighttime bracing was more effective than a 24-hour brace for avoiding surgery and preventing curve progression, but investigators attributed this finding to likely noncompliance with the 24-hour brace. A systematic review and meta-analysis reported higher success with full-time and nighttime rigid braces compared to soft bracing or observation only. Based on several factors (evidence of efficacy, lack of alternative

treatment options, professional society recommendations, potential to prevent the need for a more invasive procedure), bracing with a conventional rigid brace is considered an option for the treatment of scoliosis in patients with a high-risk of curve progression. Curves have a high-risk of progression when they measure 25° or more, and spinal growth has not been completed, or when a 20° curve is progressively worsening and at least 2 years of growth remain. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have juvenile or adolescent idiopathic scoliosis at high-risk of progression who receive a microcomputer-controlled brace, the evidence includes a pilot randomized controlled trial. Relevant outcomes are change in disease status, morbid events, quality of life, and treatment-related morbidity. A pilot randomized trial using a microcomputer-controlled brace reported improved outcomes compared with the use of a standard rigid brace; however, the low number of individuals included in the trial (n=12) ultimately limited the interpretation of these results. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have juvenile or adolescent idiopathic scoliosis at high-risk of progression who receive a flexible brace, the evidence includes a randomized and a nonrandomized comparative study. Relevant outcomes are change in disease status, morbid events, quality of life, and treatment-related morbidity. One randomized controlled trial evaluating a flexible brace did not show equivalent outcomes compared with conventional brace designs. Another study has suggested the flexible brace might improve outcomes compared with no treatment, but this study had design flaws, which interfered with drawing significant conclusions from the study. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
10/2024	Clarified coding information.
6/2024	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2023	Annual policy review. Minor editorial changes to policy statements; intent unchanged.
6/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
5/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
6/2018	Investigational statement on vertebral body stapling and vertebral body tethering removed; title changed. Effective 6/1/2018. Annual policy review. Policy section clarified; statements otherwise unchanged.
12/2016	Annual policy review. New references added.
10/2015	Annual policy review. New investigational indications described. Clarified coding information. Effective 10/1/2015.
6/2013	Annual policy review. New references added.
5/1/12	New policy describing ongoing coverage and non-coverage.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

References

1. U.S. Preventive Services Task Force (USPSTF). Final Recommendation Statement: Adolescent Idiopathic Scoliosis: Screening. 2018; <https://www.uspreventiveservicestaskforce.org/uspstf/document/RecommendationStatementFinal/adolescent-idiopathic-scoliosis-screening>. Accessed March 6, 2023.
2. American Academy of Orthopaedic Surgeons (AAOS). Idiopathic Scoliosis in Children and Adolescents. April 2021; <https://orthoinfo.aaos.org/en/diseases--conditions/idiopathic-scoliosis-in-children-and-adolescents>. Accessed March 6, 2023.
3. Margalit A, McKean G, Constantine A, et al. Body Mass Hides the Curve: Thoracic Scoliometer Readings Vary by Body Mass Index Value. *J Pediatr Orthop*. Jun 2017; 37(4): e255-e260. PMID 27861214
4. Heffernan MJ, Younis M, Song B, et al. Disparities in Pediatric Scoliosis: The Impact of Race and Insurance Type on Access to Nonoperative Treatment for Adolescent Idiopathic Scoliosis. *J Pediatr Orthop*. Sep 01 2022; 42(8): 427-431. PMID 35856501
5. Mishreky A, Parent S, Miyajiri F, et al. Body mass index affects outcomes after vertebral body tethering surgery. *Spine Deform*. May 2022; 10(3): 563-571. PMID 35013996
6. Richards BS, Bernstein RM, D'Amato CR, et al. Standardization of criteria for adolescent idiopathic scoliosis brace studies: SRS Committee on Bracing and Nonoperative Management. *Spine (Phila Pa 1976)*. Sep 15 2005; 30(18): 2068-75; discussion 2076-7. PMID 16166897
7. Negrini S, Hresko TM, O'Brien JP, et al. Recommendations for research studies on treatment of idiopathic scoliosis: Consensus 2014 between SOSORT and SRS non-operative management committee. *Scoliosis*. 2015; 10: 8. PMID 25780381
8. Janicki JA, Poe-Kochert C, Armstrong DG, et al. A comparison of the thoracolumbosacral orthoses and providence orthosis in the treatment of adolescent idiopathic scoliosis: results using the new SRS inclusion and assessment criteria for bracing studies. *J Pediatr Orthop*. Jun 2007; 27(4): 369-74. PMID 17513954
9. Fayssox RS, Cho RH, Herman MJ. A history of bracing for idiopathic scoliosis in North America. *Clin Orthop Relat Res*. Mar 2010; 468(3): 654-64. PMID 19462214
10. Schiller JR, Thakur NA, Ebersson CP. Brace management in adolescent idiopathic scoliosis. *Clin Orthop Relat Res*. Mar 2010; 468(3): 670-8. PMID 19484317
11. Wall EJ, Reynolds JE, Jain VV, et al. Spine Growth Modulation in Early Adolescent Idiopathic Scoliosis: Two-Year Results of Prospective US FDA IDE Pilot Clinical Safety Study of Titanium Clip-Screw Implant. *Spine Deform*. Sep 2017; 5(5): 314-324. PMID 28882349
12. Weinstein SL, Dolan LA, Wright JG, et al. Effects of bracing in adolescents with idiopathic scoliosis. *N Engl J Med*. Oct 17 2013; 369(16): 1512-21. PMID 24047455
13. Aulisa AG, Toniolo RM, Falciglia F, et al. Long-term results after brace treatment with Progressive Action Short Brace in adolescent idiopathic scoliosis. *Eur J Phys Rehabil Med*. Jun 2021; 57(3): 406-413. PMID 32990686
14. Aulisa AG, Guzzanti V, Falciglia F, et al. Curve progression after long-term brace treatment in adolescent idiopathic scoliosis: comparative results between over and under 30 Cobb degrees - SOSORT 2017 award winner. *Scoliosis Spinal Disord*. 2017; 12: 36. PMID 29094108
15. Costa L, Schlosser TPC, Jimale H, et al. The Effectiveness of Different Concepts of Bracing in Adolescent Idiopathic Scoliosis (AIS): A Systematic Review and Meta-Analysis. *J Clin Med*. May 15 2021; 10(10). PMID 34063540
16. Lou E, Hill D, Raso J, et al. Smart brace versus standard rigid brace for the treatment of scoliosis: a pilot study. *Stud Health Technol Inform*. 2012; 176: 338-41. PMID 22744524
17. Wong MS, Cheng JC, Lam TP, et al. The effect of rigid versus flexible spinal orthosis on the clinical efficacy and acceptance of the patients with adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)*. May 20 2008; 33(12): 1360-5. PMID 18496349
18. Guo J, Lam TP, Wong MS, et al. A prospective randomized controlled study on the treatment outcome of SpineCor brace versus rigid brace for adolescent idiopathic scoliosis with follow-up according to the SRS standardized criteria. *Eur Spine J*. Dec 2014; 23(12): 2650-7. PMID 24378629

19. Plewka B, Sibiński M, Synder M, et al. Clinical assessment of the efficacy of SpineCor brace in the correction of postural deformities in the course of idiopathic scoliosis. *Pol Orthop Traumatol*. Mar 26 2013; 78: 85-9. PMID 23535882
20. Plewka B, Sibiński M, Synder M, et al. Radiological evaluation of treatment with SpineCor brace in children with idiopathic spinal scoliosis. *Ortop Traumatol Rehabil*. Jun 28 2013; 15(3): 227-34. PMID 23897999
21. Cuddihy L, Danielsson AJ, Cahill PJ, et al. Vertebral Body Stapling versus Bracing for Patients with High-Risk Moderate Idiopathic Scoliosis. *Biomed Res Int*. 2015; 2015: 438452. PMID 26618169
22. Murray E, Tung R, Sherman A, et al. Continued vertebral body growth in patients with juvenile idiopathic scoliosis following vertebral body stapling. *Spine Deform*. Apr 2020; 8(2): 221-226. PMID 32026438
23. Bumpass DB, Fuhrhop SK, Schootman M, et al. Vertebral Body Stapling for Moderate Juvenile and Early Adolescent Idiopathic Scoliosis: Cautions and Patient Selection Criteria. *Spine (Phila Pa 1976)*. Dec 2015; 40(24): E1305-14. PMID 26655807
24. Theologis AA, Cahill P, Auriemma M, et al. Vertebral body stapling in children younger than 10 years with idiopathic scoliosis with curve magnitude of 30° to 39°. *Spine (Phila Pa 1976)*. Dec 01 2013; 38(25): E1583-8. PMID 23963018
25. Laituri CA, Schwend RM, Holcomb GW. Thoracoscopic vertebral body stapling for treatment of scoliosis in young children. *J Laparoendosc Adv Surg Tech A*. Oct 2012; 22(8): 830-3. PMID 23039706
26. O'leary PT, Sturm PF, Hammerberg KW, et al. Convex hemiepiphysiodesis: the limits of vertebral stapling. *Spine (Phila Pa 1976)*. Sep 01 2011; 36(19): 1579-83. PMID 21681138
27. Betz RR, Ranade A, Samdani AF, et al. Vertebral body stapling: a fusionless treatment option for a growing child with moderate idiopathic scoliosis. *Spine (Phila Pa 1976)*. Jan 15 2010; 35(2): 169-76. PMID 20081512
28. Zhu F, Qiu X, Liu S, et al. Minimum 3-year experience with vertebral body tethering for treating scoliosis: A systematic review and single-arm meta-analysis. *J Orthop Surg (Hong Kong)*. 2022; 30(3): 10225536221137753. PMID 36420934
29. U.S. Food and Drug Administration (FDA). SUMMARY OF SAFETY AND PROBABLE BENEFIT (SSPB): The Tether Vertebral Body Tethering System. 2019; https://www.accessdata.fda.gov/cdrh_docs/pdf19/H190005b.pdf. Accessed March 6, 2023.
30. Courvoisier A, Eid A, Bourgeois E, et al. Growth tethering devices for idiopathic scoliosis. *Expert Rev Med Devices*. Jul 2015; 12(4): 449-56. PMID 26027921
31. Samdani AF, Ames RJ, Kimball JS, et al. Anterior vertebral body tethering for idiopathic scoliosis: two-year results. *Spine (Phila Pa 1976)*. Sep 15 2014; 39(20): 1688-93. PMID 24921854
32. Samdani AF, Ames RJ, Kimball JS, et al. Anterior vertebral body tethering for immature adolescent idiopathic scoliosis: one-year results on the first 32 patients. *Eur Spine J*. Jul 2015; 24(7): 1533-9. PMID 25510515
33. Pehlivanoglu T, Oltulu I, Erdag Y, et al. Double-sided vertebral body tethering of double adolescent idiopathic scoliosis curves: radiographic outcomes of the first 13 patients with 2 years of follow-up. *Eur Spine J*. Jul 2021; 30(7): 1896-1904. PMID 33611658
34. Meyers J, Eaker L, Zhang J, et al. Vertebral Body Tethering in 49 Adolescent Patients after Peak Height Velocity for the Treatment of Idiopathic Scoliosis: 2-5 Year Follow-Up. *J Clin Med*. Jun 02 2022; 11(11). PMID 35683548
35. Negrini S, Donzelli S, Aulisa AG, et al. 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis Spinal Disord*. 2018; 13: 3. PMID 29435499
36. Scoliosis Research Society (SRS). Adolescent Idiopathic Scoliosis. n.d.; <http://www.srs.org/professionals/online-education-and-resources/conditions-and-treatments/adolescent-idiopathic-scoliosis>. Accessed March 6, 2023.
37. Scoliosis Research Society (SRS)/Pediatric Orthopaedic Society of North America (POSNA). Joint SRS/POSNA Position Statement on Payor Coverage for Anterior Fusionless Scoliosis Technologies for Immature Patients with Idiopathic Scoliosis. April 2020; <https://posna.org/POSNA/media/Documents/Position%20Statements/Why-Should-Insurance-Cover-AVBT-April-2020.pdf>. Accessed March 5, 2023.
38. National Institute of Arthritis and Musculoskeletal and Skin Diseases. Questions and Answers about Scoliosis in Children and Adolescents. December 2019; <https://www.niams.nih.gov/health-topics/scoliosis>. Accessed March 6, 2023.

39. National Institute for Health and Care Excellence (NICE). Interventional procedures guidance: Vertebral body tethering for idiopathic scoliosis in children and young people [IPG728]. June 29, 2022; <https://www.nice.org.uk/guidance/ipg728>. Accessed March 6, 2023.
40. Grossman DC, Curry SJ, Owens DK, et al. Screening for Adolescent Idiopathic Scoliosis: US Preventive Services Task Force Recommendation Statement. JAMA. Jan 09 2018; 319(2): 165-172. PMID 29318284

Endnotes

¹ Based on expert opinion