



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an Independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy Suprachoroidal Delivery of Pharmacologic Agents

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 609

BCBSA Reference Number: 9.03.19A (For Plan internal use only)
NCD/LCD: NA

Related Policies

- Photodynamic Therapy #[599](#)
- Transpupillary Thermotherapy for Treatment of Choroidal Neovascularization, #[600](#)
- Epiretinal Radiation for Age-Related Macular Degeneration, #[610](#)
- Intravitreal Angiogenesis Inhibitors for Choroidal Vascular Conditions, #[343](#)

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Suprachoroidal delivery of a pharmacologic agent is [INVESTIGATIONAL](#).

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO Blue SM	This is not a covered service.
Medicare PPO Blue SM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

CPT Codes

No specific CPT codes.

Description

Age-related macular degeneration and diabetic neuropathy are the most prevalent causes of visual impairment and are classified as posterior segment ocular diseases due to the location of the affected eye structures in the back two thirds of the eye. These posterior structures include the vitreous humor, retina, choroid, macula, and optic nerve. Delivery of pharmacologic agents to the suprachoroidal space is being investigated for treatment of posterior eye segment diseases, as current modes of drug delivery to the eye (topical and systemic) often lack bioavailability and fail to deliver therapeutic levels of drugs to the retina and posterior segment structures. Examples of drugs used via suprachoroidal injection to treat age related macular degeneration include ranibizumab and pegaptanib. For treatment of diabetic neuropathy, an example is triamcinolone acetonide.

A microcannula system (the delivery method of suprachoroidal injection) originates from this same idea and combines a drug delivery channel with a fiber-optic light source to allow transmission of light to the tip of the microcannula for illumination and guidance. One potential advantage of suprachoroidal injection would be the ability to minimize systemic side effects while delivering higher local tissue levels of drugs, assuming high local levels lead to improved outcomes. Weighed against this potential benefit is the risk of localized tissue damage from the microcannula.

An example of a device for suprachoroidal delivery of a pharmacologic agent includes the iTrack™ from iScience Interventional. All devices for suprachoroidal delivery of a pharmacologic agent are considered investigational regardless of the commercial name, the manufacturer or FDA approval status.

Summary

Evidence remains insufficient to evaluate whether suprachoroidal delivery of pharmacologic agents improves the net health outcome. Thus, this procedure is considered investigational.

Policy History

Date	Action
11/2022	Annual policy review. Policy updated with literature review through October 2022. No references added. Policy statements unchanged.
2/2020	Policy updated with literature review through February 1, 2020, references added. Policy statements unchanged.
2/2015	Annual policy review. New references added.
1/2014	Updated to remove deleted CPT code 0186T.
11/2011-4/2012	Medical policy ICD 10 remediation: Formatting, editing and coding updates. No changes to policy statements.
2/2011	Reviewed - Medical Policy Group – Psychiatry and Ophthalmology. No changes to policy statements.
1/19/2011	Medical Policy 609 -effective 1/19/2011 describing ongoing non-coverage.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

References

1. US Food and Drug Administration (FDA). Summary of Safety and Effectiveness: iScience Surgical Ophthalmic Microcannula (iTRACK) - June 22, 2004.
<http://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pmn&id=K041108>. Accessed November 24, 2014.
2. Gilger BC, Mandal A, Shah S, et al. Episcleral, intrascleral, and suprachoroidal routes of ocular drug delivery - recent research advances and patents. *Recent Pat Drug Deliv Formul.* 2014;8(2):81-91. PMID 25001638
3. Olsen, T. Drug delivery to the suprachoroidal space shows promise. *Retina Today*; March 2007. http://retinatoday.com/2007/03/0307_10.html. Accessed November 24, 2013.
4. Del Amo EM, Urtti A. Current and future ophthalmic drug delivery systems. A shift to the posterior segment. *Drug Discov Today.* Feb 2008;13(3-4):135-143. PMID 18275911
5. Rizzo S, Ebert FG, Bartolo ED, et al. Suprachoroidal drug infusion for the treatment of severe subfoveal hard exudates. *Retina.* Apr 2012;32(4):776-784. PMID 21817959
6. Tetz M, Rizzo S, Augustin AJ. Safety of submacular suprachoroidal drug administration via a microcatheter: retrospective analysis of European treatment results. *Ophthalmologica.* 2012;227(4):183-189. PMID 22354263