



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Multitarget Polymerase Chain Reaction Testing for Diagnosis of Bacterial Vaginosis

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 711

BCBSA Reference Number: 2.04.127 (For Plan internal use only)

NCD/LCD: N/A

Related Policies

Identification of Microorganisms Using Nucleic Acid Probe, #555

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity Medicare HMO BlueSM and Medicare PPO BlueSM Members

Multitarget polymerase chain reaction (PCR) testing for diagnosis of bacterial vaginosis is considered **INVESTIGATIONAL**.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.
Medicare HMO BlueSM	This is not a covered service.
Medicare PPO BlueSM	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

The following CPT codes are considered investigational for Commercial Members: Managed Care (HMO and POS), PPO, Indemnity, Medicare HMO Blue and Medicare PPO Blue:

CPT Codes

CPT codes:	Code Description
0330U	Infectious agent detection by nucleic acid (DNA or RNA), vaginal pathogen panel, identification of 27 organisms, amplified probe technique, vaginal swab
0352U	Infectious disease (bacterial vaginosis and vaginitis), multiplex amplified probe technique, for detection of bacterial vaginosis–associated bacteria (BVAB-2, Atopobium vaginae, and Megasphaera type 1), algorithm reported as detected or not detected and separate detection of Candida species (C. albicans, C. tropicalis, C. parapsilosis, C. dubliniensis), Candida glabrata/Candida krusei, and trichomonas vaginalis, vaginal-fluid specimen, each result reported as detected or not detected
0353U	Infectious agent detection by nucleic acid (DNA), Chlamydia trachomatis and Neisseria gonorrhoeae, multiplex amplified probe technique, urine, vaginal, pharyngeal, or rectal, each pathogen reported as detected or not detected
81513	Infectious disease, bacterial vaginosis, quantitative real-time amplification of RNA markers for Atopobium vaginae, Gardnerella vaginalis, and Lactobacillus species, utilizing vaginal-fluid specimens, algorithm reported as a positive or negative result for bacterial vaginosis
81514	Infectious disease, bacterial vaginosis and vaginitis, quantitative real-time amplification of DNA markers for Gardnerella vaginalis, Atopobium vaginae, Megasphaera type 1, Bacterial Vaginosis Associated Bacteria-2 (BVAB-2), and Lactobacillus species (L. crispatus and L. jensenii), utilizing vaginal-fluid specimens, algorithm reported as a positive or negative for high likelihood of bacterial vaginosis, includes separate detection of Trichomonas vaginalis and/or Candida species (C. albicans, C. tropicalis, C. parapsilosis, C. dubliniensis), Candida glabrata, Candida krusei, when reported

Description

Bacterial Vaginosis

BV is a condition caused by an imbalance in the normal bacteria vaginal flora. It is common, especially in women of reproductive age. While there is no single known etiologic agent, there is a shift in vaginal flora that involves depletion of hydrogen peroxide-producing Lactobacillus species with a rise in vaginal pH and overgrowth of other bacteria, including Gardnerella vaginalis, Mycoplasma hominis, Peptostreptococcus, Mobiluncus species, and other anaerobic gram-negative rods.

Vaginal culture is not an appropriate diagnostic method to identify BV because BV is not caused by the presence of a particular bacterial species.

Various commercial tests provide rapid and accurate pH evaluation and amine detection.

For example, automated devices that measure the volatile gases produced from vaginal samples and a colorimetric pH test are commercially available.

Nucleic acid probes of DNA fragments are available to detect and quantify specific bacteria in vaginal fluid samples. Polymerase chain reaction (PCR) methods extract and amplify the DNA fragments using either universal or specific primers. The result can be qualitative (to assess whether a specific microorganism is present) or quantitative (to assess how many microorganisms are present). The technology can be used to measure multiple organisms (eg, those known to be associated with BV) at the same time and is commercially available as multitarget PCR testing.

(Policy # [555](#) addresses the use of direct or amplified nucleic acid probes with or without quantification to detect microorganisms of clinical significance, including single microorganisms associated with BV.

Proposed Multitarget PCR Tests

The SureSwab Total (Quest Diagnostics) test involves obtaining vaginal swab specimens, extracting total DNA, and quantitating the 4 types of bacteria using PCR. Results are reported as log cells per milliliter for each organism and concentrations of all *Lactobacilli* species are reported together then classified into 1 of the following 3 categories: not supportive, equivocal, and supportive.

A classification of *not supportive* of BV diagnosis is based on:

- The presence of *Lactobacillus* species, *G. vaginalis* levels <6.0 log cells/mL, and absence of *Atopobium vaginae* and *Megasphaera* species; or
- The absence of *Lactobacillus* species, *G. vaginalis* levels <6.0 log cells/mL, and absence of *A. vaginae* and *Megasphaera* species; or
- The absence of all targeted organisms.

A classification of equivocal is based on:

- The presence of *Lactobacillus* species, plus *G. vaginalis* at least 6.0 log cells/mL, and/or presence of *A. vaginae* and/or *Megasphaera* species.

A classification of supportive of BV diagnosis is based on the absence of *Lactobacillus* species, and presence of *G. vaginalis* levels of at least 6.0 log cells/mL, and presence of *A. vaginae* and/or *Megasphaera* species.

Another product, the BD Max (Becton, Dickinson), tests for markers of BV and vaginitis. The test uses a similar process to that described for SureSwab. Vaginal swab specimens are collected, DNA is extracted, and real-time PCR is used to quantitate targeted organisms. Results of BV marker tests are not reported for individual organisms. Instead, qualitative BV results are reported as positive or negative for BV based on the relative quantity of the various organisms. The Aptima BV Assay was cleared by the U.S. Food and Drug Administration with the BD Max as the predicate device. The Aptima assay is a nucleic acid amplification test (NAAT) for detection and quantitation of ribosomal RNA.

Medical Diagnostics Laboratory offers a Bacterial Vaginosis Panel. Markers are assessed using real-time PCR and *Lactobacillus* is profiled using quantitative PCR. GenPath Diagnostics also offers a bacterial vaginosis test.

The NuSwab® Select BV test (Laboratory Corporation of America) uses semiquantitative PCR analysis of 3 predictive marker organisms of vaginal dysbiosis to generate a total score that is associated with the presence or absence of BV. In this test system, samples with a total score of 0 to 1 are considered negative for BV, samples with a score of 3 to 6 are positive for BV, and samples with a score of 2 are indeterminate for BV.

Several of the manufacturers of the BV tests also have extensions that include other causes of vaginitis such as *Trichomonas vaginalis* and *Candidiasis* species.

Summary

Bacterial vaginosis (BV) is a common medical condition resulting from an imbalance in the normal vaginal flora. Although the identification of *Gardnerella vaginalis* has traditionally been associated with BV, there is no single etiologic agent. Most cases are asymptomatic, and most symptomatic cases can be diagnosed using clinical and microscopic evaluation. Multitarget polymerase chain reaction (PCR) testing is proposed as an alternative to currently available laboratory tests to diagnose BV. This test may improve outcomes if it is a more accurate and reliable method to diagnose BV.

Summary of Evidence

In individuals who have signs or symptoms of BV who receive multitarget PCR testing, the evidence includes several prospective studies on technical performance and diagnostic accuracy. The relevant outcomes are test validity, symptoms, and change in disease status. Several studies have evaluated the diagnostic accuracy of multitarget PCR tests for BV, including 5 studies evaluating commercially available tests. The studies found sensitivities between 84% and 95% and specificities between 85% and 97% compared with standard methods of diagnosis. Most studies used a combination of the Amsel criteria and Nugent scoring as the reference standard. There is a lack of direct evidence on the clinical utility of PCR testing for BV (ie, studies showing that testing leads to better patient management decisions and/or better health outcomes than current approaches). Moreover, a chain of evidence does not currently support multitarget testing because most symptomatic women can be diagnosed with a standard workup. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
10/2022	Clarified coding information.
7/2022	Clarified coding information.
2/2022	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
1/2021	Clarified coding information.
1/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
2/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
8/2017	Annual policy review. New references added.
1/2016	Annual policy review. New references added.
3/2015	New medical policy describing investigational indications. Effective 3/1/2015.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

- Centers for Disease Control and Prevention (CDC). Bacterial Vaginosis (BV) Statistics. <http://www.cdc.gov/std/bv/stats.htm>. Accessed November 18, 2021.
- Amsel R, Totten PA, Spiegel CA, et al. Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med.* Jan 1983; 74(1): 14-22. PMID 6600371
- Landers DV, Wiesenfeld HC, Heine RP, et al. Predictive value of the clinical diagnosis of lower genital tract infection in women. *Am J Obstet Gynecol.* Apr 2004; 190(4): 1004-10. PMID 15118630
- Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. *J Clin Microbiol.* Feb 1991; 29(2): 297-301. PMID 1706728
- Ison CA, Hay PE. Validation of a simplified grading of Gram stained vaginal smears for use in genitourinary medicine clinics. *Sex Transm Infect.* Dec 2002; 78(6): 413-5. PMID 12473800
- Hilbert DW, Smith WL, Chadwick SG, et al. Development and Validation of a Highly Accurate Quantitative Real-Time PCR Assay for Diagnosis of Bacterial Vaginosis. *J Clin Microbiol.* Apr 2016; 54(4): 1017-24. PMID 26818677

7. Thompson A, Timm K, Borders N, et al. Diagnostic performance of two molecular assays for the detection of vaginitis in symptomatic women. *Eur J Clin Microbiol Infect Dis.* Jan 2020; 39(1): 39-44. PMID 31502121
8. Gaydos CA, Beqaj S, Schwebke JR, et al. Clinical Validation of a Test for the Diagnosis of Vaginitis. *Obstet Gynecol.* Jul 2017; 130(1): 181-189. PMID 28594779
9. Schwebke JR, Gaydos CA, Nyirjesy P, et al. Diagnostic Performance of a Molecular Test versus Clinician Assessment of Vaginitis. *J Clin Microbiol.* Jun 2018; 56(6). PMID 29643195
10. Aguirre-Quinonero A, Castillo-Sedano IS, Calvo-Muro F, et al. Accuracy of the BD MAX vaginal panel in the diagnosis of infectious vaginitis. *Eur J Clin Microbiol Infect Dis.* May 2019; 38(5): 877-882. PMID 30685805
11. van den Munckhof EHA, van Sitter RL, Boers KE, et al. Comparison of Amsel criteria, Nugent score, culture and two CE-IVD marked quantitative real-time PCRs with microbiota analysis for the diagnosis of bacterial vaginosis. *Eur J Clin Microbiol Infect Dis.* May 2019; 38(5): 959-966. PMID 30903536
12. Food and Drug Administration. Evaluation of Automatic Class III Designation For BD Max Vaginal Panel: Decision Summary. 2016; https://www.accessdata.fda.gov/cdrh_docs/reviews/DEN160001.pdf. Accessed November 18, 2021.
13. Cartwright CP, Pherson AJ, Harris AB, et al. Multicenter study establishing the clinical validity of a nucleic-acid amplification-based assay for the diagnosis of bacterial vaginosis. *Diagn Microbiol Infect Dis.* Nov 2018; 92(3): 173-178. PMID 29937222
14. Cartwright CP, Lembke BD, Ramachandran K, et al. Development and validation of a semiquantitative, multitarget PCR assay for diagnosis of bacterial vaginosis. *J Clin Microbiol.* Jul 2012; 50(7): 2321-9. PMID 22535982
15. Schwebke JR, Taylor SN, Ackerman R, et al. Clinical Validation of the Aptima Bacterial Vaginosis and Aptima Candida/Trichomonas Vaginitis Assays: Results from a Prospective Multicenter Clinical Study. *J Clin Microbiol.* Jan 28 2020; 58(2). PMID 31748322
16. Richter SS, Otiso J, Goje OJ, et al. Prospective Evaluation of Molecular Assays for Diagnosis of Vaginitis. *J Clin Microbiol.* Dec 23 2019; 58(1). PMID 31694966
17. Kusters JG, Reuland EA, Bouter S, et al. A multiplex real-time PCR assay for routine diagnosis of bacterial vaginosis. *Eur J Clin Microbiol Infect Dis.* Sep 2015; 34(9): 1779-85. PMID 26143346
18. Romyantseva TA, Bellen G, Romanuk TN, et al. Utility of microscopic techniques and quantitative real-time polymerase chain reaction for the diagnosis of vaginal microflora alterations. *J Low Genit Tract Dis.* Apr 2015; 19(2): 124-8. PMID 25023332
19. Romyantseva T, Shipitsyna E, Guschin A, et al. Evaluation and subsequent optimizations of the quantitative AmpliSens Florocenosis/Bacterial vaginosis-FRT multiplex real-time PCR assay for diagnosis of bacterial vaginosis. *APMIS.* Dec 2016; 124(12): 1099-1108. PMID 27714844
20. van der Veer C, van Houdt R, van Dam A, et al. Accuracy of a commercial multiplex PCR for the diagnosis of bacterial vaginosis. *J Med Microbiol.* Sep 2018; 67(9): 1265-1270. PMID 29985123
21. Committee on Practice Bulletins-Obstetrics, The American College of Obstetricians and Gynecologists. Practice bulletin no. 130: prediction and prevention of preterm birth. *Obstet Gynecol.* Oct 2012; 120(4): 964-73. PMID 22996126
22. Vaginitis in Nonpregnant Patients: ACOG Practice Bulletin, Number 215. *Obstet Gynecol.* Jan 2020; 135(1): e1-e17. PMID 31856123
23. Workowski KA, Bachmann LH, Chan PA, et al. Sexually Transmitted Infections Treatment Guidelines, 2021. *MMWR Recomm Rep.* Jul 23 2021; 70(4): 1-187. PMID 34292926
24. National Institute for Health and Care Excellence (NICE). Clinical Guideline: Antenatal Care for Uncomplicated Pregnancies [CG62]. 2008; <https://www.nice.org.uk/guidance/cg62/>. Updated February 4, 2019. Accessed November 18, 2021.
25. U.S. Preventive Services Task Force. Bacterial Vaginosis in Pregnancy to Prevent Preterm Delivery: Screening. 2020; <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/bacterial-vaginosis-in-pregnancy-to-prevent-preterm-delivery-screening#:~:text=The%20USPSTF%20recommends%20against%20screening%20for%20bacterial%20vaginosis,who%20are%20at%20increased%20risk%20for%20preterm%20delivery..> Accessed November 19, 2021.