



MASSACHUSETTS

Blue Cross Blue Shield of Massachusetts is an independent Licensee of the Blue Cross and Blue Shield Association

Medical Policy

Minimally Invasive Ablation Procedures for Morton and Other Peripheral Neuromas

Table of Contents

- [Policy: Commercial](#)
- [Policy: Medicare](#)
- [Authorization Information](#)
- [Coding Information](#)
- [Description](#)
- [Policy History](#)
- [Information Pertaining to All Policies](#)
- [References](#)

Policy Number: 719

BCBSA Reference Number: 7.01.147 (For Plan internal use only)

Related Policies

None

Policy

Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity

Minimally invasive ablation procedures, including intralesional alcohol injection, radiofrequency ablation, and cryoablation, are considered **INVESTIGATIONAL** for the treatment of Morton and other peripheral neuromas.

Prior Authorization Information

Inpatient

- For services described in this policy, precertification/preauthorization **IS REQUIRED** for all products if the procedure is performed **inpatient**.

Outpatient

- For services described in this policy, see below for products where prior authorization **might be required** if the procedure is performed **outpatient**.

	Outpatient
Commercial Managed Care (HMO and POS)	This is not a covered service.
Commercial PPO and Indemnity	This is not a covered service.

CPT Codes / HCPCS Codes / ICD Codes

Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage as it applies to an individual member.

Providers should report all services using the most up-to-date industry-standard procedure, revenue, and diagnosis codes, including modifiers where applicable.

The following codes are included below for informational purposes only; this is not an all-inclusive list.

According to the policy statement above, the following CPT codes are considered investigational for the conditions listed for **Commercial Members: Managed Care (HMO and POS), PPO, and Indemnity:**

CPT Codes

CPT codes:	Code Description
64632	Destruction by neurolytic agent; plantar common digital nerve
64640	Destruction by neurolytic agent; other peripheral nerve or branch

Description

Neuroma

A neuroma is a pathology of a peripheral nerve that develops as part of a normal reparative process. Neuromas may develop after nerve injury or result from chronic irritation, pressure, stretch, poor repair of nerve lesions or previous neuromas, laceration, crush injury, or blunt trauma.¹ Neuromas typically appear 6 to 10 weeks after trauma, with most presenting within 1 to 12 months after injury or surgery. They may gradually enlarge over 2 to 3 years and may or may not be painful. Pain from a neuroma may be secondary to traction on the nerve by scar tissue, compression of the sensitive nerve endings by adjacent soft tissues, ischemia of the nervous tissue, or ectopic foci of ion channels that elicit neuropathic pain. Patients may describe the pain as low-intensity dull pain or intense paroxysmal burning pain, often triggered by external stimuli such as touch or temperature. Neuroma formation has been implicated as a contributor of neuropathic pain in residual limb pain, postthoracotomy, postmastectomy, and postherniorrhaphy pain syndromes. Neuromas may coexist with phantom pain or can predispose to it.

Morton Neuroma

Morton neuroma is a common and painful compression neuropathy of the common digital nerve of the foot that may also be referred to as interdigital neuroma, interdigital neuritis, and interdigital or Morton metatarsalgia.^{1,2,3} It is histologically characterized by perineural fibrosis, endoneurial edema, axonal degeneration, and local vascular proliferation. Thus, some investigators do not consider Morton neuroma to be a true neuroma; instead, they consider it to be an entrapment neuropathy occurring secondary to compression of the common digital nerve under the overlying transverse metatarsal ligament. Morton neuroma appears 10-fold more often in women than in men, with an average age at presentation of around 50 years.⁴

The pain associated with Morton neuroma is usually throbbing, burning, or shooting, and localized to the plantar aspect of the foot. It is typically located between the 3rd and 4th metatarsal heads, although it may appear in other proximal locations.^{1,2} The pain may radiate to the toes and can be associated with paresthesia. The pain can be severe, and the condition may become debilitating to the extent that patients are apprehensive about walking or touching their foot to the ground. It is aggravated by walking in shoes with a narrow toe box or high heels that cause excessive pronation and excessive forefoot pressure; removal of tight shoes typically relieves the pain.

Diagnosis

Although a host of imaging methods are used to diagnosis Morton neuroma, including plain radiographs, magnetic resonance imaging, and ultrasonography, objective findings are unique to this condition and are primarily used to establish a clinical diagnosis.¹ Thus, a patient's toes often show splaying or divergence. Patients may describe the feeling of a "lump" on the foot bottom or a feeling of walking on a rolled-up or wrinkled sock. Clinical examination with medial and lateral compression may reproduce the painful symptoms with a palpable "click" on interspace compression (Mulder sign).⁵

Treatment

Management of patients diagnosed with Morton neuroma typically starts with conservative approaches, such as the use of metatarsal pads in shoes and orthotic devices that alter supination and pronation of the affected foot.³ These approaches try to reduce pressure and irritation of the affected nerve. They may provide relief, but do not alter the underlying pathology. There is scant evidence to support the effectiveness or comparative effectiveness of these practices.^{2,6,7} In a case series, Bennett et al (1995) evaluated a 3-stage protocol of "stepped care" through which private practice patients (N=115) advanced from stage I (education plus footwear modifications, and a metatarsal pad) to stage II (steroid injections with local anesthetic or local anesthetic alone), and into stage III (surgical resection) if stages I and II were not relieved within 3 months.⁸ Overall, 97 (85%) of 115 patients believed that pain had been reduced with the treatment program. However, 24 (21%) patients eventually required surgical excision of the nerve, and 23 (96%) of them had satisfactory results.

Minimally Invasive Ablation Procedures

Several minimally invasive procedures to treat refractory Morton and other peripheral neuromas are aimed at in situ destruction of the pathology, including intralesional alcohol injection, radiofrequency ablation (RFA) and cryoablation (also known as cryoneurolysis, cryolysis, and cryoanalgesia).²

Dehydrated ethanol has been shown to inhibit nerve function in vitro, has high affinity for nerve tissue, and causes direct damage to nerve cells via dehydration, cell necrosis, and precipitation of protoplasm, leading to neuritis and a pattern of Wallerian degeneration.² Technically, ethanol is a sclerosant that causes chemical neurolysis of the nerve pathology but is considered an ablative procedure for this evidence review. The use of ultrasound guidance during this procedure has been shown to increase surgical accuracy, improve outcomes, and shorten procedure duration. RFA uses heat generated by an electrode that conducts electromagnetic energy into a tissue or lesion to denature proteins and destroy cells. RFA is used to ablate a wide range of tissues or lesions, including osteoid osteoma; cardiovascular system pathologies; cervical pain syndromes; liver, lung, and other cancers; and varicosities.^{9,10,11,12,13,14,15,16,17,18,19,20} Cryoablation uses coolant to chill a cryoprobe to temperatures below -75°C, which when inserted into a lesion, freezes and kills the tissue. It has been used to treat Morton neuroma, other chronic nerve pain syndromes, and conditions for which RFA has been used.^{21,22,23,24,25,26,27,28}

This review primarily focuses on evidence for the use of intralesional alcohol injection, RFA, and cryoablation on painful neuromas, with emphasis on Morton neuroma and the comparative effectiveness of these less invasive therapies with open surgical resection of the nerve pathology.

Summary

Description

Morton neuroma is a common and painful compression neuropathy of the dorsal foot that is also referred to as intermetatarsal neuroma, interdigital neuroma, interdigital neuritis, and Morton metatarsalgia. Morton neuroma has been treated with conservative measures (pads, orthotics, drugs) or surgery. Minimally invasive procedures, including intralesional alcohol injection, radiofrequency ablation (RFA) and cryoablation, have been investigated as alternatives to open surgery. These methods have also been used to treat other peripheral neuromas.

Summary of Evidence

For individuals who have Morton neuroma who receive intralesional alcohol injection(s), the evidence includes retrospective case series. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. The body of evidence is limited, consisting of case series reporting on the treatment response of patients with refractory Morton neuroma. The available case series have generally reported that some patients experience pain relief and express satisfaction with the procedure. Some evidence has suggested that surgery after failed cases of alcohol injections is more complex and challenging than in untreated patients due to the presence of fibrosis. There is a lack of controlled trials comparing alcohol injections with alternative therapies, and there are no controlled studies comparing outcomes for alcohol injections with those for surgery in surgical candidates. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have Morton neuroma who receive RFA, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. Four case series identified reported outcomes for radiofrequency ablation (RFA) to treat Morton neuroma. The body of evidence is highly heterogeneous regarding RFA protocols, descriptions of prior conservative management, patient characteristics, follow-up durations, outcome measures, and reporting of outcomes. Variable proportions of patients require surgery after RFA, making the benefit of RFA for avoiding more invasive treatment uncertain. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have Morton neuroma who receive cryoablation, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. Only 2 retrospective case series on the use of cryoablation to treat peripheral nerve pain were identified in a literature review. The case series were heterogeneous regarding cryoablation protocols and length of follow-up. Outcome measures did not provide information on functional endpoints. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have peripheral neuroma(s) other than Morton neuroma who receive ablation no published literature was identified. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Policy History

Date	Action
8/2023	Annual policy review. Policy statement unchanged.
1/2023	Medicare information removed. See MP #132 Medicare Advantage Management for local coverage determination and national coverage determination reference.
8/2022	Annual policy review. Ongoing intralesional alcohol ablation added to investigational policy statement. Transferred from MP #642 Alcohol Injections for Treatment of Peripheral Morton Neuromas. Policy title changed to "Minimally Invasive Ablation Procedures for Morton and Other Peripheral Neuromas."
8/2021	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
8/2020	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
8/2019	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
7/2018	Annual policy review. Description, summary, and references updated. Policy statements unchanged.
7/2017	Annual policy review. New references added.
5/2015	New medical policy describing investigational indications. Effective 5/1/2015.

Information Pertaining to All Blue Cross Blue Shield Medical Policies

Click on any of the following terms to access the relevant information:

[Medical Policy Terms of Use](#)

[Managed Care Guidelines](#)

[Indemnity/PPO Guidelines](#)

[Clinical Exception Process](#)

[Medical Technology Assessment Guidelines](#)

References

1. Rajput K, Reddy S, Shankar H. Painful neuromas. Clin J Pain. Sep 2012; 28(7): 639-45. PMID 22699131
2. Jain S, Mannan K. The diagnosis and management of Morton's neuroma: a literature review. Foot Ankle Spec. Aug 2013; 6(4): 307-17. PMID 23811947

3. Thomas JL, Blitch EL, Chaney DM, et al. Diagnosis and treatment of forefoot disorders. Section 3. Morton's intermetatarsal neuroma. *J Foot Ankle Surg.* 2009; 48(2): 251-6. PMID 19232980
4. Wu KK. Morton's interdigital neuroma: a clinical review of its etiology, treatment, and results. *J Foot Ankle Surg.* 1996; 35(2): 112-9; discussion 187-8. PMID 8722878
5. MULDER JD. The causative mechanism in morton's metatarsalgia. *J Bone Joint Surg Br.* Feb 1951; 33-B(1): 94-5. PMID 14814167
6. Adams WR. Morton's neuroma. *Clin Podiatr Med Surg.* Oct 2010; 27(4): 535-45. PMID 20934103
7. Thomson CE, Gibson JN, Martin D. Interventions for the treatment of Morton's neuroma. *Cochrane Database Syst Rev.* 2004; 2004(3): CD003118. PMID 15266472
8. Bennett GL, Graham CE, Mauldin DM. Morton's interdigital neuroma: a comprehensive treatment protocol. *Foot Ankle Int.* Dec 1995; 16(12): 760-3. PMID 8749346
9. Dierselhuis EF, van den Eerden PJ, Hoekstra HJ, et al. Radiofrequency ablation in the treatment of cartilaginous lesions in the long bones: results of a pilot study. *Bone Joint J.* Nov 2014; 96-B(11): 1540-5. PMID 25371471
10. Boersma D, van Eekeren RR, Kelder HJ, et al. Mechanochemical endovenous ablation versus radiofrequency ablation in the treatment of primary small saphenous vein insufficiency (MESSI trial): study protocol for a randomized controlled trial. *Trials.* Oct 29 2014; 15: 421. PMID 25354769
11. Di Costanzo GG, Tortora R, D'Adamo G, et al. Radiofrequency ablation versus laser ablation for the treatment of small hepatocellular carcinoma in cirrhosis: a randomized trial. *J Gastroenterol Hepatol.* Mar 2015; 30(3): 559-65. PMID 25251043
12. Anchala PR, Irving WD, Hillen TJ, et al. Treatment of metastatic spinal lesions with a navigational bipolar radiofrequency ablation device: a multicenter retrospective study. *Pain Physician.* 2014; 17(4): 317-27. PMID 25054391
13. Hillen TJ, Anchala P, Friedman MV, et al. Treatment of metastatic posterior vertebral body osseous tumors by using a targeted bipolar radiofrequency ablation device: technical note. *Radiology.* Oct 2014; 273(1): 261-7. PMID 24927327
14. Wang X, Wang X, Song Y, et al. Efficiency of radiofrequency ablation for surgical treatment of chronic atrial fibrillation in rheumatic valvular disease. *Int J Cardiol.* Jul 01 2014; 174(3): 497-502. PMID 24820759
15. Huang WZ, Wu YM, Ye HY, et al. Comparison of the outcomes of monopolar and bipolar radiofrequency ablation in surgical treatment of atrial fibrillation. *Chin Med Sci J.* Mar 2014; 29(1): 28-32. PMID 24698675
16. Avery J, Kumar K, Thakur V, et al. Radiofrequency ablation as first-line treatment of varicose veins. *Am Surg.* Mar 2014; 80(3): 231-5. PMID 24666862
17. Hiraki T, Gobara H, Iguchi T, et al. Radiofrequency ablation as treatment for pulmonary metastasis of colorectal cancer. *World J Gastroenterol.* Jan 28 2014; 20(4): 988-96. PMID 24574771
18. Morillo CA, Verma A, Connolly SJ, et al. Radiofrequency ablation vs antiarrhythmic drugs as first-line treatment of paroxysmal atrial fibrillation (RAAFT-2): a randomized trial. *JAMA.* Feb 19 2014; 311(7): 692-700. PMID 24549549
19. Fuller CW, Nguyen SA, Lohia S, et al. Radiofrequency ablation for treatment of benign thyroid nodules: systematic review. *Laryngoscope.* Jan 2014; 124(1): 346-53. PMID 24122763
20. Huang XM, Hu JQ, Li ZF, et al. Symptomatic sinus tachycardia with perpetuating slow pathway: successful treatment with radiofrequency ablation. *Pacing Clin Electrophysiol.* Oct 2014; 37(10): e1-4. PMID 21077914
21. Prologo JD, Passalacqua M, Patel I, et al. Image-guided cryoablation for the treatment of painful musculoskeletal metastatic disease: a single-center experience. *Skeletal Radiol.* Nov 2014; 43(11): 1551-9. PMID 24972918
22. Kim EH, Tanagho YS, Saad NE, et al. Comparison of laparoscopic and percutaneous cryoablation for treatment of renal masses. *Urology.* May 2014; 83(5): 1081-7. PMID 24560975
23. Durand M, Barret E, Galiano M, et al. Focal cryoablation: a treatment option for unilateral low-risk prostate cancer. *BJU Int.* Jan 2014; 113(1): 56-64. PMID 24053685
24. Duarte R, Pereira T, Pinto P, et al. [Percutaneous Image-guided cryoablation for localized bone plasmacytoma treatment]. *Radiologia.* 2014; 56(5): e1-4. PMID 22621822
25. Rodriguez-Entem FJ, Expósito V, Gonzalez-Enriquez S, et al. Cryoablation versus radiofrequency ablation for the treatment of atrioventricular nodal reentrant tachycardia: results of a prospective randomized study. *J Interv Card Electrophysiol.* Jan 2013; 36(1): 41-5; discussion 45. PMID 23080326

26. Yamauchi Y, Izumi Y, Hashimoto K, et al. Percutaneous cryoablation for the treatment of medically inoperable stage I non-small cell lung cancer. *PLoS One*. 2012; 7(3): e33223. PMID 22413004
27. Collins KK, Schaffer MS. Use of cryoablation for treatment of tachyarrhythmias in 2010: survey of current practices of pediatric electrophysiologists. *Pacing Clin Electrophysiol*. Mar 2011; 34(3): 304-8. PMID 21077912
28. Kaufman CS, Bachman B, Littrup PJ, et al. Cryoablation treatment of benign breast lesions with 12-month follow-up. *Am J Surg*. Oct 2004; 188(4): 340-8. PMID 15474424
29. Pasquali C, Vulcano E, Novario R, et al. Ultrasound-guided alcohol injection for Morton's neuroma. *Foot Ankle Int*. Jan 2015; 36(1): 55-9. PMID 25367249
30. Perini L, Perini C, Tagliapietra M, et al. Percutaneous alcohol injection under sonographic guidance in Morton's neuroma: follow-up in 220 treated lesions. *Radiol Med*. Jul 2016; 121(7): 597-604. PMID 26883232
31. Musson RE, Sawhney JS, Lamb L, et al. Ultrasound guided alcohol ablation of Morton's neuroma. *Foot Ankle Int*. Mar 2012; 33(3): 196-201. PMID 22734280
32. Hughes RJ, Ali K, Jones H, et al. Treatment of Morton's neuroma with alcohol injection under sonographic guidance: follow-up of 101 cases. *AJR Am J Roentgenol*. Jun 2007; 188(6): 1535-9. PMID 17515373
33. Fanucci E, Masala S, Fabiano S, et al. Treatment of intermetatarsal Morton's neuroma with alcohol injection under US guide: 10-month follow-up. *Eur Radiol*. Mar 2004; 14(3): 514-8. PMID 14531002
34. Morgan P, Monaghan W, Richards S. A systematic review of ultrasound-guided and non-ultrasound-guided therapeutic injections to treat Morton's neuroma. *J Am Podiatr Med Assoc*. Jul 2014; 104(4): 337-48. PMID 25076076
35. Dockery GL. The treatment of intermetatarsal neuromas with 4% alcohol sclerosing injections. *J Foot Ankle Surg*. 1999; 38(6): 403-8. PMID 10614611
36. Genon MP, Chin TY, Bedi HS, et al. Radio-frequency ablation for the treatment of Morton's neuroma. *ANZ J Surg*. Sep 2010; 80(9): 583-5. PMID 20857612
37. Moore JL, Rosen R, Cohen J, et al. Radiofrequency thermoneurolysis for the treatment of Morton's neuroma. *J Foot Ankle Surg*. 2012; 51(1): 20-2. PMID 22055491
38. Chuter GS, Chua YP, Connell DA, et al. Ultrasound-guided radiofrequency ablation in the management of interdigital (Morton's) neuroma. *Skeletal Radiol*. Jan 2013; 42(1): 107-11. PMID 23073898
39. Connors JC, Boike AM, Rao N, et al. Radiofrequency Ablation for the Treatment of Painful Neuroma. *J Foot Ankle Surg*. 2020; 59(3): 457-461. PMID 32354501
40. Friedman T, Richman D, Adler R. Sonographically guided cryoneurolysis: preliminary experience and clinical outcomes. *J Ultrasound Med*. Dec 2012; 31(12): 2025-34. PMID 23197557
41. Cazzato RL, Garnon J, Ramamurthy N, et al. Percutaneous MR-Guided Cryoablation of Morton's Neuroma: Rationale and Technical Details After the First 20 Patients. *Cardiovasc Intervent Radiol*. Oct 2016; 39(10): 1491-8. PMID 27189181
42. The Association of Extremity Nerve Surgeons Clinical Practice Guidelines v. 2.0. 2020. Accessed May 8, 2023.