

# CAR T-Cell Therapy Services for Non-Hodgkin Lymphoma (Lisocabtagene Maraleucel) Prior Authorization Request Form #941

# Medical Policy #066 Chimeric Antigen Receptor Therapy for Leukemia and Lymphoma

### **CLINICAL DOCUMENTATION**

- Clinical documentation that supports the medical necessity criteria for CAR T-Cell Therapy Services for Non-Hodgkin Lymphoma (Lisocabtagene Maraleucel) must be submitted.
- If the patient does not meet all the criteria listed below, please submit a letter of medical necessity with a request for <u>Clinical Exception (Individual Consideration)</u> explaining why an exception is justified.

#### **Requesting Prior Authorization Using Authorization Manager**

Providers will need to use <u>Authorization Manager</u> to submit initial authorization requests for services. Authorization Manager, available 24/7, is the quickest way to review authorization requirements, request authorizations, submit clinical documentation, check existing case status, and view/print the decision letter. For commercial members, the requests must meet medical policy guidelines.

To ensure the request is processed accurately and quickly:

- Enter the facility's NPI or provider ID for where services are being performed.
- Enter the appropriate surgeon's NPI or provider ID as the servicing provider, *not* the billing group.

#### **Authorization Manager Resources**

• Refer to our <u>Authorization Manager</u> page for tips, guides, and video demonstrations.

Complete Prior Authorization Request Form for CAR T-Cell Therapy Services for Non-Hodgkin Lymphoma (Lisocabtagene Maraleucel) (941) using <u>Authorization Manager</u>

For out of network providers: Requests should still be faxed to 888-973-0726.

Patient Information		
Patient Name:	Today's Date:	
BCBSMA ID#:	Date of Treatment:	
Date of Birth:	Place of Service: Outpatient  Inpatient	

Physician Information	Facility Information
Name:	Name:
Address:	Address:
Phone #:	Phone #:
Fax#:	Fax#:
NPI#:	NPI#:

#### Please check off if the patient is enrolled in a Clinical Trial.

Please check off if the patient has the following diagnosis and <u>HAS RELAPSED</u> <sup>c</sup> or is <u>REFRACTORY<sup>c</sup></u> :	
Histologically confirmed diagnosis of:	
Diffuse large B-cell lymphoma not otherwise specified (including diffuse large B-cell lymphoma arising from indolent lymphoma)	
High-grade B-cell lymphoma, <b>OR</b>	
Primary mediastinal large B-cell lymphoma, <b>OR</b>	
Follicular lymphoma grade 3B.	

<sup>c</sup> Relapsed or refractory disease is defined as progression after 2 or more lines of systemic therapy (which may or may not include therapy supported by autologous cell transplant).

Pleas	e check off that the patient meets <u>ALL</u> the following criteria:	
0	listologically confirmed diagnosis of large B-cell lymphoma, including diffuse large B-cell lymphoma not therwise specified (including diffuse large B-cell lymphoma arising from indolent lymphoma), high-grade B- ell lymphoma, primary mediastinal large B-cell lymphoma, and follicular lymphoma grade 3B; <b>AND</b>	
2.	Meets at least one of the following:	
	<ul> <li>Primary refractory or relapsed disease<sup>c</sup> within 12 months of first-line chemo-immunotherapy that included an anti-CD20 monoclonal antibody and anthracycline-containing regimen; OR</li> </ul>	
	<ul> <li>Primary refractory or relapsed disease within 12 months of first-line chemo-immunotherapy that included an anti-CD20 monoclonal antibody and anthracycline-containing regimen and are not eligible for hematopoietic stem cell transplantation due to comorbidities or age; OR</li> </ul>	
	<ul> <li>Relapsed or refractory disease as defined as progression after ≥2 lines of systemic therapy including anti-CD20 monoclonal antibody for CD20-positive tumor and anthracycline-containing chemotherapy</li> </ul>	
	i. When the individual has histological transformation of follicular lymphoma or marginal zone lymphoma to diffuse large B-cell lymphoma: prior chemotherapy for follicular lymphoma or marginal zone lymphoma and ≥2 chemo-immunotherapy regimens for the transformed disease; AND	
3.	At least 18 years of age at the time of infusion; AND	
4.	Have adequate organ and bone marrow function as determined by the treating oncologist/hematologist; <b>AND</b>	
5.	Have not received prior CD19-directed chimeric antigen receptor T-cell therapy treatment, any other cell therapy, or any gene therapy or are being considered for treatment with any other cell therapy or any gene therapy; <b>AND</b>	
6.	Do not have primary central nervous system lymphoma.	

<sup>c</sup> Relapsed or refractory disease is defined as progression after 2 or more lines of systemic therapy (which may or may not include therapy supported by autologous cell transplant).

### CPT CODES/ HCPCS CODES/ ICD CODES

HCPCS codes:	Code Description	
C9076	Lisocabtagene maraleucel, up to 110 million autologous anti-cd19 car-positive viable t cells, including leukapheresis and dose preparation procedures, per therapeutic dose	
C9399	Unclassified drugs or biologicals	
J3490	Unclassified drugs	
J3590	Unclassified biologics	

J9999	Not otherwise classified, antineoplastic drugs	
-------	--	--

## Providers should enter the <u>relevant diagnosis code(s)</u> below:

Code	Description	

## Providers should enter <u>other relevant code(s)</u> below:

Code	Description	